

# Exhibit A

CFS Comments on Supplemental Environmental Impact Statement for Control of Burrowing Shrimp using Imidacloprid on Commercial Oyster and Clam Beds in Willapa Bay and Grays Harbor, Washington



**Catalogue number: H113-27/2016-20E-PDF**

**Pest Management Regulatory Agency (PMRA)  
Health Canada  
2720 Riverside Drive  
Ottawa, Ontario**

**March 22, 2017**

**Comments to Health Canada Pest Management Regulatory Agency on Proposed  
Re-evaluation Decision PRVD2016-20, Imidacloprid**

The Center for Food Safety (CFS)<sup>1</sup> in conjunction with the Save the Oak Ridges Moraine Coalition (STORM)<sup>2</sup> and Avaaz<sup>3</sup> submit the following comment in support of Health Canada Pest Management Regulatory Agency's (PMRA) proposal to prohibit the use of imidacloprid and further encourages PMRA to expedite the proposed phase out process.

We are writing on behalf of over 4.4 million people worldwide, including 200,474 Canadians, who have signed an Avaaz petition urging Canada and others to "immediately ban the use of neonicotinoid pesticides" particularly imidacloprid. Additionally CFS has 5,275 Canadian members who are increasingly concerned about the impacts that pesticides, especially neonicotinoids, are having on pollinators and our environment. Given the weight of evidence showing detrimental impacts of imidacloprid to aquatic ecosystems and the comprehensive evaluation of possible mitigation strategies, we encourage PMRA to accelerate its proposed phase out of this agent for agricultural and outdoors uses.

The Pest Control Products Act makes clear that Health Canada is responsible for regulating the use of pesticides "to protect human health and safety and the environment." Thousands of Canadians have also submitted comments to this consultation, through Avaaz, reaffirming their expectation that PMRA fulfill this responsibility by acting swiftly to prohibit the use of imidacloprid.

---

<sup>1</sup> CFS is a nonprofit, membership organization with a mission to empower people, support farmers, and protect the earth from the harmful impacts of industrial agriculture. Through groundbreaking legal, scientific, and grassroots action, CFS protects and promotes the public's right to safe food and the environment. CFS has more than 830,000 consumer and farmer supporters—including 5,275 Canadian members.

<sup>2</sup> The STORM Coalition is focused on protecting the ecological integrity of the Oak Ridges Moraine. Since 1989, STORM has been working to ensure that local and regional governments' planning decisions respect the environmental significance of the moraine and take into account its ecological and hydrological functions.

<sup>3</sup> Avaaz is a 44-million-person global campaign network that works to ensure that the views and values of the world's people shape global decision-making. "Avaaz" means "voice" or "song" in many languages. Avaaz members live in every nation of the world.

Sonya V. Thursby of Toronto, an Avaaz member, wrote:

*Demonstrate leadership by banning Imidacloprid! We know how critical a balanced ecosystem is to sustain life—the harm this chemical causes to aquatic insects throws off this significant balance. The phase out period needs to be shortened to now—the time to act is now because we have the knowledge to understand how harmful this chemical is to our ecosystems and biodiversity. What state will our ecosystem be in when my 13 and 17 year old children reach adulthood if leaders like you act now.*

**This submission asserts that:**

- a. The proposed three - five year phase out period is unnecessarily long, given the data presented by PMRA, and the weight of scientific evidence demonstrating imidacloprid’s harm to wildlife, including aquatic insects, terrestrial organisms, birds and pollinators.
- b. Neither alternative use reduction plans, nor precautionary label statements will adequately or reliably reduce the risks posed to the environment by imidacloprid.
- c. PMRA must also act to review and phase out other neonicotinoids to ensure that the environmental benefits of imidacloprid’s prohibition are not offset by increases in the use of other similarly harmful agents.

**Expedite the Phase Out: Imdacloprid is Found at Unsafe Levels Known to Cause Harm To a Variety of Species**

A growing number of studies show that Canadian waters are in jeopardy from continued contamination by neonicotinoid insecticides used widely for agricultural and outdoor uses. In fact a three-year investigation of neonicotinoid insecticide contamination in surface water sites across southern Ontario revealed three of the five neonicotinoids tested (imidacloprid, clothianidin, thiamethoxam), had more than 90 percent detection rates in over half of the sites.<sup>i</sup> The Canadian government’s threshold for imidacloprid residues in freshwater is .23 ppb, which was exceeded in 75 percent of the samples collected in two sampling sites.<sup>ii</sup> The data from the three-year investigation show a strong correlation between pesticide detection, precipitation, and stream discharge. Other studies of Canadian monitoring data by government and independent researchers revealed 98.7 percent detection frequency of thiamethoxam and 100 percent detection frequency of clothianidin in Southwestern Ontario water samples from corn-producing counties<sup>iii</sup>, and 91 percent neonicotinoid detection (imidacloprid, thiamethoxam, clothianidin, acetamiprid) in wetlands sampled across the Prairie Pothole region.<sup>iv</sup> Health Canada reported imidacloprid at concentrations as high as 290 times greater than the level of acceptable risk.<sup>v</sup> Across all studies, researchers noted neonicotinoids long- term persistence and highlighted specific concerns for wetlands in colder climates where the chemicals persist in soil and transport via snowmelt to nearby surface water.<sup>vi</sup>

A prohibition will only be implemented after the publication of the government’s final re-evaluation decision. If PMRA’s suggested three - five year phase-out period is adopted then imidacloprid will continue to be used until 2020 or 2022, depending on the re-evaluation decision’s final publication date. This is particularly troubling given the above noted findings, imidacloprid’s half-life of up to 229 days in soils,<sup>vii</sup> and PMRA’s own conclusion that “under current conditions of use, the environmental risks for most products containing imidacloprid do not meet

current safety standards.” France has committed to prohibit all neonicotinoid chemicals as early as 2018 and we strongly encourage the Canadian government to match these target dates.

### **Imidacloprid Impacts More Than Just Aquatic Insects**

PMRA correctly identifies that imidacloprid places aquatic insects at risk; however, as the agency progresses with its final re-evaluation, it must also thoroughly assess risks to other species, including pollinators, birds, and beneficial insects (such as earthworms).

For earthworms, Wang et al. 2015 note a LC<sub>50</sub> of 3.05 mg/kg of imidacloprid and that a sub-lethal dose of 2.0 mg/kg, caused an 84 percent decrease in fecundity.<sup>viii</sup> Because earthworms are critical to soil health, we feel it is imperative that harms to these beneficial insects are thoroughly evaluated in current and future risk assessments.

Hallman et al. 2014 determined that commonly-found levels of imidacloprid in Holland’s surface water correlates to a 3.5 percent annual decline in bird populations.<sup>ix</sup> Gibbons et al. 2015 found, in a comprehensive review of 150 studies, that ingestion of even a few neonicotinoid-coated seeds could cause mortality or reproductive impairment to sensitive bird species.<sup>x</sup> While, PMRA acknowledges that coated seeds may be harmful for birds, it’s suggested mitigation strategies (including label cautions and removing coated seeds from field surfaces) are inadequate when considering the minimal exposure required to cause harm.

### **Other Jurisdictions**

The U.S. Environmental Protection Agency (EPA) made similar findings about aquatic ecosystems in the 2017 *Preliminary Aquatic Risk Assessment to Support the Registration Review of Imidacloprid*. The comprehensive report incorporated data from PMRA as well as the European Food Safety Authority. Based on the evidence compiled, the EPA underlined the threat of imidacloprid to aquatic communities concluding, “It is evident ...that concentrations of imidacloprid detected in streams, rivers, lakes and drainage canals routinely exceed acute and chronic toxicity endpoints derived for freshwater invertebrates.” and explained that acute and chronic risks were identified with a majority of registered uses of imidacloprid.<sup>xi</sup> Because the farming practices and water bodies in the nations are comparable, these conclusions bolster those of PMRA and further emphasize the need for strong regulatory action.

### **Alternative Use Reduction Plans and Precautionary Labels Will Not Sufficiently Reduce Risks**

We also want to support and reiterate PMRA’s caution against an alternative use reduction plan. As outlined in the assessment, any possible mitigation plans are hindered by an inability to accurately predict the degree of reduction necessary to meet acceptable imidacloprid levels without extensive and costly water monitoring information. Further any plans are limited by difficulties in specifying the uses causing high concentration levels and finding means of ensuring that approved imidacloprid use does not increase in areas in which it is not used

extensively now. Finally, PMRA has identified immediate contamination concerns and there is no way to gauge how long an effective reduction plan would take.

The PMRA's Proposed Re-Evaluation Decision explains that water based imidacloprid is found at unsafe levels at agricultural sites, even though imidacloprid products have precautionary labels designed specifically to minimize this risk. For this reason, we again urge the agency to move forward with an expedited phase-out of imidacloprid, as current best management practices and pesticide product labels are insufficient for protecting aquatic species, birds, pollinators, and other beneficial insects.

### **Strong Actions Are Also Needed For Clothianidin and Thiamethoxam**

While imidacloprid poses a significant threat to the environment, so to, do other neonicotinoid insecticides, such as clothianidin and thiamethoxam. We strongly encourage PMRA to take immediate action to curtail the use of all neonicotinoids, so as to ensure that the environmental benefits associated with an imidacloprid phase out are not suppressed by the increased use of similarly detrimental alternatives.

A 2017 report from Chretien et al. of Agriculture and Agri-Food Canada, as well as Quebec Ministry of Sustainable Development, raises concerns about contamination from surface runoff and subsurface tile drain losses with a particular focus on the contamination by clothianidin and thiamethoxam. The report documents a two year study in which 14 surface runoff and tile drain discharge events were monitored. The researchers reported, "detection frequencies close to 100 percent in edge-of-field, surface runoff and tile drain water samples...for thiamethoxam and clothianidin even though only thiamethoxam had been applied in the first year."<sup>xii</sup> These findings highlight the persistent nature of these chemicals in certain climates and soil conditions as well as the potential harm of their degradates. The insecticides were reported at median concentrations of .46 ppb and .16 ppb and many concentrations exceeded the .0083 ppb chronic threshold for effect on aquatic life recommended by Government of Quebec.<sup>xiii</sup> The authors of the report echoed the proposal established in the Quebec Pesticide Strategy 2015-2018, and explained that any plans for reduced use or mitigation to control dust and surface runoff would not be sufficient.

Despite these findings, with a majority of environmental monitoring programs and toxicity testing dedicated to imidacloprid use, "no ecological thresholds exist for thiamethoxam and clothianidin."<sup>xiv</sup> This major shortcoming is particularly an issue in Quebec where nearly 100 percent of corn and 50 percent of soybean seeds are planted with neonicotinoid seed coatings—covering nearly 500,000 ha.<sup>xv</sup> Giroux et al. found detection frequencies of thiamethoxam and clothianidin ranging from 93 percent to 98 percent from 2012 to 2014 in four Quebec watersheds.<sup>xvi</sup> Canada's increasing documentation of this contamination supports PMRA's analysis and is an indication of the critical need for setting stronger regulatory protections for the environment, including the proposed prohibition of imidacloprid, as well as other neonicotinoid insecticides.

## Conclusions and Recommendations

Perhaps the most important point in PMRA's own conclusion was the realization that imidacloprid is indeed causing harm to aquatic environments and current uses are "not sustainable"<sup>xvii</sup>. CFS, Avaaz, and STORM fully agree with PMRA's strong conclusion - it is evident that a complete phase out is a necessary action to protect aquatic ecosystems from imidacloprid contamination. Furthermore, we support PMRA's call for similar evaluations to be conducted for additional neonicotinoid insecticides, particularly thiamethoxam and clothianidin. We commend PMRA for its decision to move forward with strong regulations based on the chemical's water solubility, persistence, and propensity for unintended contamination of critical waterbodies.

We would further like to highlight the analysis and recommendations in the two attached CFS reports, which are incorporated into this comment by reference: *2017 Updates to Water Hazard: Aquatic Contamination by Neonicotinoid Insecticides in the United States* and *Water Hazard: Aquatic Contamination by Neonicotinoid Insecticides in the United States*. Also incorporated are the attached testimonies of Avaaz members across Canada concerned with the continued use of neonicotinoid insecticides.

Joe Thauberger of Canada, an Avaaz member, wrote:

*To the PMRA, I am a retired farmer from Saskatchewan. I have seen with my own eyes the devastating affects on our wildlife. Insecticides of all kinds should be banned and used only under very special situations.*

Due to the reasons above, CFS, Avaaz, and STORM urge Health Canada to implement the proposed phase out of imidacloprid for agricultural and outdoor uses and to conduct similar evaluations of other neonicotinoid insecticides in order to put an end to the widespread contamination, the increasing concentration rates, and the observed decline in vital aquatic species. Finally, we urge you to consider shortening the phase out period to the shortest time feasible in view of the imminent hazards posed.

Thank you for your thoughtful consideration of this important matter.

Sincerely,

Center for Food Safety

Avaaz

Save the Oak Ridges Moraine Coalition

**For further information contact:**

**Larissa Walker**

**Pollinator Program Director | Policy Analyst**

**Center for Food Safety**

**660 Pennsylvania Ave. SE, Suite 302**

**Washington, DC 20003**

**(P): 202.547.9359 | (E): LWalker@CenterForFoodSafety.org**

- 
- <sup>i</sup> Struger, J., Grabuski, J., Cagampan, S., Sverko, E., McGoldrick, D., and Marvin, C. H. (2017). Factors influencing the occurrence and distribution of neonicotinoid insecticides in surface waters of southern Ontario, Canada. *Chemosphere*, 169, 516-523.
- <sup>ii</sup> Canadian Council of Ministers of the Environment. (2007). Canadian water quality guidelines for the protection of aquatic life: Imidacloprid. In: Canadian Environmental Quality Guidelines. Winnipeg, Manitoba, Canada.; Struger, J., Grabuski, J., Cagampan, S., Sverko, E., McGoldrick, D., and Marvin, C. H. (2017). Factors influencing the occurrence and distribution of neonicotinoid insecticides in surface waters of southern Ontario, Canada. *Chemosphere*, 169, 516-523.
- <sup>iii</sup> Schaafsma, A., Limay-Rios, V., Baute, T., Smith, J., and Xue, Y. (2015). Neonicotinoid insecticide residues in surface water and soil associated with commercial maize (corn) fields in southwestern Ontario. *PLoS One*, 10(2), e0118139.
- <sup>iv</sup> Main, A. R., Michel, N. L., Headley, J. V., Peru, K. M., and Morrissey, C. A. (2015). Ecological and landscape drivers of neonicotinoid insecticide detections and concentrations in Canada's prairie wetlands. *Environmental Science and Technology*, 49(14), 8367-8376.
- <sup>v</sup> Jane Philpott on Agriculture and Agri-Food (2016, November 24). Retrieved from <https://openparliament.ca/debates/2016/11/24/jane-philpott-2/only/>
- <sup>vi</sup> Main, A. R., Michel, N. L., Headley, J. V., Peru, K. M., and Morrissey, C. A. (2015). Ecological and landscape drivers of neonicotinoid insecticide detections and concentrations in Canada's prairie wetlands. *Environmental Science and Technology*, 49(14), 8367-8376.
- <sup>vii</sup> Bonmatin, J. M., Giorio, C., Girolami, V., Goulson, D., Kreuzweiser, D. P., Krupke, C., ... and Noome, D. A. (2015). Environmental fate and exposure; neonicotinoids and fipronil. *Environmental Science and Pollution Research*, 22(1), 35-67.
- <sup>viii</sup> Wang K., Pang, S., Mu, X., Qi, S., Li, D., Cui, F., and Wang, C. (2015). Biological response of earthworm, *Eisenia fetida*, to five neonicotinoid insecticides. *Chemosphere*, 132, 233-240.
- <sup>ix</sup> Hallmann, C. A., Foppen, R. P., van Turnhout, C. A., de Kroon, H., and Jongejans, E. (2014). Declines in insectivorous birds are associated with high neonicotinoid concentrations. *Nature*.
- <sup>x</sup> Gibbon, D., Morrissey, C., Mineau, P. (2015). A review of the direct and indirect effects of neonicotinoids and fipronil on vertebrate wildlife. *Environmental Science and Pollution Research*, 22(1), 103-118.
- <sup>xi</sup> Sappington, K. G., Ruhman, M. A., Housenger, J. (2016, December 22). Preliminary Aquatic Risk Assessment to Support the Registration Review of Imidacloprid. Washington: US Environmental Protection Agency Memorandum. Retrieved from: <https://www.regulations.gov/document?D=EPA-HQ-OPP-2008-0844-1086>
- <sup>xii</sup> Chrétien, F., Giroux, I., Thériault, G., Gagnon, P., and Corriveau, J. (2017). Surface runoff and subsurface tile drain losses of neonicotinoids and companion herbicides at edge-of-field. *Environmental Pollution*.
- <sup>xiii</sup> Chrétien, F., Giroux, I., Thériault, G., Gagnon, P., and Corriveau, J. (2017). Surface runoff and subsurface tile drain losses of neonicotinoids and companion herbicides at edge-of-field. *Environmental Pollution*.
- <sup>xiv</sup> Chrétien, F., Giroux, I., Thériault, G., Gagnon, P., and Corriveau, J. (2017). Surface runoff and subsurface tile drain losses of neonicotinoids and companion herbicides at edge-of-field. *Environmental Pollution*.
- <sup>xv</sup> Ministère du Développement durable, de l'Environnement et de la Lutte contre les changements climatiques, Québec. (2015 November). Quebec pesticides strategy 2015-2018. *MDDELCC*. Retrieved from: <http://www.mddelcc.gouv.qc.ca/pesticides/strategie2015-2018/index-en.htm>
- <sup>xvi</sup> Giroux, I. (2015). Présence de pesticides dans l'eau au Québec-Portrait et tendances dans les zones de maïs et de soya 2011 à 2014. *Ministère du Développement durable, de l'Environnement et de la Lutte contre les changements climatiques, Québec*, 5.
- <sup>xvii</sup> Pest Management Regulatory Agency. (2016). Proposed Re-evaluation Decision PRVD2016-20, Imidacloprid. *Consumer product safety*. Retrieved from: [http://www.hc-sc.gc.ca/cps-spc/pest/part/consultations/\\_prvd2016-20/prvd2016-20-eng.php#s3](http://www.hc-sc.gc.ca/cps-spc/pest/part/consultations/_prvd2016-20/prvd2016-20-eng.php#s3)



July 24, 2017

**Comments from Center for Food Safety on the  
EPA's Preliminary Aquatic Risk Assessment to Support the  
Registration Review of Imidacloprid, dated December 22, 2016**

Imidacloprid Docket ID: EPA-HQ-OPP-2008-0844  
Imidacloprid Document ID: EPA-HQ-OPP-2008-0844-1086

The Center for Food Safety (CFS) is a nonprofit, membership organization with a mission to empower people, support farmers, and protect the earth from the harmful impacts of industrial agriculture. Through groundbreaking legal, scientific, and grassroots action, CFS protects and promotes the public's right to safe food and the environment. CFS has more than 900,000 consumer and farmer supporters across the United States. We are pleased to submit these comments on the Preliminary Aquatic Risk Assessment to Support the Registration Review of Imidacloprid (PARA).

**Unacceptable Delays in the Registration Review Process**

Imidacloprid's Registration Review process is far behind the schedule to which the agency formally committed. The "Preliminary Work Plan" for this Registration Review, issued in 2008, had a "2014– Jul-Sep" completion date.<sup>1</sup> It also had this statement (emphasis added): "After reviewing and responding to comments and data received in the docket during this initial comment period, the Agency will develop and **commit** to a final work plan and schedule for the registration review of imidacloprid." The current "Final Work Plan" was issued in 2010.<sup>2</sup> It has this statement in the schedule: "Final Decision and Begin Post-Decision Follow-up - 2016– Jan-Mar." The agency has failed to comply with its own commitment, with a likely completion date now at least two years later than scheduled. EPA must expedite completion of this process.

**Noncompliance with the Endangered Species Act**

EPA acknowledges the lack of Endangered Species Act (ESA) analysis or compliance stating (p. 119):

"Given that the agencies are continuing to develop and work toward implementation of the Interim Approaches to assess the potential risks of pesticides to listed species and their designated critical habitat, this ecological problem formulation supporting the Preliminary Work Plan for imidacloprid does not describe the specific ESA analysis, including effects determinations for specific listed species or designated critical habitat, to be conducted during registration review."

---

<sup>1</sup> <https://www.regulations.gov/document?D=EPA-HQ-OPP-2008-0844-0003>.

<sup>2</sup> July 23, 2010. Imidacloprid Amended Final Work Plan; <https://www.regulations.gov/document?D=EPA-HQ-OPP-2008-0844-0121>.



However, EPA's PARA, taken together with an extensive amount of independent science, underscores that the ongoing contamination of aquatic ecosystems with imidacloprid run-off is adversely affecting a large variety of aquatic species – which includes ESA-listed aquatic species. Illustrative examples of ESA-listed aquatic species known to be vulnerable to these harmful effects include, but are not limited to (indeed there are scores of others): Hines emerald dragonfly (*Somatochlora hineana*); Nashville crayfish (*Orconectes shoupi*); Salt Creek tiger beetle (*Cicindela nevadica lincolniiana*); and San Diego fairy shrimp (*Branchinecta sandiegonensis*).

It is essential that EPA act contemporaneously in this Registration Review risk analysis process to also include thorough analyses of foreseeable effects to ESA-listed aquatic species now. Under the ESA implementing regulation, 50 C.F.R. § 402.14(a), agencies must review their actions at the “earliest possible time.” EPA must not delay this ESA-mandated review or else it will be in violation of the law.<sup>3</sup> Referencing alleged changes in the Interim Approaches document is not an excuse for non-compliance or for the extensive delays that have already occurred.

### **Harm to Aquatic Ecosystems and the Broader Environment**

A growing number of studies show that North American waters are in jeopardy from continued contamination by neonicotinoid insecticides used widely for agricultural and outdoor uses. A 2016 U.S. Geological Survey (USGS) review of pesticide detections in streams across the Midwest found high concentrations of imidacloprid in 98% of the sites sampled.<sup>i</sup> Of all the insecticides tested, imidacloprid was detected at the highest concentrations, with numerous detections exceeding levels known to cause harm to aquatic invertebrates.<sup>ii</sup> This USGS review is part of a growing body of research that highlights the alarming levels of contamination exposed in national and regional monitoring data,<sup>iii</sup> and builds on other reported detection frequencies such as: the 76% detection rate of one or more neonicotinoids in streams across the Midwest in 2013,<sup>iv</sup> the 70% detection frequency of downstream samples in the southern Appalachians in 2012 and 2013,<sup>v</sup> and an overall 63% detection rate in streams sampled across the United States.<sup>vi</sup> EPA recognizes this research in the PARA and yet did not conclude that such vast contamination warranted immediate action to restrict uses. This clear failure to take immediate action is particularly concerning given that numerous analyses of peer-reviewed research have shown severe risk to aquatic ecosystems—most notably Sanchez-Bayo et al. 2016, which alarmingly concluded, “Negative impacts of neonicotinoids in aquatic environments are a reality” and continues, “Solutions must be found soon if we are to save the biodiversity not only of aquatic ecosystems, but all other ecosystems linked by the food web.”<sup>vii</sup>

### **Potential Impacts to Human Health**

Furthermore, new research is emerging about the potential public health risks that imidacloprid and other persistent neonicotinoid pesticides pose. A 2017 study from USGS and the University of Iowa, *Occurrence of Neonicotinoid Insecticides in Finished Drink Water and Fate During Drinking Water Treatment*, found imidacloprid, clothianidin, and thiamethoxam in 100% of samples taken from University of Iowa tap water.<sup>viii</sup> The concentrations detected range from 0.00024 ppb to 0.0573 ppb. The report is the first peer reviewed study to examine neonicotinoid concentrations in finished drinking water. Although the study is limited to a small sampling area, the authors of the report conclude, “because of their pervasiveness in source waters, and persistence through treatment systems, neonicotinoids are likely present in other drinking water systems across the United States.” While this study is preliminary and did not expose any concentrations known to have direct impact on humans, a 2015 publication by National

---

<sup>3</sup> The scope of agency actions triggering Section 7 duties is broad, including all activities or programs of any kind authorized, licensed, funded, or carried out by federal agencies, including activities directly or indirectly causing modifications to land, water, or air. 50 C.F.R. § 402.02 (definition of “action”). The potential “effects” of an action that an agency must consider are similarly broad, and include both “direct” and “indirect” effects of the action and all activities “interrelated or interdependent” with that action. Id.

Institute of Health called for further research on the chronic human health impacts of neonicotinoids.<sup>ix</sup> Since there are currently no standards for neonicotinoids in drinking water in the United States, CFS encourages EPA to consider this route of exposure as a potential threat to human health and immediately conduct a full array of safety testing. Then, appropriate health-based restrictions on them may be needed.

### **Proposed Action to Phase-Out Uses of Imidacloprid in Canada**

In deciding the fate of the continued use of imidacloprid and other neonicotinoid insecticides, EPA should also consider the actions proposed by Health Canada's Pest Management Regulatory Agency (PMRA). PMRA's 2016 re-evaluation of imidacloprid includes a wealth of data from both government and peer-reviewed research and concludes (emphasis added):

*“The environmental assessment showed that, in aquatic environments in Canada, imidacloprid is being measured at levels that are harmful to aquatic insects. These insects are an important part of the ecosystem, including as a food source for fish, birds and other animals. Based on currently available information, the continued high volume use of imidacloprid in agricultural areas is not sustainable.”<sup>x</sup>*

Based on the documented exceedance of water quality thresholds and aquatic life benchmarks in monitoring data, PMRA proposed action necessary to protect aquatic ecosystems from imidacloprid and called for similar evaluations for other neonicotinoid insecticides. Specifically, PMRA proposed to “phase-out all the agricultural and a majority of other outdoor uses of imidacloprid over three to five years.”<sup>xi</sup> EPA relied on data from the PMRA analysis in its PARA, yet no similar proposals were made to phase-out or even restrict uses of imidacloprid in the U.S. Given that EPA, PMRA, and California Department of Pesticide Regulation have been working together on the neonicotinoid registration reviews, CFS strongly urges EPA to propose similar actions to prevent continued damages to vulnerable ecosystems.

**The following points address additional shortcomings in EPA's PARA. CFS encourages EPA to consider these shortcomings in its final review of imidacloprid:**

#### **1. Gross Underestimation of Seed Treatment Contamination and Risk**

EPA's PARA analysis proposes the unrealistic assumption that neonicotinoid chemicals applied as coatings on seeds planted below two centimeters do not move into surface waters and therefore are low risk.<sup>xii</sup> It is unacceptable that EPA's models do not account for lateral movement of these chemicals in soil and run-off. It is well documented that these chemicals move down into ground water—to assume they don't move laterally through surface soil (especially surface soil broken up by tillage) with precipitation is indefensible in view of numerous published reports showing that they do so.<sup>xiii</sup>

Roughly 1,116,000 pounds of imidacloprid were used on crops in the United States between 2004 and 2013. Fifty-six percent of this usage was as seed coatings—and more specifically 36% was as a coating on soybeans.<sup>xiv</sup> Ninety-four percent of agricultural use scenarios modeled (29 of 31) in the PARA identified acute risks to freshwater species. A majority of use scenarios were seed-coating applications—pointing to the considerable risk from this route of exposure.

The following graphic from the EPA PARA depicts the surface water contamination across the United States in relation to thresholds established for specific freshwater invertebrate species.<sup>xv</sup> As shown, concentration levels of imidacloprid detected in various water bodies are routinely exceeding benchmarks known to cause harm to critical aquatic species (with some

storm event models showing nearly 100% exceedance). **EPA in the final ecological assessment should more accurately portray the harms caused by imidacloprid seed-coatings.**<sup>4</sup>

## 2. New Endpoints but No Mandates to Ensure High Water Quality

After analyzing aquatic toxicity research, international benchmarks, and available monitoring data, and conducting acute lab testing, EPA's PARA proposed new acute and chronic endpoints for imidacloprid for freshwater invertebrates. Prior to the Assessment, EPA's endpoints were exponentially higher than other regulatory and non-regulatory benchmarks from around the world.<sup>xvi</sup> The new proposed endpoints of 0.39 ppb (acute) and 0.01 ppb (chronic) are not only more in line with the conclusions of PMRA, but they also are more consistent with the thresholds proposed by Morrissey et al., and discussed in CFS's 2015 *Water Hazard Report*. Yet, these endpoints have not been updated on EPA's Aquatic Life Benchmarks for Pesticide Registration website.<sup>5</sup> Moreover, there is no mandate by which toxicity benchmarks are enforced. According to its website, EPA's Office of Water may use the "aquatic toxicity data to develop ambient water quality criteria that can be adopted by states and tribes to establish water quality standards under the Clean Water Act,"<sup>xvii</sup> however there are no mandates to establish such standards. Given that current monitoring data shows exceedances of the proposed thresholds across the United States in various surface water bodies, **EPA should formally update proposed water quality standards.**

## 3. No Mention of Pesticide Synergies

EPA's PARA contains almost no mention of pesticide synergies and the particular threat of chemical combinations to aquatic ecosystems unable to escape continued exposure to multiple pesticide stressors. According to Morrissey et al. 2015, "neonicotinoids are known to be additively or synergistically toxic when they occur together or when combined with certain fungicides..."<sup>xviii</sup> These combined "tank mixes" of pesticide formulations are patented and even encouraged by agrichemical companies for their increased toxicity. In fact, a 2016 Center for Biological Diversity analysis of recently approved products from major pesticide companies found that 69% of patent applications claimed or demonstrated synergistic action.<sup>xix</sup> Additionally, when neonicotinoids were tested together for impacts on *Daphnia magna* species, a species known to be highly *tolerant* to neonicotinoid toxicity, the effects included notable impacts on reproduction, growth, and survival, in correlation to chemical synergism.<sup>xx</sup> Due to the tendency for aquatic ecosystems to be contaminated by several neonicotinoid chemicals from a range of application sites as well as other chemicals present in surface water bodies, **EPA's final risk assessment should include the threat from combined exposure and synergistic effects of multiple pesticides.**

## 4. Limited Field Realistic Conditions and Lack of Evaluation of Sub-lethal Impacts to Ecosystem Functioning and Food Chains

The PARA addresses the lack of higher-tier data stating that the final risk assessment will include "*an independent review of mesocosm data,*" however this delay in analysis poses a significant risk to aquatic ecosystems. EPA, in its assessment of impacts to fish and aquatic phase amphibians notes:

---

<sup>4</sup> Imidacloprid, Clothianidin, Thiamethoxam, Dinotefuran, Acetamiprid

<sup>5</sup> Not updated as of July 11, 2017 <https://www.epa.gov/pesticide-science-and-assessing-pesticide-risks/aquatic-life-benchmarks-pesticide-registration>

*“While the risk of direct effects of imidacloprid to fish and amphibians is considered low, the potential exists for indirect risks to fish and aquatic-phase amphibians through reduction in their invertebrate prey base.”*

A more thorough analysis of available peer-reviewed research will show that the indirect risks to fish and aquatic-phase amphibians are a reality and that the continued use of imidacloprid and other persistent neonicotinoid chemicals weakens the base of the food-web and is detrimental to entire watershed ecosystems—including birds. **If EPA continues to disregard the indirect but significant impacts, then the repercussions will extend far beyond the aquatic invertebrate prey base.**

#### **5. Ignores Risks to Non-aquatic Species**

Initially intended to be a complete ecological risk assessment of imidacloprid, EPA justified its decision to only include aquatic risks, stating:

*“... a substantial body of aquatic monitoring and toxicity data have been generated for imidacloprid since the Agency’s last comprehensive risk assessment was conducted. In contrast, very little new data have been generated on the toxicity of imidacloprid to birds and mammals since the Agency’s most recent ecological risk assessments.”*

This is an underestimation of the research that has emerged showing risks to non-aquatic species—particularly birds, which are impacted by the use of neonicotinoid chemicals as shown in the findings of the comprehensive Palmer and Mineau report, *The Impact of the Nation’s Most Widely Used Insecticides on Birds*, as well as substantial other journal-published bird research. It also is a setback in finalizing the registration review and initiating regulatory action on these environmental contaminants.<sup>xxi</sup> **Rather than wait on the full ecological risk assessment, EPA should recognize the risks to aquatic species as well as the interconnection of aquatic and terrestrial environments and immediately restrict uses of imidacloprid to prevent these harms.**

#### **6. Strong Evidence of Risk, Yet No Regulatory Action**

EPA concluded in its PARA (emphasis added):

*“It is evident, however that **concentrations of imidacloprid** detected in streams, rivers, lakes and drainage canals **routinely exceed acute and chronic toxicity endpoints** derived for freshwater invertebrates.”*

Again, based on the substantial impacts to aquatic invertebrates, including ESA-protected species, happening on a wide-scale by registered uses, it is clear that EPA needs to take immediate action to restrict uses of imidacloprid and other neonicotinoid insecticides to prevent further damage to ecosystem services.

Furthermore, EPA identifies that:

*“...the risk findings summarized in this assessment are in general agreement with recent findings published by Canada’s Pest Management Regulatory Agency and the European Food Safety Authority.”*

EPA should follow PMRA's example in proposing a prompt full phase-out of imidacloprid for agricultural and outdoor uses. PMRA recognizes that due to imidacloprid's persistence and water solubility, regional restrictions will not be sufficient in mitigating risks. **EPA needs to enforce strong action now to prevent continued, potentially irreparable, damages to vulnerable species and ecosystems.**

**Due to the reasons above, as well as those outlined in detail in the attached two reports, which are incorporated into this comment by reference, *Water Hazard 2.0: Continued Aquatic Contamination by Neonicotinoid Insecticides in the United States (2017)* and *Water Hazard: Aquatic Contamination by Neonicotinoid Insecticides in the United States (2015)*, CFS urges EPA to take action to immediately restrict uses of imidacloprid to prevent further adverse impacts to aquatic ecosystems, pollinators, other vulnerable species, and the broader environment.**

### **Recommendations to EPA**

**The agency should:**

- 1. Expedite completion of the final risk assessment and the overall Registration Review for Imidacloprid, which is now at least two and likely three years behind the schedule to which EPA had committed.**
- 2. Conduct full ESA Sec. 7 compliance now, contemporaneous with the risk assessments in the Registration Review process, rather than afterwards, which would violate the ESA.**
- 3. In the final risk assessment, more accurately portray the risk posed by seed-coatings and include a thorough field-realistic analysis of imidacloprid seed-coatings to aquatic systems.**
- 4. Update its water quality benchmarks for imidacloprid using the newly proposed thresholds referenced in this comment.**
- 5. In the final risk assessment, include a comprehensive examination of the threats from additive and synergistic effects of combined exposure of imidacloprid and multiple other pesticides, fungicides, inerts and other compounds.**
- 6. Include higher-tier and mesocosm analyses to fully determine the risk to fish, amphibian, and bird species.**
- 7. Immediately enforce strong action to restrict uses of imidacloprid and other neonicotinoid insecticides to prevent continued, potentially irreparable, damages to vulnerable aquatic ecosystems.**

CC: California Department of Pesticide Regulation

**Attachments** –*Water Hazard 2.0: Continued Aquatic Contamination by Neonicotinoid Insecticides in the United States (2017)*; *Water Hazard: Aquatic Contamination by Neonicotinoid Insecticides in the United States (2015)*

For further information contact:  
Larissa Walker  
Pollinator Program Director | Policy Analyst  
Center for Food Safety  
660 Pennsylvania Ave. SE, Suite 302  
Washington, DC 20003  
(P): 202.547.9359 | (E): LWalker@CenterForFoodSafety.org

- 
- <sup>i</sup> Van Metre, P. C., Alvarez, D. A., Mahler, B. J., Nowell, L., Sandstrom, M., and Moran, P. (2017). Complex mixtures of Pesticides in Midwest US streams indicated by POCIS time-integrating samplers. *Environmental Pollution*, 220, 431-440.
- <sup>ii</sup> Morrissey, C. A., Mineau, P., Devries, J. H., Sanchez-Bayo, F., Liess, M., Cavallaro, M. C., and Liber, K. (2015). Neonicotinoid contamination of global surface waters and associated risk to aquatic invertebrates: a review. *Environment International*, 74, 291-303.
- <sup>iii</sup> Van Metre, P. C., Alvarez, D. A., Mahler, B. J., Nowell, L., Sandstrom, M., and Moran, P. (2017). Complex mixtures of Pesticides in Midwest US streams indicated by POCIS time-integrating samplers. *Environmental Pollution*, 220, 431-440.
- <sup>iv</sup> Hladik, M. L., Kolpin, D. W., and Kuivila, K. M. (2014). Widespread occurrence of neonicotinoid insecticides in streams in a high corn and soybean producing region, USA. *Environmental Pollution*, 193, 189-196.
- <sup>v</sup> Benton, E. P., Grant, J. F., Mueller, T. C., Webster, R. J., and Nichols, R. J. (2016). Consequences of imidacloprid treatments for hemlock woolly adelgid on stream water quality in the southern Appalachians. *Forest Ecology and Management*, 360, 152-158.
- <sup>vi</sup> Hladik, M. L., and Kolpin, D. W. (2016). First national-scale reconnaissance of neonicotinoid insecticides in streams across the USA. *Environmental Chemistry*, 13(1), 12-20.
- <sup>vii</sup> Sánchez-Bayo, F., Goka, K., and Hayasaka, D. (2016). Contamination of the Aquatic Environment with Neonicotinoids and its Implication for Ecosystems. *Frontiers in Environmental Science*, 4, 71.
- <sup>viii</sup> <http://pubs.acs.org/doi/abs/10.1021/acs.estlett.7b00081>
- <sup>ix</sup> <https://ehp.niehs.nih.gov/ehp515/>
- <sup>x</sup> Pest Management Regulatory Agency. (2016). Proposed Re-evaluation Decision PRVD2016-20, Imidacloprid. *Consumer product safety*. Retrieved from: <http://www.hc-sc.gc.ca/cps-spc/pest/part/consultations/prvd2016-20/prvd2016-20-eng.php#s3>
- <sup>xi</sup> Pest Management Regulatory Agency. (2016). Proposed Re-evaluation Decision PRVD2016-20, Imidacloprid. *Consumer product safety*. Retrieved from: <http://www.hc-sc.gc.ca/cps-spc/pest/part/consultations/prvd2016-20/prvd2016-20-eng.php#s3>
- <sup>xii</sup> Sappington, K. G., Ruhman, M. A., Housenger, J. (2016, December 22). Preliminary Aquatic Risk Assessment to Support the Registration Review of Imidacloprid. Washington: US Environmental Protection Agency Memorandum. Retrieved from: <https://www.regulations.gov/document?D=EPA-HQ-OPP-2008-0844-1086>
- <sup>xiii</sup> Bonmatin, J. M., Giorio, C., Girolami, V., Goulson, D., Kreutzweiser, D. P., Krupke, C., ... and Noome, D. A. (2015). Environmental fate and exposure; neonicotinoids and fipronil. *Environmental Science and Pollution Research*, 22(1), 35-67.
- <sup>xiv</sup> Sappington, K. G., Ruhman, M. A., Housenger, J. (2016, December 22). Preliminary Aquatic Risk Assessment to Support the Registration Review of Imidacloprid. Washington: US Environmental Protection Agency Memorandum. Retrieved from: <https://www.regulations.gov/document?D=EPA-HQ-OPP-2008-0844-1086>
- <sup>xv</sup> Sappington, K. G., Ruhman, M. A., Housenger, J. (2016, December 22). Preliminary Aquatic Risk Assessment to Support the Registration Review of Imidacloprid. Washington: US Environmental Protection Agency Memorandum. Retrieved from: <https://www.regulations.gov/document?D=EPA-HQ-OPP-2008-0844-1086>
- <sup>xvi</sup> Morrissey, C. A., Mineau, P., Devries, J. H., Sanchez-Bayo, F., Liess, M., Cavallaro, M. C., and Liber, K. (2015). Neonicotinoid contamination of global surface waters and associated risk to aquatic invertebrates: a review. *Environment International*, 74, 291-303.
- <sup>xvii</sup> US Environmental Protection Agency. [n.d.]. Aquatic life benchmark registration. Retrieved online at: <https://www.epa.gov/pesticide-science-and-assessing-pesticide-risks/aquatic-life-benchmarks-pesticide-registration>
- <sup>xviii</sup> Andersch, W., Jeschke, P., Thielert, W. (2010). Combination of methiocarb and one or more compounds selected from thiacloprid, thiamethoxam, acetamiprid, nitenpyram, and dinotefuran; effective animal pests control and for plant seed dressing. Google Patents. United States: Bayer CropScience AG; Iwasa, T., Motoyama, N., Ambrose, J. T., and Roe, R. M. (2004). Mechanism for the differential toxicity of neonicotinoid insecticides in the honey bee, *Apis mellifera*. *Crop Protection*, 23(5), 371-378.; Morrissey, C. A., Mineau, P., Devries, J. H., Sanchez-Bayo, F., Liess, M., Cavallaro, M. C., and Liber, K. (2015). Neonicotinoid contamination of global surface waters and associated risk to aquatic invertebrates: a review. *Environment International*, 74, 291-303.

---

<sup>xix</sup> Donley, N. (2016 July). Toxic concoctions: How the EPA ignores the dangers of pesticide cocktails. *Center for Biological Diversity*. Retrieved from:

[https://www.biologicaldiversity.org/campaigns/pesticides\\_reduction/pdfs/Toxic\\_concoctions.pdf](https://www.biologicaldiversity.org/campaigns/pesticides_reduction/pdfs/Toxic_concoctions.pdf)

<sup>xx</sup> Pavlaki, M. D., Ferreira, A. L., Soares, A. M., and Loureiro, S. (2014). Changes of chemical chronic toxicity to *Daphnia magna* under different food regimes. *Ecotoxicology and environmental safety*, 109, 48-55.

<sup>xxi</sup> Mineau, P., and Palmer, C. (2013 March). The impact of the nation's most widely used insecticides on birds. *American Bird Conservancy*. Retrieved from:

<https://extension.entm.purdue.edu/neonicotinoids/PDF/TheImpactoftheNationsMostWidelyUsedInsecticidesonBirds.pdf>