Henrika Stich

Noise Pollution, Public Health, and Potential Disproportionate Impacts on Persons with Disabilities and Sensory Sensitivities

To the Department of Ecology, Clean Energy Coordination:

I respectfully submit this additional scoping comment concerning the Programmatic Environmental Impact Statement ("PEIS") for stand-alone, utility-scale battery energy storage system ("BESS") facilities in Washington State.

The Washington State Department of Ecology Battery Storage PEIS states that Ecology is studying potential negative impacts of stand-alone, utility-scale BESS facilities, including measures to avoid, reduce, or mitigate those impacts. Ecology also acknowledges that previous battery studies identified higher noise levels among potential impacts.

For these reasons, I respectfully request that the PEIS specifically evaluate operational noise pollution as a potential public-health and accessibility concern, including whether residents with disabilities, neurodevelopmental conditions, sensory-processing differences, sleep disorders, and other conditions associated with sound sensitivity may experience disproportionate functional effects when utility-scale BESS facilities are located near residences.

This issue warrants specific consideration rather than being addressed solely through a generalized community-noise analysis.

1. Environmental Noise Should Be Evaluated as a Public-Health Issue

Noise should not be assessed solely as an issue of inconvenience, nuisance, or compliance with an average exterior decibel limit.

The World Health Organization Guidance on Environmental Noise identifies excessive environmental noise as capable of causing annoyance and reports evidence concerning increased risks of ischemic heart disease and hypertension, sleep disturbance, hearing impairment, tinnitus, and cognitive impairment, with increasing evidence concerning additional health outcomes including mental-health effects.

WHO's environmental-noise materials address environmental sources including transportation and industry rather than being limited to occupational noise.

I recognize that the WHO evidence principally concerns environmental noise generally and does not establish that BESS facilities specifically cause these health effects. Nevertheless, this evidence demonstrates why persistent environmental noise near residences warrants careful public-health evaluation when assessing a new industrial noise source.

Because the PEIS is intended to identify potential adverse impacts and measures to avoid, reduce, or mitigate them, I request that residential noise exposure, including potential impacts on noise-sensitive populations, be expressly included within the scope of the human-health and noise

analysis.

2. Persons With Disabilities and Sensory-Processing Differences Require Specific Consideration

An environmental analysis based solely upon the response of an "average" resident may fail to identify impacts experienced by individuals whose disabilities or health conditions affect their response to environmental stimuli.

The PEIS should investigate whether potentially noise-sensitive populations may include individuals with conditions such as:

attention-deficit/hyperactivity disorder ("ADHD");
sensory-processing differences;
autism spectrum conditions;
hyperacusis or tinnitus;
migraines associated with sound sensitivity;
post-traumatic stress disorder;
anxiety or panic disorders;
traumatic brain injury;
cognitive disabilities; and
insomnia and other sleep disorders.

Scientific evidence supporting examination of sensory-processing differences in certain populations.

For example, a 2025 systematic review and meta-analysis addressing ADHD and sensory processing evaluated 30 studies encompassing 5,374 participants. The researchers reported significantly greater sensory atypicalities among individuals with ADHD than control populations across several domains, including sensory sensitivity and sensory avoiding. The researchers also reported substantial heterogeneity among the studies.

This scientific evidence supports the more limited and relevant conclusion that sensory response is not uniform across the population and that an environmental review should investigate whether residents with heightened sensory sensitivity may experience persistent environmental noise differently from the general population.

3. The Residential Environment Deserves Particular Protection in the Noise Analysis

For individuals with sensory sensitivities or other noise-sensitive conditions, the home may be one of the most important environments in which external stimulation can be controlled.

Consequently, evaluation of a BESS facility near residences should not end with determining whether sound levels approach thresholds associated with hearing damage.

The PEIS should investigate whether chronic, recurring, intermittent, or tonal operational noise could interfere with:

sleep and nighttime recovery;

concentration;
learning and homework;
working from home;
communication;
ordinary indoor activities;
use of yards and other outdoor residential areas; and
the ability of noise-sensitive residents to obtain relief from unwanted environmental stimulation within their homes.

WHO materials identify sleep disturbance, annoyance, and cognitive impairment among the adverse outcomes associated with excessive environmental noise.

This general environmental-noise evidence provides a scientifically legitimate basis for asking Ecology to evaluate the issue before large industrial battery facilities are situated near residential communities.

4. The Analysis Should Not Rely Exclusively on a Single Average dBA Measurement

I respectfully request that Ecology evaluate whether reliance upon a single A-weighted average decibel measurement adequately characterizes the potential effect of BESS operations on nearby residences and noise-sensitive populations.

The PEIS should evaluate whether project-level acoustic assessments should consider additional characteristics including:

daytime and nighttime sound separately;
existing background or ambient sound conditions;
continuous versus intermittent sound;
tonal characteristics;
low-frequency sound components;
equipment cycling;
maximum sound events;
simultaneous operation of multiple pieces of equipment;
cumulative sound from the BESS together with relevant existing sources;
sound levels at nearby residences and other sensitive receptors;
potential indoor residential and bedroom exposure; and
reasonably foreseeable operating conditions associated with elevated noise generation.

This is particularly important in residential areas where the pre-existing nighttime acoustic environment may be substantially quieter than an industrial environment.

The question should therefore not merely be whether a facility satisfies an applicable numerical noise standard. The PEIS should also evaluate whether the methodology used to characterize noise adequately represents the actual residential exposure that nearby communities may experience.

5. Nighttime Noise and Sleep Deserve Separate Evaluation

Nighttime exposure should receive particular attention.

WHO identifies sleep disturbance as an adverse effect associated with excessive environmental noise and has incorporated sleep disturbance and annoyance into its work assessing environmental-noise health burdens.

For individuals who already experience neurological, sensory, psychological, or sleep-related vulnerabilities, disruption of the residential nighttime acoustic environment warrants explicit investigation.

I therefore request that Ecology evaluate whether project-level noise studies should assess BESS sound reaching nearby residences during the quietest nighttime conditions, including potential indoor bedroom exposure where technically appropriate.

Nighttime impacts should not be obscured by reliance solely upon a longer-term average that combines louder daytime ambient conditions with quieter nighttime conditions.

6. The PEIS Should Establish a Protective Noise-Assessment Framework

Because Ecology states that this study will lay the foundation for future project-level environmental reviews, the methodology developed at the programmatic stage is particularly important.

I respectfully request that Ecology evaluate whether future project-level BESS acoustic assessments should include:

preconstruction baseline ambient-noise measurements;
separate daytime and nighttime analysis;
evaluation at the nearest residences and sensitive receptors;
frequency-specific analysis where appropriate;
assessment of tonal and low-frequency characteristics;
intermittent and cycling noise;
simultaneous operation of relevant equipment;
reasonably foreseeable higher-noise operating conditions;
appropriate consideration of weather, topography, and sound propagation;
cumulative noise from relevant sources;
post-construction verification measurements; and
enforceable corrective mitigation where predicted acoustic performance is not achieved.

Establishing an appropriate methodology at the PEIS level would allow subsequent environmental reviews to evaluate residential noise impacts more consistently and transparently.

7. Disability and Accessibility Impacts Should Be Expressly Considered

Ecology's BESS scoping materials expressly state the Department's commitment to providing people with disabilities access to information and services and reference the Americans with Disabilities Act, Sections 504 and 508 of the Rehabilitation Act, and Washington State Policy #188.

I do not contend through this comment that those provisions independently require Ecology to reach a particular substantive conclusion concerning BESS noise.

Rather, consistent with Ecology's stated commitment to accessibility, I respectfully ask the Department to consider disability and sensory vulnerability as part of its evaluation of potential human-health and residential-noise impacts.

The PEIS should specifically examine whether siting a continuously or intermittently operating industrial facility near residences could create disproportionate functional impacts for individuals whose disabilities or health conditions make their residential acoustic environment particularly important.

The analysis should recognize individual variability and should not assume that everyone sharing a particular diagnosis will respond to sound in the same manner.

8. Mitigation Should Focus on the Facility Rather Than Shifting the Burden to Residents

Where potentially significant residential noise impacts are identified, I request that the PEIS evaluate avoidance and mitigation measures during planning and siting rather than waiting until residents experience adverse effects after facility construction.

Potential measures for evaluation should include, where technically appropriate:

adequate separation between major noise-producing equipment and residences;
facility layout and equipment orientation;
selection of quieter equipment;
acoustic enclosures or barriers;
nighttime operational controls where warranted;
post-construction noise monitoring;
accessible complaint procedures; and
enforceable corrective measures where actual operating noise exceeds modeled or permitted conditions.

Most importantly, mitigation should focus first on reducing noise at its source or through facility design and siting, rather than transferring responsibility to nearby residents.

Residents, particularly those with disabilities or sensory sensitivities, should not be expected to rely upon permanently closed windows, continuous personal hearing protection, relocation of bedrooms, or abandonment of normal use of outdoor areas as substitutes for appropriate facility siting and effective noise control.

9. Requested PEIS Finding and Scope of Analysis

I respectfully request that Ecology recognize the following principle within the PEIS:

Operational noise from utility-scale BESS facilities located near residences warrants evaluation as a potential public-health and accessibility concern, including consideration of nighttime exposure, existing ambient conditions, tonality, low-frequency characteristics, intermittency, cumulative noise, indoor residential exposure, and potentially noise-sensitive populations.

I further request that the PEIS establish an evidence-based technical framework for evaluating these factors during subsequent project-level environmental review and identify measures capable of avoiding, reducing, or mitigating significant residential noise impacts.

Ecology states that the BESS study is examining potential negative impacts of stand-alone utility-scale battery storage and methods for avoiding, reducing, or mitigating those impacts, and that the study will provide the foundation for later project-level environmental reviews.

For precisely this reason, noise impacts on persons with disabilities and other noise-sensitive residents should be examined at the programmatic stage rather than first being addressed after a specific facility has been designed, permitted, constructed, or placed into operation.

A scientifically responsible assessment does not require Ecology to presume that BESS noise will cause a particular disease or disability. It requires the Department to recognize that environmental-noise exposure can have health and functional consequences, that sensitivity to environmental sound varies across the population, and that potentially vulnerable populations should not be overlooked when evaluating industrial facilities proposed near residential communities.

Thank you for considering this comment and for ensuring that Washington's evaluation of utility-scale battery energy storage systems considers both the general population and residents who may experience heightened vulnerability to changes in their residential acoustic environment.

Respectfully submitted,

Henrika Stich
36702 SE Gravenstein Ct.
Snoqualmie, Washington

Supporting authorities

Washington State Department of Ecology, Scoping Document for Battery Energy Storage Systems Programmatic Environmental Impact Statement, Publication 26-06-017. Ecology's scoping materials describe the BESS PEIS and its accessibility commitment.

Washington State Department of Ecology, Battery Storage. Ecology states that it is studying potential negative impacts from stand-alone utility-scale BESS facilities and identifies higher noise levels among impacts found in its previous battery studies.

World Health Organization, Guidance on Environmental Noise. WHO identifies a range of health effects associated with excessive environmental noise, including annoyance, sleep disturbance and cognitive impairment, among others.

World Health Organization, Environmental Noise. WHO discusses environmental noise exposure and associated health considerations.

Sensory Processing in Individuals With Attention-Deficit/Hyperactivity Disorder Compared With Control Populations: A Systematic Review and Meta-analysis. The 2025 analysis reports greater

sensory-processing atypicalities among people with ADHD across multiple domains while also reporting heterogeneity in the available studies.

DEPARTMENT OF ECOLOGY State of Washington http://www.ecy.wa.gov

Publication Summary

Our Ecology website has changed, which can cause broken links.
To report these, please *contact us* with the publication and broken link.

TITLE	**Scoping document for Battery Energy Storage Systems Programmatic Environmental Impact Statement**
	Publication number: 26-06-017 — Date Published: August 2026
VIEW NOW	Scoping document for Battery Energy Storage Systems Programmatic Environmental Impact Statement (Number of pages: 23) (Publication Size: 1117KB) Trouble viewing? Try these free options. • Get the latest Adobe Reader for PDFs. • For Excel or Word viewing get Open Office, Microsoft OneDrive, DropBox Basic or a mobile app at your favorite app store.
AUTHOR(S)	Clean Energy Coordination
DESCRIPTION	Ecology is studying the potential adverse impacts of battery energy storage systems in Washington – including ways potential negative effects can be avoided, reduced or mitigated. The scoping document provides initial information about the evaluation process and a general overview of battery energy storage.
REQUEST A COPY	The mission of the Department of Ecology is to protect, preserve, and enhance Washington's environment. To help us meet that goal, please consider the environment before you print or request a copy. **ADA Accessibility** The Department of Ecology is committed to providing people with disabilities access to information and services by meeting or exceeding the requirements of the Americans with Disabilities Act (ADA), Section 504 and 508 of the Rehabilitation Act, and Washington State Policy #188. Visit Ecology's website for more information. • Shorelands & Environmental Assistance Order Form
CONTACT	Clean Energy Coordination at 360-407-6600 or cleanenergy@ecy.wa.gov
KEYWORDS	Programmatic Environmental Impact Statements, battery energy storage systems, PEIS, BESS, scoping document

WEB PAGE	Programmatic environmental impact statements
RELATED PUBLICATIONS	Title: State Environmental Policy Act Final Programmatic Environmental Impact Statement for Utility-Scale Solar Energy Facilities in Washington State State Environmental Policy Act Final Programmatic Environmental Impact Statement for Utility-Scale Onshore Wind Energy Facilities in Washington State State Environmental Policy Act Final Programmatic Environmental Impact Statement for Green Hydrogen Energy Facilities in Washington State

Copyright © Washington State Department of Ecology. See https://ecology.wa.gov/About-us/Accountability-transparency/Our-website/Copyright-information.

META-ANALYSIS

Sensory Processing in Individuals With Attention-Deficit/Hyperactivity Disorder Compared With Control Populations: A Systematic Review and Meta-analysis

Lucie Jurek[a,b,c,*], MD, PhD, Arnaud Duchier[d], MD, Christophe Gauld[e], MD, PhD, Léonie Hénault[a], MSc, Caroline Giroudon[f], MSc, Pierre Fourneret[f], MD, PhD, Samuele Cortese[c,g,h,i], MD, PhD, Mikail Nourredine[b,c,f], MD, MSc

Objective: To evaluate the association between attention-deficit/hyperactivity disorder (ADHD) and atypical sensory processing patterns.

Method: For this systematic review and meta-analysis, PubMed, Embase, and Web of Science were searched from their inception until June 30, 2024. Studies examining sensory processing patterns using questionnaires in participants with a diagnosis of ADHD compared with healthy controls were included. The study risk of bias was assessed using a modified Newcastle-Ottawa Scale. A random-effects meta-analysis was conducted using R Version 4.3.1 software, considering sensory modulation severity atypicities as the primary outcome. The risk of publication bias was explored, and sensitivity analyses were conducted to test the robustness of findings.

Results: After screening 10,750 references, 30 studies (reported in 32 publications) encompassing 5,374 participants were included, with 23 studies focusing on children and 7 focusing on adults. Patients with ADHD presented with significantly more severe sensory atypicities than the control group in multiple domains: sensory sensitivity (k = 13, SMD = 1.17, 5% CI [0.75, 1.59], I^2 = 87%), sensory avoiding (k = 11, SMD = 1.15, 95% CI [0.66, 1.64], I^2 = 92%), low sensory registration (k = 23, SMD = 1.22, 95% CI [0.89, 1.56], I^2 = 92%), and sensory seeking (k = 15, SMD = 1.23, 95% CI [0.48, 1.97], I^2 = 97%). Heterogeneity was high in all analyses. Only 9 studies were deemed at overall low risk of bias.

Conclusion: Even though available ADHD clinical guidelines do not specifically mention the need to assess sensory processing in ADHD, this meta-analysis suggests that this should be systematically explored in the evaluation of children and adults referred for ADHD.

Study registration information: Association between ADHD and sensory processing disorder: A systematic review and meta-analysis; https://www.crd.york.ac.uk/PROSPERO/view/CRD42022325271.

Diversity & Inclusion Statement: We actively worked to promote sex and gender balance in our author group. One or more of the authors of this paper self-identifies as a member of one or more historically underrepresented sexual and/or gender groups in science. One or more of the authors of this paper self-identifies as a member of one or more historically underrepresented racial and/or ethnic groups in science.

Key words: attention-deficit/hyperactivity disorder; meta-analysis; sensory modalities; sensory processing; systematic review

J Am Acad Child Adolesc Psychiatry 2025;■(■):■-■.

Attention-deficit/hyperactivity disorder (ADHD) is a neurodevelopmental disorder characterized by symptoms of inattention, hyperactivity-impulsivity, or both that interfere with daily functioning and development.[1] It is the most prevalent neurodevelopmental condition, is more common in males, and is estimated to affect about 5% of youth[2] and 3% of adults.[3] ADHD is frequently associated with other physical and psychiatric disorders, such as neurodevelopmental, conduct, anxiety, and mood disorders.[4,5]

Theorized initially in the 1970s by Ayres,[6] sensory integration is the process by which one registers, modulates, and discriminates sensations received through sensory systems (visual, auditory, gustatory, olfactory, tactile, proprioceptive, and vestibular) to produce purposeful, adaptive behaviors in response to the environment. Atypical sensory processing patterns (SPPs) can impact the development of behavioral, emotional, motor, and cognitive domains.[7,8] Dunn's model[9] describes sensory modulation as the interaction between the neurological threshold and self-

regulation. A low neurological threshold means that a person's neurons are activated and cause a response even to low-intensity stimuli. Conversely, a high neurological threshold indicates that triggering the same neurological response requires more intense stimuli. According to Dunn, sensory modulation can be classified into four domains (or quadrants): sensitivity (low threshold and passive self-regulation strategy); avoiding (low threshold and active self-regulation strategy); seeking (high threshold and active self-regulation strategy); and registration (high threshold and passive self-regulation strategy).

The estimated prevalence of atypical SPPs ranges from 5% to 16% in neurotypical people[10,11] and from 30% to 80% in people with developmental disabilities.[12-14] Atypical SPPs may affect cognitive functions and psychosocial development,[9] impact the intensity and enjoyment of engaging in social relationships,[15] and increase emotional and attentional regulation difficulties.[16]

In *DSM-5-TR*, atypical SPPs are not included as a specific diagnosis category.[1] However, sensory processing disorder has been included in some diagnostic classifications for developmental disorders in early childhood (eg, Developmental Disorders of Infancy and Early Childhood [DC: 0-5][17]) and is now listed among the possible symptoms of autism spectrum disorders in *DSM-5-TR*.[1]

Previous studies showed that individuals with ADHD may also be at a higher risk of having atypical SPPs.[18,19] However, the exact extent to which SPPs are atypical in, namely, ADHD is unclear, and indeed current clinical guidelines for ADHD do not include the assessment of SPPs.

Thus, investigating SPPs in ADHD has high clinical relevance, as it may inform future clinical guidelines and recommendations. This systematic review and meta-analysis aimed to quantitatively synthesize studies comparing SPPs in participants with ADHD and controls from the general population.

METHOD

This study is reported according to the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines[20] (see Supplement 1, available online, for the checklists). This review was registered in PROSPERO in May 2022 (CRD42022325271).

Literature Search

The electronic databases Medline, Embase, and Web of Science were searched from inception until June 30, 2024. The search strategy was designed by an information specialist and specialists in the field and included terms related to ADHD and SPPs. The terms were combined with MEDLINE filters to exclude nonrelevant review articles. The search terms were adapted for use with other bibliographic databases in combination with database-specific filters where these are available. From the last search until September 1, 2024, monitoring of the 3 databases was performed, but no new study was included. We used a combination of free text and thesaurus terms for concepts relevant to the topic. The search algorithms are described in Supplement 1, available online. The reference lists of reviews related to our subject were also screened.[14,21-24]

Study Selection

The inclusion and exclusion criteria were as follows:

1. We included studies in which ADHD was diagnosed by a clinician with or without use of tools based on established diagnostic criteria (including *DSM-III* and later editions and *ICD-9* and later revisions). Studies including participants with ADHD symptoms only without a formal clinical diagnosis were excluded.
2. Studies including participants with more than 20% of comorbid autism spectrum disorder, intellectual disability, or genetic disorders were not included.
3. Studies included a control group without ADHD. A clinical control group (homogeneous diagnosis in the control group) was not accepted.
4. Original cohort, case-control, and cross-sectional studies were included. Case reports were excluded.

Outcomes were the severity of SPP atypicities and the risk of being hyposensitive or hypersensitive. Assessment of each outcome could occur before or after the diagnosis of ADHD using questionnaires (self-reports, parent reports, teacher reports or semistructured interviews). Sensory-based motor disorders (dyspraxia and postural disorder) were not evaluated in this systematic review.

Screening and Data Extraction

Two independent reviewers (A.D. or Ch.G. and L.J.) conducted the literature search. Duplicate reports of the same studies were removed. Duplicates were managed using Zotero for the first search and Covidence for the update. Two reviewers (A.D. or Ch.G. and L.J.) checked eligibility of the studies from titles and abstracts first, followed by full-text reading.

The following variables were extracted by 2 independent reviewers (A.D. or Ch.G. and L.J.) using a standardized extraction form: study design; sample size and description (eg, clinical-based, population-based);

comparison group description (eg, matching criteria, sibling controls, healthy volunteers); ADHD and SPP assessment methods; population characteristics (eg age, gender, race/ethnicity, IQ, medication, comorbidities); and outcome results, including means, SDs, and prevalence in each group.

In case of discrepancies between the 2 reviewers during the selection or the data extraction process, a third author (M.N.) was consulted. In case of missing information to assess the eligibility criteria or to complete data extraction, the study's corresponding author was contacted.

Risk-of-Bias Assessment

Two reviewers (A.D. or Ch.G. and L.J.) independently assessed the quality of each study by using a modified Newcastle Ottawa Scale (NOS). The NOS contains 7 items categorized into 3 dimensions: selection, comparability, and outcome. The maximum total score was 10 points. To better reflect efforts to ensure comparability across study populations, we assigned an additional point for representative samples. The evaluation of the NOS outcome dimension was modified to allow a maximum of 1 point, as only unblinded studies were included (see Supplement 1, available online, for more details on the NOS). As in previous studies, we considered a study that scored $\geq$7 to have a low risk of bias. Studies scoring between 4 and 7 points were at an unclear risk of bias, whereas studies scoring less than 4 points were at a high risk of bias.[25]

Data Analysis

Outcomes. Outcomes were the severity of SPP atypicities and the risk of being hyposensitive or hypersensitive. The severity of SPP atypicities was divided as follows: the sensory modulation disorder according to each Dunn quadrant (sensory sensitivity, low sensory registration, sensory seeking, and sensory avoiding), using continuous measures, and the sensory discrimination disorder according to each sensory modality (tactile, auditory, visual, oral [taste/smell], movement [vestibular], and body position [proprioception]), using continuous measures.

The assessment tools for each continuous outcome are outlined in Supplement 1, available online. In one study[26] using the Sensory Over-Responsivity scale, the taste and smell domains were combined as "oral sensory." To calculate the SD, a correlation of 0.41 between the 2domains was used.[27] For the risk of being hyposensitive or hypersensitive (binary outcome), the number of participants classified as hypersensitive ("more than others" or "much more than others" on sensory sensitivity) and hyposensitive ("less than others" or "very much less than others" on low registration) in both groups were retrieved. Primary outcomes were sensory modulation disorder severity according to each Dunn quadrant.

Statistical Analyses. For continuous outcomes, effect size estimates were based on standardized mean differences (SMDs) between the ADHD and control groups using Hedges g small sample bias adjustment.[28] For binary outcomes, pooled odds ratios (ORs) were estimated using the number of hypersensitive or hyposensitive diagnoses in participants with ADHD and controls. We conducted a random-effects meta-analysis for all outcomes. The Hartung-Knapp adjustment was used to estimate CI.[29] Pooled SMDs and ORs were estimated using the inverse variance method. Heterogeneity was quantified using τ^2 (estimated using restricted maximum likelihood) and I^2 statistics.

Prespecified subgroup analyses were conducted for all outcomes, with subgroups defined by study population (children or adults) and risk of bias (low, unclear, or high). We also conducted 3 post hoc subgroup analyses. The first analysis was based on the type of SPP assessment tool used; the second analysis separated studies in which ASD was clearly excluded from studies where the exclusion of ASD was unclear (eg, not reported, self-reported diagnosis, or use of a screening tool); and the third analysis separated studies into those that excluded participants with low IQ, studies that did not exclude participants with low IQ, and studies where information on possible exclusion based on IQ was not available.

We ran prespecified meta-regressions with the following variables: mean age, proportion of female participants, and proportion of ADHD participants taking medication. A post hoc meta-regression with publication year was performed. Publication bias was visually assessed using a funnel plot and the Egger regression test for asymmetry. In case of asymmetry, a trim-and-fill method was used to adjust for publication bias in the meta-analysis.[30] Regression-based tests were conducted if at least 10 studies were available per predictor. All analyses were conducted using the meta package version 7.0-0[31] in R version 4.3.1.[32]

Deviation From Protocol. We were unable to conduct a sensitivity analysis on subtypes of ADHD or age at diagnosis of ADHD, as this information was rarely reported. The impact of the study design and different control groups was not assessed, as study designs and control groups tended to be the same across studies.

RESULTS

Study Selection

The literature search generated 10,750 references. After removal of duplicates, 6,526 records were screened based on

the title and abstract, of which 84 were assessed for eligibility by full-text review. Of these 84 articles, 52 were excluded. A total of 30 studies (32 reports) were included in the quantitative synthesis. The PRISMA flowchart and the reasons for exclusions are presented in Supplement 1, available online. Of the 15 authors who were contacted to retrieve missing information from their study, only 1 author responded.

Study Characteristics

The characteristics of the included studies are presented in Table 1.[18,19,26,33-59] The total population included 5,374 participants (ADHD participants = 1,656 and controls = 3,718). Studies were published between 2001 and 2024. They were conducted in North America (k = 8), Europe (k = 7), Middle East (k = 6), Asia (k = 8), and South America (k = 1). All were cross-sectional studies.

Female participants were less represented than male participants, with a mean of 30% female participants (range 0%-67%). The mean age ranged from 4.7 to 34 years in the ADHD group and 4.8 to 32.1 in the control group. Of studies, 23 focused on children, 6 focused on adults, and 1 focused on adolescents and young adults (>16 years old).

Ethnicity of all participants was reported in 4 studies, 1 study provided information on ethnicity only for the control group, and 25 studies did not report ethnicity at all. Among the studies reporting ethnicity, the majority of the population was White (the term Caucasian is used Table 1 as it appears in the original studies included in this review).

Concerning IQ, 20 studies excluded participants with low IQ, 2 did not exclude participants with low IQ, and 9 did not provide any information on this criterion. The mean IQ in the ADHD group ranged from 72.06 to 119.4, whereas mean IQ in the control group ranged from 94.3 to 119.9.

The ADHD diagnosis was assessed using diagnostic criteria (*DSM-IV*, *DSM-IV-TR*, *DSM-5*, or *ICD-10*) in 18 studies or by a clinician without indicating the precise tool used. Other studies used chart reviews or mixed methods to assess diagnosis.

For participants with ADHD, medication was an exclusion criterion in 2 studies. Nine studies included participants with medication (see Supplement 1, available online, for details), and 19 studies did not provide any information on medication.

Comorbidities (excluding ASD) were described in 9 studies. Oppositional defiant disorder was the most frequent comorbidity. Comorbidities were excluded from 7 studies, and 14 studies did not provide any information. Concerning ASD, 20 studies excluded participants with ASD, 1 study included 4 participants with ADHD and ASD, and 9 studies did not provide any information regarding ASD.

The included studies used the following questionnaires to assess SPP: the Sensory Profile (SP) (k = 8, of which 2 studies used an adapted Chinese version [C-SP]), the Short Sensory Profile (SSP) (k = 6), the Adolescent/Adult Sensory profile (AASP) (k = 7), the Sensory Profile-2 (SP-2) (k = 3), the Sensory Over-Responsivity (SOR) scale (k = 1), the Sensory Processing 3-Dimensions Scale (SP-3DS) (k = 1), the Sensory Processing Measure (k = 2), the Touch Inventory for Elementary-School-Aged Children (TIE) (k = 1). One study used a newly designed and validated tool, the Sensory Perception Questionnaire (SPQ).

Risk of Bias

The overall quality of each study is reported in Supplement 1, available online. Nine studies were considered to have a low risk of bias, 19 studies were considered to have an unclear risk of bias, and 2 studies were considered to have a high risk of bias.

Major issues were found in the population selection. Only 3 studies estimated an a priori sample size, and only 2 studies described nonrespondents. The sampling strategy was rarely based on randomization or selection of all eligible participants. In most studies, the ADHD and control groups were sampled from different populations (outpatients for the ADHD group and volunteers from the general population for the control group, sometimes from the same region/schools).

Meta-analyses: Sensory Quadrants

This section presents the main results. All sensitivity analyses (subgroups, meta-regressions, leave-one-out analysis) and publication bias analyses plots are presented in Figures S1 to S33, available online.

Sensory Sensitivity. The meta-analysis on sensory sensitivity included 13 studies (ADHD participants = 626, controls = 991). Sensory sensitivity was higher in the ADHD group compared with the control group (SMD = 1.17, 95% CI [0.75,1.59]), observed in both adults and children (Figure 1). Heterogeneity was high (I^2 = 87%). A meta-regression on the year of the study revealed a significant effect with a larger SMD in most recent studies (estimate = 0.096, 95% CI [0.047, 0.144]), reducing the heterogeneity to 68.65%. Subgroup analyses indicated a significant difference based on the risk of bias ($p < .01$). When focusing only on studies with a low risk of bias, the difference between groups was no longer significant, with large CI explained by the insufficient number of studies

TABLE 1 Characteristics of Populations in Included Studies

Authors, year, reference	Country	Setting	N	% Female	Age, y, mean (SD))	Race/ethnicity	IQ, mean (SD)	Type of ADHD diagnosis	Comorbidity	ADHD medication	Sensory assessment
Mangeot *et al.*, 2001[33]	USA	Control: volunteers	30	30	8.2 (2)	NR	108.91 (10.68)	Clinical diagnosis (CAP)	NR	Total = 15; 8 MPH, 2 clonidine, 5 others	SSP
		ADHD: outpatients	26	31	8.3 (2.4)		72.06 (9.69)				
Dunn and Bennett, 2002[19]	USA	Control: randomly selected matched controls	70	13	8.6	NR, discussed by authors as a limitation	NR	*DSM-IV*	ADHD: 23 ODD, 1 posttraumatic stress disorder, 1 adjustment disorder, 10 learning disorders, 2 encopresis ASD excluded	Total = 52	SP
		ADHD: outpatients	70	13	8.6						
Yochman *et al.*, 2004[18]	Israel	Control: matched controls, cohort	46	20	4.8 (0.62)	NR	Low IQ excluded	Clinical diagnosis (pediatrician)	ASD excluded	NR	SP
		ADHD: cohort (same as control)	48	19	4.7 (0.76)						
Bröring *et al.*, 2008[34] (males)	Europe	Control: volunteers from same region	16	0	9.7 (0.6)	All children were European Caucasian[a] descent	Low IQ excluded	*DSM-IV-TR*	ASD excluded	Total = 28; 25 psychostimulant, 1 nonstimulant, 2 combination	TIE
		ADHD: outpatients	35	0	10 (1.7)						
Bröring *et al.*, 2008[34] (females)	Europe	Control: volunteers from same region	19	100	9.4 (0.8)	All children were European Caucasian[a] descent	Low IQ excluded	*DSM-IV-TR*	ASD excluded	Total = 8; 6 psychostimulant, 2 nonstimulant	TIE
		ADHD: outpatients	12	100	10 (1.1)						
Cheung and Siu, 2009[35]	China	Control: randomly selected from schools	1,840	50.3	7.25 (2.8)	NR	NR	*DSM-IV-TR*	ASD excluded	NR	C-SP
		ADHD: inpatients	114	NA	7.9 (1.4)						
Miller *et al.*, 2012[36]	USA	Control: volunteers	57	49	8.09 (2.68)	African American (3%), Asian (3%), Caucasian[a] (89%), Hispanic (2%), Native American (1%), other (2%)	100.95 (11.3)	Clinical diagnosis	Excluded (ASD excluded as well)	Total = 17 9 MPH, 4 dextroamphetamine, 1 fluoxetine, 3 combination	SSP
		ADHD: outpatients	37	24	9.99 (2.26)		73.92 (8.65)				

(*continued*)

TABLE 1 Continued

Authors, year, reference	Country	Setting	N	% Female	Age, y, mean (SD))	Race/ethnicity	IQ, mean (SD)	Type of ADHD diagnosis	Comorbidity	ADHD medication	Sensory assessment
						ethnicities No differences between groups with respect to ethnicity ($p > .05$)					
Lin *et al.*, 2013[37]	Taiwan	Control: matched children from local schools	20	0	9.1 (1.79)	NR	Low IQ excluded	*DSM-IV-TR*	NR	NR	C-SP
		ADHD: outpatients	20	0	8.64 (2.57)						
Lufi and Tzischinsky, 2014[38]	Israel	Control: volunteers from local schools	28	50	15.64 (1.5)	NR	NR	Clinical diagnosis (CAP)	NR	Total = 27 MPH only	AASP
		ADHD: outpatients	30	10	15.1 (1.27)						
Shimizu *et al.*, 2014[39]	Brazil	Control: volunteers from same school	37	19	8.9 (1.49)	NR	Low IQ excluded	*DSM-IV-TR*	13 affective disorder, 13 anxiety disorder, 21 ODD, 22 CD ASD excluded	NR	SP
		ADHD: outpatients	37	19	8.9 (1.49)						
Pfeiffer *et al.*, 2015[40]	USA	Control: mixed (outpatients, volunteers, etc)	27	52	8.3 (1.6)	African American (12.8%), Asian (12.8%), Caucasian[a] (61.7%), Hispanic (6.4%), Native American (2.1%), other (4.3%) ethnicities The group without ADHD included more Asian American children ($\chi^2 = 12.34$, $p < .05$)	Low IQ excluded	Mixed (clinician, *DSM-IV-TR*)	3 ADHD and comorbid diagnosis ASD excluded	Total = 12	SPM
		ADHD: outpatients	20	25	9.1 (1.3)						
Bijlenga *et al.*, 2017[41]	Netherlands	Control: norm group from AASP	496	51	NA	92% Caucasian in norm group; no information for ADHD group	NR	DIVA-5	42 sleep disorders, 22 depression, 17 anxiety disorder, 20 drug dependence, 4 autism, 11 impaired hearing, 1 visual disability, 5 olfactory and/or gustatory disabilities, 3 tactile disabilities	Total = 48; 30 ADHD medication and 18 antidepressant	AASP
		ADHD: outpatients	116	38	32 (10.2)						

(*continued*)

TABLE 1 Continued

Authors, year, reference	Country	Setting	N	% Female	Age, y, mean (SD))	Race/ethnicity	IQ, mean (SD)	Type of ADHD diagnosis	Comorbidity	ADHD medication	Sensory assessment
Sanz-Cervera *et al.*, 2017[42]	Spain	Control: volunteers from local schools	27	30	6.28 (1.11)	NR	98.04 (17.21)	*DSM-IV-TR*	Excluded (ASD excluded as well)	NR	SPM
		ADHD: outpatients	21	14	6.81 (1.1)		87.43 (14.38)				
Little *et al.*, 2018[43]	USA	Control: matched controls (cohort)	84	25	8.77	Asian (3.1%), Black (16%), Hispanic (20.4%), White (51.2%), other (9.3%) ethnicity	NR	Chart review	ASD excluded	NR	SP-2
		ADHD: cohort (volunteers), same cohort as control	78	22	9.7						
Mimouni-Bloch, 2018[44,b]	Israel	Control: volunteers	39	59	9.46 (0.9)	NR	Low IQ excluded	*DSM-5*	Excluded (ASD excluded as well)	NR	SSP
		ADHD: outpatients	38	29	9.31 (0.85)						
Jegadeesan *et al.*, 2020[45]	India	Control: volunteers from local schools	50	NA	NA	NR	Low IQ excluded	Clinical diagnosis (pediatrician)	NR	NR	SSP
		ADHD: outpatients	50	NA	NA						
Kamath *et al.*, 2020[46]	USA	Control: cohort	22	55	21.14 (2.25)	NR	111.18 (10.14)	*DSM-5*	Excluded (except depression and anxiety)	NR	AASP
		ADHD: cohort (same as control)	23	61	21.87 (1.98)		109.1 (8.75)				
Ohta *et al.*, 2020[47]	Japan	Control: volunteers	58	16	29.4 (6.7)	NR	107.7 (7.7)	*DSM-IV-TR*	Excluded (ASD excluded as well)	Total = 32 30 ADHD medication and 2 antipsychotics	AASP
		ADHD: outpatients	55	24	31.2 (8.8)		106.3 (12.5)				
Adra *et al.*, 2021[48]	USA	Control: volunteers from general population	29	69	27 (5.8)	NR	119.9 (13.9)	*DSM-5*	None reported, ASD excluded	NR	SP-3DS
		ADHD: volunteers from general population	25	56	26.4 (6.7)		119.4 (13.5)				
Dellapiazza *et al.*, 2021[49]	France	Control: matched children from local schools	31	39	9.3 (1.6)	NR	NR	*DSM-5*	ASD excluded	No medication	SP
		ADHD: outpatients	28	18	9.2 (1.8)		109 (16.2)				
Gomez *et al.*, 2021[50]	Philippines	Control: volunteers from local schools	42	0	8.9 (1.52)	NR	Low IQ excluded	Clinical diagnosis (therapy center)	None reported, ASD excluded	NR	SP
		ADHD: outpatients	24	0	9.4 (1.71)						
Grinblat and Rosenblum, 2022[51,b]	Israel	Control: matched volunteers	52	69	32.1 (6.56)	NR	Low IQ excluded	Clinical diagnosis	75% of ADHD group had at least 1 psychiatric comorbidity	NR	AASP
		ADHD: volunteers	69	65	33.42 (6.39)						
Sobhy *et al.*, 2022[52]	Egypt	Control: volunteers from nursery and schools	40	65	4.53 (0.98)	NR	NR	*DSM-5*	ASD excluded	NR	SPQ
		ADHD: outpatients	30	27	4.51 (0.96)						
Darnal *et al.*, 2023[53]	India	Control: volunteers from local schools	139	NA	6-18	NR	Low IQ excluded	*ICD-10*	Excluded (ASD excluded as well)	NR	SSP
		ADHD: NR	100	NA							
Panda *et al.*, 2023[54]	India	Control: matched controls	33	9	10.8 (1.6)	NR	94.3 (2.1)	*DSM-5*	More than 50% of ADHD group had	Total = 63; 34 atomoxetine, 22	SP-2

(*continued*)

TABLE 1 Continued


Authors, year, reference	Country	Setting	N	% Female	Age, y, mean (SD))	Race/ethnicity	IQ, mean (SD)	Type of ADHD diagnosis	Comorbidity	ADHD medication	Sensory assessment
		ADHD: outpatients	66	9	10.7 (1.9)		90.1 (3.5)		a learning disorder or ODD; 30% had sleep problems; some reported CD, anxiety, or depression	MPH, 2 clonidine, 5 atomoxetine + MPH	
Rani *et al.*, 2023[55]	India	Control: matched controls	34	6	8.35 (2.47)	NR	94.8 (5.8)	*DSM-5*	ASD excluded 31.6% had a psychiatric comorbidity (ODD, anxiety, or CD)	NR	SP-2
		ADHD: outpatients	38	5	8.45 (2.27)		92.1 (6.4)				
Schramm *et al.*, 2023[56]	Germany	Control: matched controls	14	36	32.14 (9.93)	NR	Low IQ excluded	Clinical diagnosis	ASD excluded Excluded	Total = 10; 9 MPH and 1 bupropion	AASP
		ADHD: outpatients	15	33	34 (7.1)						
Schulz *et al.*, 2023[57]	Canada	Control: cohort	163	44	11.29 (3.46)	NR	109.56 (12.14)	Clinical diagnosis	NR	NR	SSP
		ADHD: cohort (same as control)	317	27	10.82 (3.43)		93.15 (14.87)				
Fabio *et al.*, 2024[58]	Italy	Control: matched children, cohort	69	35	16.2 (1.9)	NR	NR	*DSM-5*	Excluded	NR	AASP
		ADHD: cohort (same as controls)	69	35	Age from global cohort						
Martínez-González *et al.*, 2024[26]	Spain	Control: volunteers randomly selected	76	62	19.18 (4.44)	NR	Low IQ excluded	Clinical diagnosis	Excluded (ASD excluded as well)	NR	SOR
		ADHD: outpatients	15	53	18.87 (2.41)						
Mohammed *et al.*, 2024[59]	Egypt	Control: control from nursery and schools	30	27	6.04 (2.29)	NR	NR	*DSM-5*	NR	No medication	SP
		ADHD: outpatients	30	27	6.03 (2.17)						

Note: *ADHD = attention-deficit/hyperactivity disorder; AASP = Adolescent/Adult Sensory Profile; ASD = autism spectrum disorder; CAP = child and adolescent psychiatrist; CD = conduct disorder; C-SP = Chinese Version Sensory Profile; DIVA-5 = Diagnostic Interview for ADHD in Adults; MPH = methylphenidate; NA = not available; NR = not reported; ODD = oppositional defiant disorder; SOR = Sensory Over-Responsivity (scale); SP = Sensory Profile; SP-2 = Sensory Profile-2; SP-3DS = Sensory Processing 3-Dimensions Scale; SPM = Sensory Processing Measure; SPQ = Sensory Perception Questionnaire; SSP = Short Sensory Profile; TIE = Touch Inventory for Elementary-School-Aged Children.*

[a]*The term Caucasian is used here as it appears in the original studies included in this review.*

[b]*The Grinblat and Rosenblum study from 2022[51] and the Mimouni-Bloch study from 2018[44] were kept for the analyses, but results are also reported in Grinblat and Rosenblum from 2021[60] and Mimouni-Bloch et al. in 2017.[61]*

FIGURE 1 Forest Plot of Meta-analysis on Sensory Sensitivity



Note: *ADHD = attention-deficit disorder with or without hyperactivity; SMD = standardized mean difference. Please note color figures are available online.*

(n = 4) and the high heterogeneity. No outliers were detected in the leave-one-out sensitivity analysis, and no publication bias was found on funnel plot visual inspection or the Egger test.

Sensory Avoiding. The meta-analysis on sensory avoiding included 11 studies (ADHD participants = 558, controls = 925). Sensory avoiding was higher in the ADHD group compared with the control group (SMD = 1.15, 95% CI [0.66, 1.64]) in adults as well as children (Figure 2). Heterogeneity was high (I^2 = 92%). Subgroup analyses indicated a significant difference based on the risk of bias (p < .01). When focusing only on studies with a low risk of bias, the difference between groups was no longer significant with large CI explained by the insufficient number of studies (n = 3) and the high heterogeneity.


No outliers were detected in leave-one-out sensitivity analysis. Publication bias was found on funnel plot visual inspection. The adjusted result with trim-and-fill procedure showed no difference between the groups (SMD = 0.57, 95% CI [−0.03, 1.16]).

Low Sensory Registration. The meta-analysis on low sensory registration included 23 studies (ADHD participants = 1,374, controls = 1,677). Low sensory registration was higher in the ADHD group compared with the control group (SMD = 1.22, 95% CI [0.89, 1.56]), in both adults and children (Figure 3). Heterogeneity was high (I^2 = 92%). Meta-regression on age revealed a significant effect (estimate = 0.04, 95% CI [0.003, 0.076]) with higher SMD with higher age. Accounting for age reduced the heterogeneity to 88.52%.

No outliers were detected in leave-one-out sensitivity analysis. Publication bias was found on funnel plot visual inspection and the Egger test. The adjusted result with the trim-and-fill procedure remained higher in the ADHD group (SMD=0.67, 95%CI [0.26, 1.9]).

Sensory Seeking. The meta-analysis on sensory sensitivity included 15 studies (ADHD participants = 968, controls = 1183). Sensory seeking was significantly higher in the ADHD group compared with the control group (SMD = 1.23, 95% CI [0.48, 1.97]). Heterogeneity was high (I^2 = 97%). No difference in sensory-seeking behaviors was found

FIGURE 2 Forest Plot of Meta-analysis on Sensory Avoiding



Note: ADHD = attention-deficit disorder with or without hyperactivity; SMD = standardized mean difference. Please note color figures are available online.

between participants with ADHD and controls for the adult subgroup (SMD = 0.52, 95% CI [1.06, 2.10]) (Figure 4).

Subgroup analyses indicated a significant difference based on the risk of bias ($p < .01$), with only studies with low risk of bias studies showing a significant difference between groups; on the sensory tools ($p < .01$), with studies using AASP showing no difference; and on the exclusion of participants with ASD ($p = .02$), with no difference between group when the exclusion of ASD was unclear. No outliers were detected in leave-one-out sensitivity analysis. No publication bias was found on funnel plot visual inspection and the Egger test.


Meta-analyses: Sensory Modalities

The results on sensory modalities are synthesized in Figure 5. The detailed results, sensitivity, and publication bias analyses are presented in Figures S34 to S64, available online. Patients with ADHD showed greater difficulties than the control population in tactile processing (k = 25, SMD = 0.85, 95% CI [0.65, 1.24], I^2 = 88%), visual processing (k = 16, SMD = 1.02, 95% CI [0.75, 1.30], I^2 = 77%), oral processing (k = 20, SMD = 0.78, 95% CI [0.50, 1.06], I^2 = 85%), auditory processing (k = 16, SMD = 1.40, 95% CI [0.95, 1.85], I^2 = 89%), and movement processing (k = 10, SMD = 1.28, 95% CI [0.56, 2.00], I^2 = 97%). There was no difference in body position processing between the ADHD and control groups (k = 4, SMD = 0.91, 95% CI [−0.16, 1.98], I^2 = 89%).

In subgroup analyses (Supplement 1, available online), differences between participants with ADHD and controls in tactile, visual, and auditory processing were found for children only. Movement and body position processing analyses did not include studies with an adult population. Adults and children with ADHD showed greater difficulties in oral processing compared with controls.

Heterogeneity was high in the different analyses. Its exploration of heterogeneity in subgroup analyses showed a significant difference based on risk of bias for visual processing ($p < .01$) and movement processing ($p = .02$) outcomes. When focusing on studies with a low risk of bias, the difference between groups concerning movement processing was no longer significant. Subgroup analyses showed a significant difference in heterogeneity based on sensory tools for visual processing ($p = .03$), auditory processing ($p = .01$), and movement processing ($p < .01$) outcomes. Meta-regression using age, year, or proportion of women as predictor was not significant for any of the sensory modality outcomes. No outliers were detected in leave-one-out

FIGURE 3 Forest Plot of Meta-analysis on Low Sensory Registration



Note: ADHD = attention-deficit/hyperactivity disorder, SMD = standardized mean difference. Please note color figures are available online.

sensitivity analysis. No publication bias was found in the different funnel plot visual inspections and Egger tests (Supplement 1, available online).

Meta-analyses: Binary Outcomes


Hypersensitivity. The meta-analysis on hypersensitivity included 9 studies (ADHD participants = 372, controls = 753). The ADHD group had an increased risk of hypersensitivity (OR = 9.0, 95% CI [2.92, 27.71]) in both children and adults. Heterogeneity was moderate (I^2 = 61%) (Figure S65, available online). The funnel plot suggested a publication bias (Figure S66, available online). The adjusted result with the trim-and-fill procedure still favored an increased risk of hypersensitivity in ADHD (OR = 7.73, 95% CI [2.57, 23.24]). We did not explore the heterogeneity for this outcome due to paucity of studies.

Hyposensitivity. The meta-analysis on hyposensitivity included 5 studies (ADHD participants = 284, controls = 664). The ADHD population had an increased risk of hyposensitivity (OR = 9.71, 95% CI [2.80, 33.61]). Heterogeneity was moderate (I^2 = 43%). This difference was observed only in adults in the subgroup analyses (Figure S67, available online). We did not explore heterogeneity and publication bias for this outcome given the limited number of studies.

DISCUSSION

This is the first systematic review and meta-analysis to our knowledge exploring the association between ADHD and atypical SPP. Using data from 31 studies, we found that compared with controls, individuals with ADHD had

FIGURE 4 Forest Plot of Meta-analysis on Sensory Seeking




Note: *ADHD = attention-deficit/hyperactivity disorder; SMD = standardized mean difference. Please note color figures are available online.*

significantly higher levels of atypical SPP in 4 sensory modulation domains (sensory sensitivity, sensory seeking, low registration, and sensory avoiding), although sensory avoiding was not different between groups after adjustment using the trim-and-fill procedure.

Sensory sensitivity and low registration were different between groups in both adults and children. Nevertheless, sensory seeking did not differ between the ADHD and control groups in the adult studies subgroup. This variability may be due to the inconsistent conceptualization of sensory

FIGURE 5 Forest Plot of Meta-analysis Results for the Sensory Modalities



Note: *The detailed forest plots for each sensory modality meta-analysis are presented in Supplement 1, available online. ADHD = attention-deficit/hyperactivity disorder; SMD = standardized mean difference. Please note color figures are available online.*

seeking across different studies. Scales for young children focus on intense sensory exploration, whereas adult questionnaires, such as AASP, address personal sensory preferences.[14] This also raises the question of overlap in the symptoms between SPP and ADHD. The higher atypical SPP in the ADHD group may be explained, at least for some domains, by this overlap.[23,62] For example, on the sensory profile, participants with a high level of sensory-seeking behaviors often touch objects and tap their pencils in a way that increases the sensory input and stays alert. This mechanism has been frequently described in ADHD (hyperactivity). Therefore, it may be difficult to differentiate hyperactivity symptoms from sensory seeking. As hyperactivity symptoms tend to decrease with age, it is also possible that adult patients with ADHD present with fewer sensory-seeking behaviors. However, our meta-regression on age was not significant for this outcome, making it difficult to draw conclusions.

Low registration was moderated by age, with an increasing difference between individuals with ADHD and controls as age increased. This result is somewhat surprising, as the opposite trend has been described for ASD.[14,63] As inattention could reduce the perception of sensory stimuli and increase low registration scores, one hypothesis could be that in the younger population, certain contextual factors, such as the presence of parents, may naturally enhance children's attention toward stimuli. Another hypothesis could be that due to a longer history of ADHD and therefore the development of adaptive strategies, adults could filter sensory stimuli more effectively, leading to an increase in the low-registration domain.

Concerning sensory modalities, compared with the control group, participants with ADHD had a higher level of atypical SPP in 4 sensory discrimination domains/modalities (tactile, visual, auditory, oral, and movement). No difference was found in body position processing (or proprioception); however, this analysis included only 3 studies, so further replication is needed.

Our results on sensory modalities can be explained by shared neurophysiological pathways between ADHD and SPP, particularly involving disruptions in sensory integration systems. Deficits in white matter integrity in key tracts responsible for sensory integration can contribute to multisensory impairments.[64] A recent meta-analysis described alterations in the splenium and the body of the corpus callosum in ADHD, and these alterations could contribute to sensory processing difficulties.[65] Additionally, dopamine dysregulation, a hallmark of ADHD, can impair sensory gating and filtering mechanisms, further contributing to sensory discrimination challenges. This hypothesis suggests that treatment with psychostimulants may offer potential benefits for regulating sensory processing.[66]

When separating studies on adult and child populations, no difference between participants with ADHD and controls was found in adults concerning tactile, visual, and auditory processing. However, only 2 studies analyzed these outcomes and presented high between-study heterogeneity. More studies are therefore needed to reach conclusions on these specific domains in adult populations.

Considering the binary outcomes, individuals with ADHD, either adults or children, had a higher risk of being hypersensitive and hyposensitive. These analyses had a lower level of heterogeneity. This result is not surprising given the differences between patients with ADHD and controls across various sensory quadrants. Furthermore, this binary outcome demonstrates that clear distinctions between the groups remain even when focusing solely on individuals with extreme deviations from the norm. This underscores that these hypersensitivity and hyposensitivity patterns in ADHD represent an important clinical issue, complementing the significant differences already observed in continuous sensory scores.

More generally, our findings illustrate the complexity of sensory processing difficulties, with some individuals exhibiting both hypersensitivity and hyposensitivity, depending on the specific sensory modality. For instance, a person may be hypersensitive to auditory stimuli and hyposensitive to tactile input. This coexistence and the atypical sensory patterns across the 4 sensory quadrants reflect the heterogeneity of sensory profiles in ADHD, where sensory processing difficulties are not uniformly distributed but manifest as unique combinations of strengths and challenges across modalities.

This systematic review and meta-analysis has several strengths. We included studies with a clinical diagnosis of ADHD rather than ADHD symptoms to ensure that the results were pertinent to this clinical population. We included all validated sensory processing questionnaires to enable generalizability. Our work should also be considered in light of some limitations.

Concerning the limitations of the primary studies included in our systematic review, substantial methodological heterogeneity was observed, and a few studies were rated as having a low risk of bias. We could not retrieve missing data for several studies even after contacting the corresponding authors, preventing us from processing all preplanned sensitivity analyses. The presence of ADHD or another neurodevelopmental disorder was not systematically verified in the control groups, and some studies did not assess comorbidities or ADHD medications. We were unable to test the impact of medication, severity of ADHD, and type of ADHD, although these factors could be potential moderators of sensory processing. Information on IQ

was frequently missing in 30% of the studies; however, no significant differences were observed between the subgroups in the IQ assessment subgroup analysis. The sample size of the included studies was often small and not defined a priori. Information on ethnicity was reported in only 5 studies (17%), which limits the generalizability of our results across diverse populations.

Most of our analyses found high heterogeneity, limiting our confidence in the results. Multiple subgroup and meta-regression analyses were conducted to explore this heterogeneity, but we did not find any specific explanation.

Concerning the limitations of the systematic review and meta-analysis itself, as stated above, we wanted to synthesize sensory processing constructs across multiple tools; therefore, we created a classification with an expert in the field. This classification could nevertheless introduce variability and bias the results beyond what we could identify in this meta-analysis.

Suspicion of publication bias emerged from the sensitivity analyses on several outcomes (sensory avoiding, low registration, hypersensitivity). We used an adjustment method to correct this bias.

The findings from this meta-analysis strongly indicate that both children and adults with ADHD are more likely to experience atypical SPP than the general population. Currently, the main guidelines for ADHD management (ie, National Institute for Health and Care Excellence [NICE] guidelines, American Academy of Pediatrics [AAP], American Academy of Child and Adolescent Psychiatry [AACAP] practice parameters[67-69]) do not include specific information on associated SPP. Given the significant functional impact of atypical SPP, early identification and targeted interventions are crucial to improve the overall prognosis of individuals with ADHD. Importantly, assessing SPP in individuals with ADHD is relatively straightforward due to the availability of validated assessment tools, making it easier for clinicians to identify and address these issues.

The effectiveness of sensory-based interventions for atypical SPP remains a subject of debate. Whereas some studies have suggested potential benefits, available studies present important methodological limitations, such as insufficient samples or nonrandomized assessment.[70] Lang *et al.*[71] reviewed sensory-based treatments and reported mixed results: 3 studies suggested effectiveness, 8 found mixed outcomes, and 14 found no significant benefits. Importantly, most of these studies evaluated isolated sensory procedures (eg, weighted vests, sensory brushing) rather than comprehensive, multicomponent approaches. Due to the current lack of solid evidence supporting direct therapeutic interventions, some in the field have highlighted the importance of applying sensory processing knowledge to make practical modifications in daily routines and environments, thereby enhancing accessibility and alleviating sensory-related difficulties.[72]

Crucially, the purpose of assessing SPP in individuals with ADHD is not solely to categorize them as normal or pathological. Instead, the goal is to better understand their unique functional profiles, including their strengths and weaknesses, and to leverage this understanding to develop comprehensive, individualized interventions. SPP assessments can provide clinicians and caregivers with valuable insights into how to support individuals by building on their strengths and addressing their sensory challenges in meaningful ways. Beyond identification, clinicians can guide patients in adapting their environments to better manage sensory sensitivities, such as using noise-reduction headphones for patients with high auditory sensitivity, as well as advocate for inclusive policies, such as quiet hours in supermarkets. Additionally, understanding one's sensory processing profile through such assessments can empower individuals with ADHD to engage in self-adaptation strategies, thereby enhancing their ability to cope with and mitigate the impact of SPD on their daily lives. By integrating these insights, clinicians may more effectively offer tailored interventions that could significantly enhance outcomes.

SPP has been proposed as a translational model in the Research Domain Criteria framework to better understand ASD.[73,74] These research efforts could be extended to ADHD, with SPP representing a novel avenue for advancing our understanding of the disorder and its associated comorbidities.

To fully realize the potential of this emerging field, future studies should prioritize larger, well-defined cohorts of ADHD and control participants, with careful attention paid to variables such as medication status, IQ, and comorbid conditions. Longitudinal studies, in particular, would be essential for examining the developmental trajectory of sensory processing in ADHD across the life span. Studies on the prevalence of atypical SPP could provide valuable insights into the extent and significance of sensory difficulties in ADHD. Additionally, studies comparing individuals with ADHD and other clinical groups would help to better understand the specificity of this association.

CRediT authorship contribution statement

Lucie Jurek: Writing – review & editing, Writing – original draft, Visualization, Project administration, Methodology, Formal analysis, Data curation, Conceptualization. **Arnaud Duchier:** Writing – review & editing, Data curation. **Christophe Gauld:** Writing – review & editing, Data

curation. **Léonie Hénault:** Writing – review & editing, Methodology. **Caroline Giroudon:** Writing – review & editing, Software, Data curation, Conceptualization. **Pierre Fourneret:** Writing – review & editing. **Samuele Cortese:** Writing – review & editing, Writing – original draft, Supervision. **Mikail Nourredine:** Writing – review & editing, Writing – original draft, Supervision, Methodology, Formal analysis, Conceptualization.

Accepted April 9, 2025.

[a]Centre Hospitalier Le Vinatier, Bron, France; [b]Université de Lyon 1, Lyon, France; [c]University of Southampton, Southampton, United Kingdom; [d]Hôpital Charles Perrens, Bordeaux, France; [e]Hôpital Femme Mère Enfant, Hospices Civils de Lyon, Lyon, France; [f]Hospices Civils de Lyon, Lyon, France; [g]Solent NHS Trust, Southampton, United Kingdom; [h]New York University Child Study Center, New York City, New York; [i]University of Bari Aldo Moro, Bari, Italy

The authors have reported no funding for this work.

Data Sharing: Data and code are available upon request to the corresponding author.

Disclosure: Lucie Jurek has received a research mobility grant from the Planiol Foundation. The Planiol Foundation was not involved in any part of this work. Pierre Fourneret has received honoraria from HAC Pharma. Samuele Cortese, NIHR Research Professor (NIHR303122), is funded by the NIHR for this research project. The views expressed in this publication are those of the author(s) and not necessarily those of the NIHR, NHS, or the UK Department of Health and Social Care. He is also supported by NIHR grants NIHR203684, NIHR203035, NIHR130077, NIHR128472, and RP-PG-0618-20003 and by grant 101095568-HORIZONHLTH-2022-DISEASE-07-03 from the European Research Executive Agency. He has declared reimbursement for travel and accommodation expenses from the Association for Child and Adolescent Mental Health (ACAMH) in relation to lectures delivered for ACAMH, the Canadian AADHD Alliance Resource, the British Association of Psychopharmacology, and Healthcare Convention for educational activity on ADHD and has received honoraria from Medice. Arnaud Duchier, Christophe Gauld, Léonie Hénault, Caroline Giroudon, and Mikail Nourredine have reported no biomedical financial interests or potential conflicts of interest.

*Correspondence to Lucie Jurek, MD, PhD, CETESLA, bat 504, CH Le Vinatier, 69500 Bron, France; e-mail: lucie.jurek@ch-le-vinatier.fr

0890-8567/$36.00/©2025 American Academy of Child and Adolescent Psychiatry. Published by Elsevier Inc. This is an open access article under the CC BY-NC-ND license (http://creativecommons.org/licenses/by-nc-nd/4.0/).

https://doi.org/10.1016/j.jaac.2025.02.019

REFERENCES

1. American Psychiatric Association. Diagnostic and Statistical Manual of Mental Disorders. 5th ed, text rev. Washington, DC: American Psychiatric Association; 2022.
2. Cortese S, Song M, Farhat LC, *et al*. Incidence, prevalence, and global burden of ADHD from 1990 to 2019 across 204 countries: data, with critical re-analysis, from the Global Burden of Disease study. Mol Psychiatry. 2023;28(11):4823-4830. https://doi.org/10.1038/s41380-023-02228-3
3. Ayano G, Tsegay L, Gizachew Y, *et al*. Prevalence of attention deficit hyperactivity disorder in adults: umbrella review of evidence generated across the globe. Psychiatry Res. 2023;328:115449. https://doi.org/10.1016/j.psychres.2023.115449
4. Arrondo G, Solmi M, Dragioti E, *et al*. Associations between mental and physical conditions in children and adolescents: an umbrella review. Neurosci Biobehav Rev. 2022;137:104662. https://doi.org/10.1016/j.neubiorev.2022.104662
5. Faraone SV, Asherson P, Banaschewski T, *et al*. Attention-deficit/hyperactivity disorder. Nat Rev Dis Primers. 2015;1(1):15020. https://doi.org/10.1038/nrdp.2015.20
6. Ayres AJ. Characteristics of types of sensory integrative dysfunction. Am J Occup Ther. 1971;25(7):329-334.
7. Neufeld J, Hederos Eriksson L, Hammarsten R, *et al*. The impact of atypical sensory processing on adaptive functioning within and beyond autism: the role of familial factors. Autism. 2021;25(8):2341-2355. https://doi.org/10.1177/13623613211019852
8. Miller L. Perspectives on sensory processing disorder: a call for translational research. Front Integr Neurosci. 2009;3:22. https://doi.org/10.3389/neuro.07.022.2009
9. Dunn W. The sensations of everyday life: empirical, theoretical, and pragmatic considerations. Am J Occup Ther. 2001;55(6):608-620. https://doi.org/10.5014/ajot.55.6.608
10. Ben-Sasson A, Hen L, Fluss R, Cermak SA, Engel-Yeger B, Gal E. A meta-analysis of sensory modulation symptoms in individuals with autism spectrum disorders. J Autism Dev Disord. 2009;39(1):1-11. https://doi.org/10.1007/s10803-008-0593-3
11. Gouze KR, Hopkins J, LeBailly SA, Lavigne JV. Re-examining the epidemiology of sensory regulation dysfunction and comorbid psychopathology. J Abnorm Child Psychol. 2009;37(8):1077-1087. https://doi.org/10.1007/s10802-009-9333-1
12. Ahn RR, Miller LJ, Milberger S, McIntosh DN. Prevalence of parents' perceptions of sensory processing disorders among kindergarten children. Am J Occup Ther. 2004;58(3):287-293. https://doi.org/10.5014/ajot.58.3.287
13. Baranek GT, Foster LG, Berkson G. Sensory defensiveness in persons with developmental disabilities. Occup Ther J Res. 1997;17(3):173-185. https://doi.org/10.1177/153944929701700302
14. Ben-Sasson A, Gal E, Fluss R, Katz-Zetler N, Cermak SA. Update of a meta-analysis of sensory symptoms in ASD: a new decade of research. J Autism Dev Disord. 2019;49(12):4974-4996. https://doi.org/10.1007/s10803-019-04180-0
15. Cosbey J, Johnston SS, Dunn ML. Sensory processing disorders and social participation. Am J Occup Ther. 2010;64(3):462-473. https://doi.org/10.5014/ajot.2010.09076
16. DelCarmen-Wiggins R, Carter A. Handbook of Infant, Toddler, and Preschool Mental Health Assessment. New York, NY: Oxford University Press; 2004.
17. Zeanah CH, Carter AS, Cohen J, *et al*. Introducing a New Classification of Early Childhood Disorders: DC:0–5. 2017. Accessed October 1, 2024. https://mi-aimh.org/wp-content/uploads/2017/11/2017-01-Zeanah-S.-intro-article.pdf
18. Yochman A, Parush S, Ornoy A. Responses of preschool children with and without ADHD to sensory events in daily life. Am J Occup Ther. 2004;58(3):294-302. https://doi.org/10.5014/ajot.58.3.294
19. Dunn W, Bennett D. Patterns of sensory processing in children with attention deficit hyperactivity disorder. Occupational Therapy Journal of Research. 2002;22(1):4-15. https://doi.org/10.1177/153944920202200102
20. Page MJ, McKenzie JE, Bossuyt PM, *et al*. The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. BMJ. 2021;372:n71. https://doi.org/10.1136/bmj.n71
21. Fuermaier ABM, Hüpen P, De Vries SM, *et al*. Perception in attention deficit hyperactivity disorder. Atten Defic Hyperact Disord. 2018;10(1):21-47. https://doi.org/10.1007/s12402-017-0230-0
22. Ghanizadeh A. Sensory processing problems in children with ADHD, a systematic review. Psychiatry Investig. 2011;8(2):89-94. https://doi.org/10.4306/pi.2011.8.2.89
23. Keating J, Gaffney R, Bramham J, Downes M. Sensory modulation difficulties and assessment in children with attention deficit hyperactivity disorder: a systematic review. Eur J Dev Psychol. 2022;19(1):110-144. https://doi.org/10.1080/17405629.2021.1889502
24. Van Den Boogert F, Klein K, Spaan P, *et al*. Sensory processing difficulties in psychiatric disorders: a meta-analysis. J Psychiatr Res. 2022;151:173-180. https://doi.org/10.1016/j.jpsychires.2022.04.020
25. Ribeiro CM, Beserra BTS, Silva NG, *et al*. Exposure to endocrine-disrupting chemicals and anthropometric measures of obesity: a systematic review and meta-analysis. BMJ Open. 2020;10(6):e033509. https://doi.org/10.1136/bmjopen-2019-033509
26. Martínez-González AE, Rodríguez-Jiménez T, Piqueras JA, *et al*. Cross-disorder comparison of sensory reactivity, pain, gastro-intestinal symptoms and obsessive-compulsive symptoms in adolescents and young adults with autism and other neurodevelopmental disorders. Int J Dev Disabil. Published online February 28, 2024. https://doi.org/10.1080/20473869.2024.2318529
27. Falkenstein MJ, Conelea CA, Garner LE, Haaga DAF. Sensory over-responsivity in trichotillomania (hair-pulling disorder). Psychiatry Res. 2018;260:207-218. https://doi.org/10.1016/j.psychres.2017.11.034
28. Higgins JPT, Thomas J, Chandler J, *et al*. Cochrane Handbook for Systematic Reviews of Interventions Version 6.4 (updated August 2023). Cochrane; 2023. Accessed October 1, 2024. www.training.cochrane.org/handbook
29. IntHout J, Ioannidis JP, Borm GF. The Hartung-Knapp-Sidik-Jonkman method for random effects meta-analysis is straightforward and considerably outperforms the standard DerSimonian-Laird method. BMC Med Res Methodol. 2014;14(1):25. https://doi.org/10.1186/1471-2288-14-25

30. Thornton A. Publication bias in meta-analysis its causes and consequences. J Clin Epidemiol. 2000;53(2):207-216. https://doi.org/10.1016/S0895-4356(99)00161-4
31. Balduzzi S, Rücker G, Schwarzer G. How to perform a meta-analysis with R: a practical tutorial. Evid Based Mental Health. 2019;22(4):153-160. https://doi.org/10.1136/ebmental-2019-300117
32. R Core Team. R: A Language and Environment for Statistical Computing. R Foundation for Statistical Computing, Vienna, Austria. Accessed January 18, 2025. https://www.R-project.org/
33. Mangeot SD, Miller LJ, McIntosh DN, *et al*. Sensory modulation dysfunction in children with attention-deficit-hyperactivity disorder. Dev Med Child Neurol. 2001;43(6):399-406. https://doi.org/10.1111/j.1469-8749.2001.tb00228.x
34. Bröring T, Rommelse N, Sergeant J, Scherder E. Sex differences in tactile defensiveness in children with ADHD and their siblings. Dev Med Child Neurol. 2008;50(2):129-133. https://doi.org/10.1111/j.1469-8749.2007.02024.x
35. Cheung PPP, Siu AMH. A comparison of patterns of sensory processing in children with and without developmental disabilities. Res Dev Disabil. 2009;30(6):1468-1480. https://doi.org/10.1016/j.ridd.2009.07.009
36. Miller LJ, Nielsen DM, Schoen SA. Attention deficit hyperactivity disorder and sensory modulation disorder: a comparison of behavior and physiology. Res Dev Disabil. 2012;33(3):804-818. https://doi.org/10.1016/j.ridd.2011.12.005
37. Lin CY, Yang AL, Su CT. Objective measurement of weekly physical activity and sensory modulation problems in children with attention deficit hyperactivity disorder. Res Dev Disabil. 2013;34(10):3477-3486. https://doi.org/10.1016/j.ridd.2013.07.021
38. Lufi D, Tzischinsky O. The relationships between sensory modulation and sleep among adolescents with ADHD. J Atten Disord. 2014;18(8):646-653. https://doi.org/10.1177/1087054712457036
39. Shimizu VT, Bueno OFA, Miranda MC. Sensory processing abilities of children with ADHD. Braz J Phys Ther. 2014;18(4):343-352. https://doi.org/10.1590/bjpt-rbf.2014.0043
40. Pfeiffer B, Daly BP, Nicholls EG, Gullo DF. Assessing sensory processing problems in children with and without attention deficit hyperactivity disorder. Phys Occup Ther Pediatr. 2015;35(1):1-12. https://doi.org/10.3109/01942638.2014.904471
41. Bijlenga D, Tjon-Ka-Jie JYM, Schuijers F, Kooij JJS. Atypical sensory profiles as core features of adult ADHD, irrespective of autistic symptoms. Eur Psychiatry. 2017;43:51-57. https://doi.org/10.1016/j.eurpsy.2017.02.481
42. Sanz-Cervera P, Pastor-Cerezuela G, González-Sala F, Tárraga-Mínguez R, Fernández-Andrés MI. Sensory processing in children with autism spectrum disorder and/or attention deficit hyperactivity disorder in the home and classroom contexts. Front Psychol. 2017;8:1772. https://doi.org/10.3389/fpsyg.2017.01772
43. Little LM, Dean E, Tomchek S, Dunn W. Sensory processing patterns in autism, attention deficit hyperactivity disorder, and typical development. Phys Occup Ther Pediatr. 2018;38(3):243-254. https://doi.org/10.1080/01942638.2017.1390809
44. Mimouni-Bloch A. Association between sensory modulation and daily activity function of children with attention deficit/hyperactivity disorder and children with typical development. Res Dev Disabil. 2018;83:69-76. https://doi.org/10.1016/j.ridd.2018.08.002
45. Jegadeesan T, Nagalakshmi P, Renuchitra R. Developing children and children with attention deficit. Indian J Public Health Res Dev. 2020;11(8):167-173.
46. Kamath MS, Dahm CR, Tucker JR, Huang-Pollock CL, Etter NM, Neely KA. Sensory profiles in adults with and without ADHD. Res Dev Disabil. 2020;104:103696. https://doi.org/10.1016/j.ridd.2020.103696
47. Ohta H, Aoki YY, Itahashi T, *et al*. White matter alterations in autism spectrum disorder and attention-deficit/hyperactivity disorder in relation to sensory profile. Molecular Autism. 2020;11(1):77. https://doi.org/10.1186/s13229-020-00379-6
48. Adra N, Cao A, Makris N, Valera EM. Sensory modulation disorder and its neural circuitry in adults with ADHD: a pilot study. Brain Imaging Behav. 2021;15(2):930-940. https://doi.org/10.1007/s11682-020-00302-w
49. Dellapiazza F, Michelon C, Vernhet C, *et al*. Sensory processing related to attention in children with ASD, ADHD, or typical development: results from the ELENA cohort. Eur Child Adolesc Psychiatry. 2021;30(2):283-291. https://doi.org/10.1007/s00787-020-01516-5
50. Gomez IN, Domondon LM, Tsang HW, Chan CC, Lai CY. Sensory behaviours and resting parasympathetic functions among children with and without ADHD. ScientificWorldJournal. 2021;2021:6615836. https://doi.org/10.1155/2021/6615836
51. Grinblat N, Rosenblum S. Work participation, sensory processing and sleep quality in adults with attention-deficit hyperactive disorder. Work. 2022;73(4):1235-1244. https://doi.org/10.3233/WOR-211129
52. Sobhy N, Afsah O, Baz H. Developing an Arabic questionnaire to assess sensory processing disorders among preschool Egyptian children. Res Dev Disabil. 2022;126:104238. https://doi.org/10.1016/j.ridd.2022.104238
53. Darnal S, Panda S, Namrata Chettri A. Sensory processing, emotional and behavioural challenges in children with autism spectrum disorder and attention deficit hyperactive disorder. Int J Dev Disabil. Published online December 18, 2023. https://doi.org/10.1080/20473869.2023.2277601
54. Panda PK, Ramachandran A, Kumar V, Sharawat IK. Sensory processing abilities and their impact on disease severity in children with attention-deficit hyperactivity disorder. J Neurosci Rural Pract. 2023;14(3):509-515. https://doi.org/10.25259/JNRP_22_2023
55. Rani I, Agarwal V, Arya A, Mahour P. Sensory processing in children and adolescents with attention deficit hyperactivity disorder. J Atten Disord. 2023;27(2):145-151. https://doi.org/10.1177/10870547221129306
56. Schramm M, Goregliad Fjaellingsdal T, Aslan B, *et al*. Electrophysiological evidence for increased auditory crossmodal activity in adult ADHD. Front Neurosci. 2023;17:1227767. https://doi.org/10.3389/fnins.2023.1227767
57. Schulz SE, Kelley E, Anagnostou E, *et al*. Sensory processing patterns predict problem behaviours in autism spectrum disorder and attention-deficit/hyperactivity disorder. Adv Neurodev Disord. 2023;7(1):46-58. https://doi.org/10.1007/s41252-022-00269-3
58. Fabio RA, Orsino C, Lecciso F, Levante A, Suriano R. Atypical sensory processing in adolescents with attention deficit hyperactivity disorder: a comparative study. Res Dev Disabil. 2024;146:104674. https://doi.org/10.1016/j.ridd.2024.104674
59. Mohammed HO, Elsayed SAA, Mahmoud NF. Sensory processing profile among a sample of Egyptian children with different types of delayed language development: correlations of different variables. Egypt J Otolaryngol. 2024;40(1):20. https://doi.org/10.1186/s43163-023-00554-5
60. Grinblat N, Rosenblum S. Sleep quality, sensory processing abilities and work performance for adults with attention deficit hyperactive disorder. Eur Psychiatry. 2021;64(Suppl 1):S553. https://doi.org/10.1192/j.eurpsy.2021.1475
61. Mimouni-Bloch A, Offek H, Rosenblum S, Posener E, Silman Z, Engel-Yeger B. Association between sensory processing disorder and daily function of children with attention deficit/hyperactive disorder and controls. Eur J Paediatr Neurol. 2017;21(Suppl 1):e171. https://doi.org/10.1016/j.ejpn.2017.04.1157
62. Koziol LF, Budding D. ADHD and sensory processing disorders: placing the diagnostic issues in context. Appl Neuropsychol Child. 2012;1(2):137-144. https://doi.org/10.1080/21622965.2012.709422
63. Schauder KB, Bennetto L. Toward an interdisciplinary understanding of sensory dysfunction in autism spectrum disorder: an integration of the neural and symptom literatures. Front Neurosci. 2016;10:268. https://doi.org/10.3389/fnins.2016.00268
64. Narayan A, Rowe MA, Palacios EM, *et al*. Altered cerebellar white matter in sensory processing dysfunction is associated with impaired multisensory integration and attention. Front Psychol. 2021;11:618436. https://doi.org/10.3389/fpsyg.2020.618436
65. Parlatini V, Itahashi T, Lee Y, *et al*. White matter alterations in attention-deficit/hyperactivity disorder (ADHD): a systematic review of 129 diffusion imaging studies with meta-analysis. Mol Psychiatry. 2023;28(10):4098-4123. https://doi.org/10.1038/s41380-023-02173-1
66. Sommer JL, Low AM, Jepsen JRM, *et al*. Effects of methylphenidate on sensory and sensorimotor gating of initially psychostimulant-naïve adult ADHD patients. Eur Neuropsychopharmacol. 2021;46:83-92. https://doi.org/10.1016/j.euroneuro.2021.02.004
67. Wolraich ML, Hagan JF, Allan C, *et al*. Clinical Practice Guideline for the Diagnosis, Evaluation, and Treatment of Attention-Deficit/Hyperactivity Disorder in Children and Adolescents. Pediatrics. 2019;144(4):e20192528. https://doi.org/10.1542/peds.2019-2528
68. Pliszka S. Practice Parameter for the Assessment and Treatment of Children and Adolescents With Attention-Deficit/Hyperactivity Disorder. J Am Acad Child Adolesc Psychiatry. 2007;46(7):894-921. https://doi.org/10.1097/chi.0b013e318054e724
69. Attention Deficit Hyperactivity Disorder: Diagnosis and Management. London: National Institute for Health and Care Excellence (NICE); 2019. (NICE Guideline, No. 87.). Accessed October 1, 2024. https://www.ncbi.nlm.nih.gov/books/NBK493361/
70. Camarata S, Miller LJ, Wallace MT. Evaluating sensory integration/sensory processing treatment: issues and analysis. Front Integr Neurosci. 2020;14:556660. https://doi.org/10.3389/fnint.2020.556660
71. Lang R, O'Reilly M, Healy O, *et al*. Sensory integration therapy for autism spectrum disorders: a systematic review. Res Autism Spectr Disord. 2012;6(3):1004-1018. https://doi.org/10.1016/j.rasd.2012.01.006
72. Bailliard A, Dunn W, Brown C, Engel-Yeger B. Editorial: Meaningful participation and sensory processing. Front Psychol. 2022;13:1045650. https://doi.org/10.3389/fpsyg.2022.1045650
73. Robertson CE, Baron-Cohen S. Sensory perception in autism. Nat Rev Neurosci. 2017;18(11):671-684. https://doi.org/10.1038/nrn.2017.112
74. Siemann JK, Veenstra-VanderWeele J, Wallace MT. Approaches to understanding multisensory dysfunction in autism spectrum disorder. Autism Res. 2020;13(9):1430-1449. https://doi.org/10.1002/aur.2375

Compendium of WHO and other UN guidance on health and environment

2022 update

Chapter 11. Environmental noise



WHO/HEP/ECH/EHD/22.01

© World Health Organization 2022

Some rights reserved. This work is available under the Creative Commons Attribution-NonCommercial-ShareAlike 3.0 IGO licence (CC BY-NC-SA 3.0 IGO; https://creativecommons.org/licenses/by-nc-sa/3.0/igo).

Suggested citation. Environmental noise. In: Compendium of WHO and other UN guidance on health and environment, 2022 update. Geneva: World Health Organization; 2022 (WHO/HEP/ECH/EHD/22.01). Licence: CC BY-NC-SA 3.0 IGO.

11. Environmental noise

This section addresses exposure of the general population to environmental noise, such as noise from various forms of traffic or industry. It includes amplified music in the framework of leisure activities as well. It does not specifically include occupational noise exposure. Occupational risks, including noise exposure, are covered in section 11.3 on workplaces.

Overview

In 2011, an estimated one million healthy life years were lost from traffic-related noise in the western part of Europe only *(1)*. Important sources for environmental noise exposure are road, railway and air traffic, or building sites. Noise exposure can also occur through other sources such as wind turbines, and leisure activities such as listening to loud music or other audio content including participation in e-sports (video and computer game competitions). Excessive noise can cause annoyance; in addition research shows it increases the risk for IHD and hypertension, sleep disturbance, hearing impairment, tinnitus and cognitive impairment, with increasing evidence for other health impacts such as adverse birth outcomes and mental health problems *(2)*.

What is the proportion of people impacted by environmental noise in my country?

The noise indicators below are taken from guidelines that were developed for the WHO European Region. In terms of their health implications, the recommended exposure levels can be considered applicable in other regions and suitable for a global audience *(2)*.

Noise indicators are based on the European Union Directive 2002/49/EC *(3)* in the European Region.

- L_{den} is an average sound pressure level over all days, evenings and nights in a year.
- L_{night} is the equivalent continuous sound pressure level when the reference time interval is the night.
- $L_{Aeq,T}$ is the A-weighted (a frequency weighting to better reflect the human ear), equivalent continuous sound pressure level during a stated time interval starting at t_1 and ending at t_2, expressed in decibels (dB), at a given point in space.

The first two indicators are used particularly for noise monitoring and exposure assessment. The third is used for measuring leisure noise exposure. For more information on these and other noise indicators consult the *Environmental noise guidelines for the European Region (2)*. These noise indicators can be converted to other indicators used in other settings *(4)*.

What is the proportion of people impacted by environmental noise in my country?

Several countries use surveys to assess the perception of noise in the general population. The last European Quality of Life survey, carried out 2016–2017, found that 32% of more than 30 000 participants across Europe reported problems with noise in the immediate neighbourhood of their home *(5)*.

What are the levels of noise exposure we want to achieve?

Based on the systematic review of evidence available at the time of the development of the environmental noise guidelines *(2)*, the following recommended levels for specific noise sources can be defined.

For average noise exposure, the following sound pressure levels are recommended *(2, 6)*:

- < 53 dB L_{den} for road traffic noise
- < 54 dB L_{den} for railway noise
- < 45 dB L_{den} for aircraft noise
- < 45 dB L_{den} for wind turbine noise
- yearly average from all leisure source noises combined to ≤ 70 dB $L_{Aeq, 24h}$
- weekly average from leisure sources (such as personal listening devices [1]) ≤ 80 dB(A) or 1.6 Pa^2h
- short-term average from occasional exposure to leisure source noise ≤ 100 dB $L_{Aeq, 15min}$.

For **night noise exposure**, the following sound pressure levels are recommended *(2)*:

- < 45 dB L_{night} for road traffic noise
- < 44 dB L_{night} for railway noise
- < 40 dB L_{night} for aircraft noise.

Different categories of noise mitigation interventions along a continuum from source reduction to behaviour change can be defined. Interventions in the guidance section below are marked with A–E as defined hereafter *(2)*.

A. Source intervention:
   - change in emission levels of sources
   - time restrictions on source operators.

B. Path intervention:
   - change in the path between source and receiver
   - path control through insulation of receiver/receiver's dwelling

C. New/closed infrastructure:
   - opening of a new infrastructure noise source
   - closure of an existing one
   - planning controls between (new) receivers and sources.

D. Other physical intervention:
   - change in other physical dimensions of dwelling/neighbourhood.

E. Behaviour change intervention:
   - change in individual behaviour to reduce exposure
   - avoidance of exposure or reduced duration of exposure
   - community education and communication.

[1] A personal listening or audio device is a portable device designed to be worn on the body or in a pocket. It is designed to allow the user to listen to various forms of media.

Guidance	Sector principally involved in planning/ implementation	Level of implementation	Instruments
Road traffic noise: policies and actions			
Recommended actions are available for specific noise sources and do not cover all potentially important noise exposures.			
1. Improve the choice of appropriate tyres and road surface (A) *(2)*.	Transport Land use planning	National; community	Regulation; infrastructure, technology and built environment
2. Reduce traffic flow and restrict truck traffic (A) *(2)*.	Transport Land use planning	National; community	Regulation; taxes and subsidies; infrastructure, technology and built environment
3. Insulate dwellings, construct barriers (B) *(2)*.	Housing Land use planning	National; community	Regulation; taxes and subsidies; infrastructure, technology and built environment
4. Construct road tunnels (C) *(2)*.	Transport Land use planning	National; community	Infrastructure, technology and built environment
5. Design/make available a “quiet side” in the dwelling; create nearby green space (D) *(2)*.	Housing Land use planning	National; community	Infrastructure, technology and built environment
Railway noise: policies and actions			
6. Apply rail grinding procedures to remove deformations and corrosions on railway tracks (A) *(2)*.	Transport	National; community	Infrastructure, technology and built environment
Railway noise: awareness raising and capacity building			
7. Inform the community about interventions being implemented to potentially reduce noise annoyance (E) *(2)*.	Health Environment Transport	Community Universal health coverage	Information, education and communication
Aircraft noise: policies and action			
8. Adapt opening and closing of runways (C) *(2)*.	Transport	National; community	Regulation; other management and control

Guidance	Sector principally involved in planning/ implementation	Level of implementation	Instruments
9. Rearrange flight paths (C) *(2)*.	Transport	National; community	Regulation; other management and control
Leisure noise: policies and actions			
10. Implement sound exposure monitoring (volume level and time spent listening) in all personal listening devices to allow for self-control with reference to a standard. In every listening device, the user should be allowed to select two different operational modes of reference exposure *(6)*, and track the percentage of exposure used vs the reference exposure for every seven days. The two operational modes include the following. • Mode 1: WHO standard level for adults • Mode 2: WHO standard level for sensitive users (e.g. children).	Health Sports and leisure	National	Regulation; infrastructure, technology and built environment
11. Implement options for volume limitation and parental volume control in every device *(6)*.	Sports and leisure	National	Regulation; infrastructure, technology and built environment
12. Enact and enforce legislation/regulations/policies for limiting sound levels and exposure in entertainment venues and events such as clubs, bars, fitness centres, concerts, etc.*(3, 7)*. Legislation should focus on: • limiting sounds to 100 dB(A) averaged over 15 minutes; • conducting regular sound monitoring to ensure and document compliance; • optimizing venue acoustics and sound system design to ascertain optimal listening conditions for all audience members in the venue/event; • create quiet zones allowing audience members to rest; • ensuring provision of hearing protection (earplugs); • ensuring provision of training on noise reduction strategies and information about noise.	Sports and leisure	National	Regulation
Leisure noise: awareness raising and capacity building			
13. Provide information on personal sound exposure to the user of personal listening devices through the device interface or other means *(6)*.	Health Sports and leisure	National	Information, education and communication
14. Provide personalized recommendations and cues for action for safe listening through personal listening devices, customized to a user's listening profile through the device interface or other means *(6)*.	Health Sports and leisure	National	Information, education and communication
15. Provide instructions on how to use safe listening features on the specific device through the device interface or other means *(6)*.	Health Sports and leisure	National	Information, education and communication

Guidance	Sector principally involved in planning/ implementation	Level of implementation	Instruments
16. Provide general information on safe listening and ways to practise it through the device interface or other means *(6)*.	Health Sports and leisure	National	Information, education and communication

Selected tools
WHO 2021: WHO is developing a global standard for safe listening entertainment venues *(7)* This guidance will promote safe listening among attendees of entertainment venues to mitigate their risk of hearing loss.
WHO Regional Office for Europe 2018: *Environmental noise guidelines for the European Region (2)* Results of the noise guidelines are also available as an executive summary in different languages.
WHO/ITU 2019: *Safe listening devices and systems – a WHO-ITU standard (6)* This document outlines the key features and requirements that personal audio systems must have in order to facilitate safe listening practices among users.
WHO 2015: *Make listening safe (8)* This webpage provides access to advocacy material around safe listening such as infographic, poster, banner and brochure.
WHO/ITU 2019: *Toolkit for safe listening devices and systems (9)* This toolkit provides practical guidance to support countries, industry partners and civil society groups in the use and implementation of the global standard on safe listening devices and systems (ITU-T H.870) *(10)*.
WHO Regional Office for Europe 2012: *Methodological guidance for estimating the burden of disease from environmental noise (11)*
WHO Regional Office for Europe 2011: *Burden of disease from environmental noise. Quantification of healthy life years lost in Europe (1)*
WHO Regional Office for Europe 2009: *Night noise guidelines for Europe (12)*

References

1. Burden of disease from environmental noise: Quantification of healthy life years lost in Europe. Copenhagen: WHO Regional Office for Europe; 2011 (https://apps.who.int/iris/handle/10665/326424).
2. Environmental noise guidelines for the European Region. Copenhagen: WHO Regional Office for Europe; 2018 (https://apps.who.int/iris/handle/10665/279952).
3. Directive 2002/49/EC of the European Parliament and of the Council of 25 June 2002 relating to the assessment and management of environmental noise. EUR-Lex; 2002 (https://eur-lex.europa.eu/legal-content/EN/TXT/?uri=CELEX%3A02002L0049-20210729).
4. Brink M, Schäffer B, Pieren R, Wunderli JM. Conversion between noise exposure indicators $Leq_{24}h$, $L_{Evening}$, L_{Night}, L_{dn} and L_{den} principles and practical guidance. Int J Hyg Environ Health. 2018;221(1):54–63. doi:10.1016/j.ijheh.2017.10.003.
5. Eurofound. European Quality of Life Survey 2016: quality of life, quality of public services, and quality of society. Luxembourg: Publications Office of the European Union; 2017 (https://www.eurofound.europa.eu/surveys/european-quality-of-life-surveys/european-quality-of-life-survey-2016).
6. Safe listening devices and systems: a WHO-ITU standard. Geneva: World Health Organization and International Telecommunication Union; 2019 (https://www.who.int/publications/i/item/safe-listening-devices-and-systems-a-who-itu-standard, accessed 2 December 2021).
7. Information session on the upcoming WHO Global standard for safe listening entertainment venues [website]. Geneva: World Health Organization; 2021 (https://www.who.int/news-room/events/detail/2021/09/21/default-calendar/information-session-on-the-upcoming-who-global-standard-for-safe-listening-entertainment-venues, accessed 10 December 2021).
8. Making listening safe [website]. Geneva: World Health Organization; 2021 (https://www.who.int/activities/making-listening-safe, accessed 2 December 2021).
9. Toolkit for safe listening devices and systems. Geneva: World Health Organization and International Telecommunication Union; 2019 (https://www.itu.int/en/ITU-D/Digital-Inclusion/Pages/Digital_Inclusion_Resources/Strategies,%20policies,%20toolkits/Toolkit_safe_listening_devices/safe_listening.aspx, accessed 2 December 2021).
10. Guidelines for safe listening devices/systems. In: SERIES H: AUDIOVISUAL AND MULTIMEDIA SYSTEMS: E-health multimedia systems, services and applications – safe listening. Geneva: International Telecommunication Union; 2018 (https://www.itu.int/rec/T-REC-H.870-201808-I/en, accessed 2 December 2021).
11. Hellmuth T, Classen T, Kim R, Kephalopoulos S, editors. Methodological guidance for estimating the burden of disease from environmental noise. Copenhagen: WHO Regional Office for Europe; 2012 (https://www.euro.who.int/__data/assets/pdf_file/0008/179117/Methodological-guidance-for-estimating-the-burden-of-disease-from-environmental-noise-ver-2.pdf, accessed 2 December 2021).
12. Night noise guidelines for Europe. Copenhagen: WHO Regional Office for Europe; 2009 (https://www.euro.who.int/__data/assets/pdf_file/0017/43316/E92845.pdf).

World Health Organization
Department of Environment, Climate Change and Health
Division of Universal Health Coverage / Healthier Populations
20, Avenue Appia
CH-1211 Geneva 27
Switzerland

www.who.int/teams/environment-climate-change-and-health