

# Puget Soundkeeper

Please see attached comments on Rulemaking - Aquatic Life Toxics Criteria, Chapter 173-201A WAC submitted on behalf of Puget Soundkeeper.

Thank you!  
Emily Gonzalez



May 7, 2024

Marla Koberstein  
Department of Ecology  
Water Quality Program  
PO Box 47696  
Olympia, WA 98504-7696

Protecting and  
Preserving  
Puget Sound

*Delivered via Ecology Online Portal*

130 Nickerson Street,  
Suite 107  
Seattle, WA 98109

**Re: Toxics Criteria for Aquatic Life - Comments**

Dear Ms Koberstein:

P 206.297.7002

F 206.297.0409

[www.pugetsoundkeeper.org](http://www.pugetsoundkeeper.org)

Puget Soundkeeper (Soundkeeper) respectfully submits these comments on the Department of Ecology's (Ecology) proposed updates to the Toxics Criteria for Aquatic Life. Soundkeeper also incorporates by reference herein comments submitted by Washington Council of Trout Unlimited regarding 6PPD-quinone. Soundkeeper has been working for decades to enhance and protect surface waters across the state burdened by a host of impacts, including toxic pollution and the effects of toxicity on the communities and ecosystems of the Puget Sound watershed. Soundkeeper in general supports the long-overdue update proposed by Ecology; our understanding of aquatic ecosystems has evolved significantly in the three decades since these criteria were last amended. In several key areas, however, more needs to be done.

#### I. Overall Update

First, Soundkeeper applauds Ecology for prioritizing this update to the broad list of criteria for toxicants found in Table 240. The role these criteria play in developing water quality standards cannot be overstated. Water quality standards are a fundamental and scientifically sound part of how pollution is managed and ecosystems are protected. They also provide a basis for how entities understand the practical expectations of regulatory compliance. These standards and expectations are relied upon by communities in an ever-evolving scientific landscape, but environmental regulations anticipate accommodating, in real-time or at the very least every few years, advances in science and technology. Thus, regular, periodic updates are both necessary and required. Soundkeeper is especially pleased with the proposed updated criteria where Washington has leaned into its role as a leader in toxics regulation.

Washington's Toxic Criteria for Aquatic Life is administered according to WAC 173-201A-240 and states "[t]he department may revise the criteria in Table 240 for aquatic life on a statewide or water body-specific basis *as needed to protect aquatic life* occurring in waters of the state and to *increase the technical accuracy of the criteria* being applied."<sup>1</sup> Significantly, the most recent large-scale update to the Aquatic Life Toxics Criteria occurred 31 years ago in 1993.<sup>2</sup> For perspective, notable scientific, political, and cultural events from 1993 include: the European Union was created, first female US Attorney General, the release of *Sleepless* in Seattle, Pearl Jam's *Jeremy* won MTV's Best Video award, the *London Convention* banned

<sup>1</sup> WAC 173-201A-240(5)(a), emphasis added.

<sup>2</sup> See Focus Sheet, available at <https://apps.ecology.wa.gov/publications/documents/2310040.pdf>

dumping radioactive waste at sea, NAFTA was signed into law, the Convention on Biological Diversity went into effect, and the planet resolved the hole in the ozone layer of our atmosphere. Things change over 31 years, and Soundkeeper asserts that there must not be this kind of gap between updates to regulatory schema like the toxics criteria which are meant to respect and rely on up-to-date knowledge.

## II. Proposed 6PPD-quinone Acute Freshwater Criteria

Soundkeeper has a particular interest in stormwater and wastewater management, 6PPD, and the derivative 6PPD-quinone (6PPD-Q). Accordingly, we are very supportive of Ecology's proposed criteria for 6PPD-Q in stormwater. Finalization of this criteria will be invaluable as we move forward in trying to regulate, monitor, and address this pollutant in runoff, stormwater, combined sewage overflows, and other discharges. Soundkeeper's toxicologists have reviewed the supporting data and approve of Ecology's foundation for determining the acute freshwater limit. We applaud Ecology's efforts here and in the State's role as a leader in this global issue.

## III. Chronic Freshwater and Acute and Chronic Marine Criteria

Acute freshwater toxicity is often the first, and arguably the least complex, of four important criteria.<sup>3</sup> The remaining three are chronic freshwater, acute marine, and chronic marine. Each of these play an important role for the quality of Washington's waters because it is a land of rivers, creeks, lakes, ponds, and streams, as well as estuaries, marine shoreline, and the inland waters of the Salish Sea. Effective environmental management requires understanding and mitigating each of these types of effects to preserve aquatic life and biodiversity.

### Acute and Chronic

"Acute conditions" are changes in the physical, chemical, or biologic environment which are expected or demonstrated to result in injury or death to an organism as a result of short-term exposure to the substance or detrimental environmental condition.<sup>4</sup> Typically, the exposure duration might range from a few minutes to about 96 hours. Acute toxicity can cause immediate harm to aquatic organisms, often leading to rapid symptoms or death. The effects are generally severe and occur soon after exposure. Acute toxicity is usually measured by lethal concentration values (LC50), which indicate the concentration of a substance that kills 50% of the test organisms within a specific short period.<sup>5</sup> For example, a chemical spill or runoff containing pesticides following a storm can lead to high concentrations of toxins in water bodies, causing acute effects to fish, invertebrates, and other aquatic life.

Alternatively, "chronic conditions" are changes in the physical, chemical, or biologic environment which are expected or demonstrated to result in injury or death to an organism as a result of repeated or constant exposure over an extended period of time to a substance or

---

<sup>3</sup> See Appendix A, ITRC What We Know: 6PPD and 6PPD-quinone (6PPD-q...ranks as one of the most potent acute aquatic toxicants when compared to chemicals with existing Clean Water Act Aquatic Life Ambient Water Quality Criteria. Most of the ecological toxicity data generated thus far focuses on the acute freshwater aquatic toxicity of 6PPD-q.)

<sup>4</sup> WAC 173-201A-020

<sup>5</sup> Sublethal effects refer to those impacts from toxins that do not result in immediate death. These effects can impair the health, behavior, or biological functions of organisms but are not necessarily fatal in the short term. Sublethal effects are more often associated with chronic toxicity.

detrimental environmental condition.<sup>6</sup> Chronic effects are the result of long-term exposure to a low or moderate concentration of a toxic substance.<sup>7</sup> This exposure can extend over weeks, months, or even the entire lifespan of the aquatic organism. Chronic exposure often results in long-term health effects that may not be immediately noticeable.<sup>8</sup> These effects can include reduced reproductive success, slower growth, impaired immune functions, and accumulated bioaccumulation of toxins.<sup>9</sup> Chronic toxicity is evaluated through metrics like the no observed effect concentration (NOEC) or the lowest observed effect concentration (LOEC), which are determined from long-term studies. For example, and relevant to the ubiquity of 6PPD-quinone in the environment, continuous discharge of industrial wastes, agricultural runoff, or polluted stormwater into water bodies can expose aquatic life to persistent low levels of harmful substances, leading to chronic toxicity effects.

While not defined in the WAC, subchronic effects are toxicologically relevant and are related to exposures that last for an intermediate duration sometime between acute and chronic exposure. Subchronic studies are designed to observe the effects of exposure over these intermediate periods to detect any potential health impacts that might not be apparent in short-term (acute) studies but do not require long-term (chronic) exposure to manifest.

### Freshwater and Marine

Acute and chronic effects of toxic substances can differ between freshwater and marine environments, influenced by several factors including the physical, chemical, and biological differences between these two types of aquatic systems.<sup>10</sup> Understanding these differences is crucial for effectively assessing and managing the ecological impacts of pollutants in different aquatic environments. Tailored approaches are necessary to protect the diverse and unique life forms found in freshwater and marine systems. Significantly, receiving waters of the Salish Sea are marine/estuarine while stormwater is usually fresh. Some of the marine organisms in the Salish Sea may be even more sensitive than freshwater receptors used to test in stormwater. In addition, threatened or endangered salmonids that are vulnerable to 6PPD-quinone live varying life stages in freshwater, brackish estuarine, and marine environments.

## IV. Conclusion

For the reasons above, Soundkeeper asks Ecology to use the power of momentum around 6PPD-quinone to develop chronic marine and freshwater criteria. Given the extreme mortality

---

<sup>6</sup> WAC 173-201A-020

<sup>7</sup> Adverse effects of chronic toxicity can manifest as direct lethality but more commonly refers to sublethal endpoints such as decreased growth, reduced reproduction, or behavioral changes such as impacted swimming performance.

<sup>8</sup> Chronic exposure might extend to much of an organism's lifetime.

<sup>9</sup> See e.g. Xin Hua, Xiao Feng, Geyu Liang, Jie Chao, Dayong Wang, Long-term exposure to tire-derived 6-PPD quinone causes intestinal toxicity by affecting functional state of intestinal barrier in *Caenorhabditis elegans*, *Science of The Total Environment*, Volume 861, 2023,160591,ISSN 0048-9697, available at <https://doi.org/10.1016/j.scitotenv.2022.160591> (environmentally relevant concentrations of 6-PPDQ were  $\leq 10$   $\mu\text{g/L}$ ...suggested that long-term exposure to 6-PPDQ at environmentally relevant concentrations potentially results in intestinal toxicity by disrupting functional state of intestinal barrier in organisms). See also Greer, J. B., Dalsky, E. M., Lane, R. F., & Hansen, J. D. (2023). Establishing an *in vitro* model to assess the toxicity of 6PPD-quinone and other tire wear transformation products. *Environmental Science & Technology Letters*, 10(6), 533–537. <https://doi.org/10.1021/acs.estlett.3c00196> (absence of mortality in Chinook salmon and sockeye salmon exposed to 6PPDQ...does not preclude the potential for sublethal effects that could have detrimental impacts on behavior, growth, or reproduction).

<sup>10</sup> Examples include: salinity; buffering capacity; temperature and pressure; species variation; species adaptation and tolerance; structure and diversity of biological communities; and types, sources, and concentration of pollution.

caused by this chemical wherever it is discharged – including into estuaries, rivers, marshes, and in places at the very mouths of salmon spawning streams, Soundkeeper also asks Ecology to apply the current freshwater acute criteria in marine waters as well. Soundkeeper further encourages Ecology to fully occupy its leadership role on this issue and take advantage of the process provided for in the Administrative Code to update the Toxics Criteria for Aquatic Life as needed protect aquatic life and to increase the technical accuracy of the criteria being applied with these additional freshwater and marine values.

Sincerely,

A handwritten signature in black ink that reads "Emily Gonzalez". The signature is written in a cursive, flowing style.

Emily Gonzalez  
Director of Law and Policy  
Puget Soundkeeper  
[emily@pugetsoundkeeper.org](mailto:emily@pugetsoundkeeper.org)

# **APPENDIX A**

## What We Know: 6PPD and 6PPD-quinone

In the short time since 6PPD-quinone (6PPD-q) was isolated and characterized, scientists have been working to understand its prevalence and behaviors in the environment. This focus sheet provides environmental officials with a brief overview of the current understanding of 6PPD-q sources, exposure, fate, transport, toxicity, and mitigation strategies. In-depth ITRC guidance will be released in summer 2024.

In 2020, researchers in Washington State discovered and identified 6PPD-quinone (6PPD-q) as the stormwater chemical responsible for urban runoff mortality syndrome observed in coho salmon (*Oncorhynchus kisutch*) around Puget Sound over the last 25 years.<sup>1,2</sup> Research has demonstrated that 6PPD-q is also acutely lethal to brook trout<sup>3</sup> and rainbow trout/steelhead.<sup>3-5</sup> 6PPD is the primary anti-degradant in tires and has been in use since the 1960s. 6PPD-q is one of

the products formed by the reaction of 6PPD and ozone (Figure 1). 6PPD-q may be present in many places impacted by tire use. 6PPD and 6PPD-q have been detected in stormwater and surface waters on many continents<sup>1,6-10</sup> and have been found in airborne particulates,<sup>11-14</sup> sediment,<sup>15</sup> soil,<sup>11</sup> rubber products other than tires,<sup>16</sup> and human urine.<sup>17</sup>





Figure 1. Chemical structures for 6PPD and 6PPD-quinone.

### How Is 6PPD-q Entering Surface Waters?

Tire wear particles (TWPs) containing 6PPD-q are transported via stormwater to surface water (Figure 2). Many urban stormwater systems are designed to control flooding, not capture and treat contaminants. In separate storm sewer systems, rainwater is transported to natural receiving waters through a network of ditches and pipes without natural or engineered green

spaces to remove pollutants prior to entering surface waters. Additionally, some areas with installed stormwater best management practices (BMPs) are failing to contain stormwater due to increased urbanization and storm events that are larger than the infrastructure was designed for, leading to direct conveyance of 6PPD-q to vulnerable aquatic ecosystems.

## What We Know: 6PPD and 6PPD-quinone

Multiple aspects of the lifecycle of 6PPD-q are under investigation. This includes the factors that influence the formation of 6PPD-q in tires and tire wear particles (TWPs) in the environment, 6PPD-q's leaching rates from TWPs, and its persistence and bioaccumulation potential. Programs that

divert scrap tires from landfills recycle the tires into crumb rubber materials used on sports fields, rubber-modified asphalt, tire-derived aggregate used in civil engineering projects, and more. The levels of 6PPD-q released from recycled tire products is also actively being researched.

### Conceptual Transport and Exposure Model




Figure 2. 6PPD in tires is converted to 6PPD-quinone (6PPD-q) when exposed to ozone. 6PPD-q is contained in tire wear particles that can be transported in the air and potentially inhaled by people. The particles can also be deposited on surfaces, soils, and plants, including foods, leading to potential plant uptake and human dermal exposure and ingestion. Tire wear particles can also stay near the roadway and be transported to surface waters through stormwater drains and runoff. 6PPD-q in surface waters can be ingested and absorbed by fishes. Exposed organisms can be ingested by humans and other species. 6PPD-q can potentially be mitigated by green stormwater infrastructure. Research is ongoing to further define 6PPD-q's environmental behaviors, exposures, and the potential development of adverse health outcomes. Figure credit: Hannah Vinyard, Washington State Department of Ecology.

### Ecological Toxicity

Both 6PPD and 6PPD-q surpass the threshold for *very high* acute aquatic toxicity using the Globally Harmonised System of Classification and Labeling of Chemicals.<sup>18</sup> This section focuses on 6PPD-q, which ranks as one of the most potent acute aquatic toxicants when compared to chemicals with existing Clean Water Act<sup>2</sup> Aquatic Life Ambient Water Quality Criteria. Most of the

ecological toxicity data generated thus far focuses on the acute freshwater aquatic toxicity of 6PPD-q. Two studies on the toxicity of 6PPD-q to marine organisms have been conducted,<sup>19,20</sup> but no studies have been done on the toxicity of 6PPD-q to the estuarine and marine stages of salmonids, which represents a significant data gap. Sublethal effects and chronic toxicity of



## What We Know: 6PPD and 6PPD-quinone

6PPD-q are being investigated. There is limited research regarding its toxicity to terrestrial species (e.g., *Caenorhabditis elegans*<sup>21-23</sup>).

**Acute Toxicity.** Research to date has demonstrated acute toxicity to 6PPD-q in only a few species within the salmonid family, which includes salmon, char, and trout (Table 1). Coho salmon are the most sensitive species documented, with a median LC<sub>50</sub> concentration (50% mortality in lab tests) of 0.08 µg/L and death occurring within hours.<sup>1,2,4,24</sup> Toxicity to 6PPD-q does not follow a phylogenetic relationship. Species within the *Oncorhynchus* genus show radically different acute toxicities, from an LC<sub>50</sub> as low as 0.040 µg/L in coho hatchlings<sup>24</sup> to no mortality observed in sockeye at 50 µg/L.<sup>25</sup> Some

salmonids in the *Salvelinus* genus (white-spotted char<sup>26</sup> and brook trout<sup>3</sup>) are acutely sensitive at relatively low concentrations (see Table 1), while others are not.<sup>3</sup> *Oncorhynchus mykiss*, which encompasses rainbow trout (freshwater only) and steelhead (ocean-going), show mortality at higher doses and a slower onset of symptoms in response to 6PPD-q.<sup>3</sup> The LC<sub>50</sub> for Chinook salmon is well above environmentally relevant concentrations<sup>25</sup>; however, Chinook had a low level of mortality when exposed to undiluted roadway runoff.<sup>27</sup> Salmonids that do not experience acute toxicity to 6PPD-q include sockeye salmon,<sup>25</sup> Arctic char,<sup>3</sup> Atlantic salmon, and brown trout,<sup>28</sup> as well as two varieties of Asiatic salmon: southern Dolly Varden and cherry salmon.<sup>26</sup>

**Table 1. Reported 6PPD-quinone LC<sub>50</sub> concentrations (50% observed mortality) of salmonids.**

Species	LC <sub>50</sub> (µg/L)	Test duration (h)	Toxicity Key
Coho salmon ( <i>Oncorhynchus kisutch</i> )	0.04, <sup>24</sup> 0.08, <sup>25</sup> 0.095 <sup>2</sup>	24	Higher
White-spotted char ( <i>Salvelinus leucomaenis pluvius</i> )	0.51 <sup>26</sup>	24	Higher
Brook trout ( <i>Salvelinus fontinalis</i> )	0.59 <sup>3</sup>	24	
Rainbow trout/steelhead ( <i>Oncorhynchus mykiss</i> )	0.64, <sup>29</sup> 1.0, <sup>3</sup> 2.26 <sup>5</sup>	96	
Chinook salmon ( <i>Oncorhynchus tshawytscha</i> )	67.3 <sup>24</sup> , 82.1 <sup>25</sup>	24	Lower
Sockeye salmon ( <i>Oncorhynchus nerka</i> )	Not acutely toxic at 50 <sup>25</sup>	24	
Atlantic salmon ( <i>Salmo salar</i> )	Not acutely toxic at 12.2 <sup>28</sup>	48	
Brown trout ( <i>Salmo trutta</i> )	Not acutely toxic at 12.2 <sup>28</sup>	48	
Arctic char ( <i>Salvelinus alpinus</i> )	Not acutely toxic at 12.7 <sup>3</sup>	24	
Southern Dolly Varden ( <i>Salvelinus curilus</i> )	Not acutely toxic at 3.8 <sup>26</sup>	48	
Cherry salmon ( <i>Oncorhynchus masou masou</i> )	Not acutely toxic at 3.5 <sup>26</sup>	48	

Note: Example species in the table are listed from very high to low across a toxicity gradient based on the LC<sub>50</sub> value, with the following ratings: coho = very high; white-spotted char and brook trout = high; rainbow trout / steelhead = medium high; Chinook salmon = medium low; and sockeye salmon, Atlantic salmon, brown trout, Arctic char, southern Dolly Varden, and cherry salmon = low. Chinook salmon were assigned medium-low toxicity out of an abundance of caution. They have an LC<sub>50</sub> above environmentally relevant concentrations and potentially above some of the salmonids listed below it in the table. Nevertheless, Chinook showed low levels of mortality in undiluted roadway runoff, which could be a result of 6PPD-q or another contaminant. Until further research clarifies whether any life stage of Chinook experiences acute mortality in response to 6PPD-q at potentially environmentally relevant exposures, they were assigned medium-low toxicity.

## What We Know: 6PPD and 6PPD-quinone

---

Chum salmon (*Oncorhynchus keta*) do not show toxicity to roadway runoff<sup>27</sup> or tire leachate<sup>30</sup> but have not been tested with 6PPD-q. Outside the salmonid family, environmentally relevant concentrations of 6PPD-q (up to 2.8 µg/L) are not fatal to several fishes (white sturgeon,<sup>3</sup> zebrafish, and medaka<sup>31</sup>) and aquatic invertebrates (*Daphnia* and the crustacean (*Hyalella azteca*)).<sup>31</sup> Toxicity studies on 6PPD-q in marine invertebrates, *Brachionus koreanus*<sup>19</sup> and *Parhyale hawaiiensis*,<sup>20</sup> indicated no acute toxicity.

Acute symptoms mimic respiratory distress and include gasping at the water's surface, fin splaying, and loss of equilibrium<sup>32</sup>; onset of symptoms can occur within 90 minutes.<sup>1</sup>

Examples of the effects are shown in the [Puget Soundkeeper — Longfellow Creek coho salmon video](#). Scientists are still working to determine how 6PPD-q causes mortality in fish. Researchers have demonstrated exposure to roadway runoff causes fluid to leak out of the blood vessels in the gills and brain of coho, demonstrating that the blood–brain barrier is compromised in coho.<sup>33</sup> Mahoney and colleagues provide evidence that energy production at the cellular level may be disrupted.<sup>34</sup> The researchers further suggest that the potential inability of sensitive species to metabolize 6PPD-q into a less toxic form may contribute to its selective toxicity.<sup>34</sup>

**Sublethal toxicity.** It is still unknown whether 6PPD-q causes sublethal toxicity in wild fish populations. Sublethal effects could impact growth and reproduction and make fishes susceptible to other stressors, such as pathogens, higher temperatures, or other poor water quality parameters. Additional studies are needed to determine the concentrations of 6PPD-q that could result in adverse effects to salmonids, particularly because some populations are protected under the Endangered Species Act.

Limited work has been done studying sublethal effects in zebrafish, where 6PPD-q influenced

embryo movement and heart rate.<sup>35</sup> In addition, environmentally relevant concentrations of 6PPD-q have been shown to alter the central nervous system of zebrafish, changing their exploratory behavior, wake/sleep cycle, and heart rate.<sup>36</sup> Beyond fish, chronic toxicity of 6PPD-q has been studied in *C. elegans*, a soil-dwelling round worm. The worms have neurobehavioral changes and show symptoms of oxidative stress at concentrations starting as low as 0.1 µg/L.<sup>21,22</sup> At 1 µg/L the worms have diminished reproductive capacity.<sup>23</sup> How these results translate to salmonids that are more susceptible to acute toxicity and how these sublethal effects relate to survival of aquatic species require further research.

### Human Health

This section provides the most salient (e.g., not comprehensive) toxicological information on 6PPD and 6PPD-q. The health effects of 6PPD are better characterized than 6PPD-q. The health hazards of other 6PPD transformation products remain another notable data gap.<sup>37</sup>

**6PPD.** 6PPD is a well-documented skin sensitizer, resulting in allergic contact dermatitis in sensitized individuals.<sup>38</sup> 6PPD is also listed as a category 1B reproductive toxicant by the European Chemicals Agency (ECHA).<sup>39</sup> Exposed rats experienced prolonged and difficult birth, including some pregnant rats to the point of death.<sup>39</sup> The no adverse effect level for reproduction is 7 mg/kg body weight per day for females.<sup>39</sup> 6PPD increased fat accumulation in liver in mice that were given oral doses of 10 mg/kg body weight per day for 6 weeks.<sup>40</sup> Similarly, ECHA identified the liver and blood cells as targets of toxicity in a 28-day oral exposure rat study. Effects on the liver were reversible at 20 mg/kg body weight per day, and both sexes showed fat deposition in the liver and anemia at 100 mg/kg body weight per day.<sup>39</sup>

## What We Know: 6PPD and 6PPD-quinone

---

**6PPD-quinone.** 6PPD-q is predicted to cause oxidative stress.<sup>41</sup> 6PPD-q increased lipid accumulation in the livers of mice that were given oral doses of 10 mg/kg body weight per day for 6 weeks.<sup>40</sup> In addition, 6PPD-q increased liver triglycerides at all doses tested (10, 30, and 100 mg/kg body weight per day).<sup>40</sup>

**Human Exposure.** 6PPD and 6PPD-q were detected in human urine in a Chinese study.<sup>17</sup> Pregnant women's urine had the highest levels of 6PPD and 6PPD-q out of all the demographic groups in the study.<sup>17</sup> One predicted route of exposure to these chemicals is inhalation of particulates, with the highest potential for exposure occurring near traffic (see Figure 2). Ingestion and incidental contact with rubber products or dust may be other sources of exposure to the chemicals.<sup>42</sup> 6PPD and 6PPD-q are present in tire crumb rubber, and these compounds have been identified in the bioaccessible fraction after extraction of crumb rubber with simulated gastrointestinal fluid, implying they may be absorbed after ingestion.<sup>43</sup> The risk of 6PPD and 6PPD-q to people who consume high levels of aquatic species has yet to be characterized. There are limited studies on the bioaccumulative properties of 6PPD and 6PPD-q. Fang and colleagues suggest that 6PPD and 6PPD-q may bioaccumulate in the livers of lab mice<sup>40</sup>; however, additional information regarding absorption, distribution, metabolism, and excretion by the exposed mice is needed to draw this conclusion. Contaminated sources for drinking water could potentially result in exposure depending on the source water and treatment method. Research is ongoing to address this question. Johannessen and colleagues reported negative findings in treated drinking water from two Canadian facilities.<sup>44</sup> No test results for U.S. drinking waters have been reported.

## Environmental Justice and Tribal Government Considerations

The extent of 6PPD and 6PPD-q impacts on vulnerable populations and overburdened communities will be determined as knowledge advances. Communities near roadways are disproportionately comprised lower-income people and people of color, making the potential impacts of airborne 6PPD and 6PPD-q on these communities a notable environmental justice concern.<sup>45,46</sup> Environmental justice considerations include but are not limited to food safety of fish consumption, drinking and recreational water safety, use of recycled rubber products, traffic proximity and air particulate matter exposure, socioeconomic impacts to subsistence and commercial fishers, and cumulative impacts.

Salmonid mortality, which can be caused by 6PPD-q, other toxic chemicals, climate change, habitat loss, and additional factors,<sup>18</sup> disproportionately impacts tribal nations in North America by threatening tribal treaty rights, access to traditional foods, and the cultural and economic well-being of Indigenous peoples. Fishing rights for many tribal nations are guaranteed by treaties that have been signed, ratified, and reaffirmed by the U.S. government. Concerns for tribal nations around 6PPD-q include impaired salmon recovery and hatchery efforts, sublethal impacts to fishes, reduced ecosystem resilience, and cumulative impacts to fishes and peoples.

## What We Know: 6PPD and 6PPD-quinone

### Fate and Transport

Stormwater is the primary transport mechanism for 6PPD-q to surface water. TWPs are generated as tires roll across the road, particularly during acceleration, braking, and turning. These particles, and the chemicals they contain, collect in road dust until stormwater transports them into the aquatic environment. In many cities in the United States, stormwater is diverted to wastewater treatment plants (WWTPs) through combined sewer and stormwater systems. Studies investigating 6PPD-q removal in WWTPs' have had mixed results. Several studies showed a strong reduction or removal of 6PPD-q to nondetect levels,<sup>10,47,48</sup> and another study showed an increase in mass in the effluent from the WWTP.<sup>49</sup> More research is needed to follow up on this. The presence of 6PPD and 6PPD-q in biosolids from WWTP remains a data gap.

The levels of 6PPD-q are highest during or following rain or snowmelt runoff<sup>8,47</sup> and have been measured in U.S. surface waters at

concentrations above the LC<sub>50</sub> values (see Table 1) for coho, brook trout, and potentially rainbow trout.<sup>1,2</sup> The levels of 6PPD-q in the water column can stay elevated for days<sup>6,8</sup>; the duration depends on the frequency of inputs, the site, and the characteristics of the receiving water. Fate and transport of the chemicals in estuaries and saltwater has not yet been characterized. 6PPD-q is expected to sorb to sediment or particles<sup>18,50</sup> and has been measured in sediment in China.<sup>15</sup> Additionally, TWPs may be airborne initially and could be transported long distances. The chemical and physical properties of 6PPD-q in the atmosphere are currently unknown. Notably, 6PPD-q has been measured in particulate matter, including in airborne particles less than 2.5 µm (PM<sub>2.5</sub>),<sup>14,51,52</sup> road dust,<sup>12,53</sup> and household dust.<sup>12</sup> The highest detections and concentration ranges measured in various environmental media are provided in Table 2).Table 2Table 2. 6PPD-quinone concentrations measured in roadway runoff, surface water, sediment, and particulate matter-2.5.

Media	Concentration Range	Notes	References
Roadway runoff	ND – 2.43 µg/L	Highest detection was in China by Cao et al.	2,6,9,10,54
Surface water	ND – 2.8 µg/L*	Highest detection was in the Don River in Toronto, Canada, roughly 35× higher than median coho LC <sub>50</sub> . Loading generally correlates with the amount of rainfall.	2,7,8,54–56
Sediment	ND – 18.2 ng/g	Highest in urban river sediment, present in deep sea sediment in China.	15
Particulate matter (up to PM <sub>2.5</sub> )	0.1 – 7,250 pg/m <sup>3</sup>	Highest detection alongside a road in Guangzhou, China.	14,51,52

\*Median LC<sub>50</sub> for coho (0.08 µg/L), brook trout (0.59 µg/L), and rainbow trout (1.0 µg/L) (see Table 1).

Notes: µg/L = microgram per liter, ng/g = nanogram per gram dry weight, ND = nondetect, pg/m<sup>3</sup> = picogram per cubic meter

## What We Know: 6PPD and 6PPD-quinone

---

### Monitoring and Analytical Methods

Monitoring for 6PPD and 6PPD-q in air, soil, surfaces, and water is challenging because environmentally relevant concentrations may be low and presence may be intermittent. Sampling studies indicate that 6PPD-q is detected at higher concentrations during or following rain and snowmelt events that occur following an extended dry period.<sup>8,56</sup> The rate of transport of stormwater and exposure to aquatic life increase with percent impervious surface within a watershed.<sup>8,56</sup> Studies have shown that 6PPD-q can persist for days in urban areas during or following storm events.<sup>8,57</sup> Efforts are underway by the state of Washington to evaluate passive sampling technologies and effectiveness.

The U.S. Environmental Protection Agency is developing a 6PPD-q test method for surface water and stormwater that is projected to be available late in 2023. *Standard Operating Procedure (SOP): Extraction and Analysis of 6PPD-q (Mel730136, Version 1.2)*<sup>58</sup> contains procedures for the extraction and the qualitative and quantitative analysis of 6PPD-q by triple quadrupole mass spectrometry. The standard operating procedure recommends the sample collection, preservation, storage, and holding times. In addition, several commercial and research laboratories can test for 6PPD-q in water.

**Other media.** Standardized methods are currently in development for sediment and biological tissues. 6PPD-q has been measured in the air by academic researchers,<sup>14,51,52</sup> but there is not a verified method for regulatory testing.

### Stormwater Best Management Practices

Stormwater research is focused on determining the effectiveness of existing and new BMPs at 6PPD-q removal, modifying stormwater systems to improve 6PPD-q removal, adding BMP retrofits to urban roadways that lack adequate space for green infrastructure, and refining green stormwater infrastructure to maximize 6PPD-q filtration.

**Effective Stormwater Mitigation Technologies.** A recent Washington State publication<sup>59</sup> evaluated stormwater BMP treatment mechanisms and rated their expected 6PPD and 6PPD-q removal effectiveness. Several source control, flow control, and runoff treatment stormwater control measures were found to be potentially effective solutions. Washington State is funding research to verify the efficacy of these BMPs and stormwater control measures.

Researchers have demonstrated that running stormwater through the bioretention soil mix (stormwater compost and sand) that is designed as a component of a bioretention system (Figure 3) prevents acute mortality in coho.<sup>60,61</sup> Research to optimize the depth and composition of the bioretention soil mix to maximize the effectiveness and longevity of the system is ongoing. Additionally, different media are being tested to reduce potential nutrient leaching from bioretention BMPs. Preliminary results of a study representing an accelerated timeline of 10 water years by passing water contaminated with 6PPD-q through a laboratory-managed bioretention soil mix shows prevention of coho mortality over the 18-month study period; the results of this study are being prepared by McIntyre and colleagues.

## What We Know: 6PPD and 6PPD-quinone




Figure 3. Typical bioretention system with design features. Current research is focused on optimizing the depth and composition of the bioretention soil mix. Courtesy of AHBL, Inc.

Researchers are also using compost-amended biofiltration swales comprising topsoil, compost, and vegetation (Figure 4) to determine the effectiveness of biofiltration systems alongside roadways; the results of this study are being prepared by Tian and colleagues. Preliminary results of the study show variability in compost-amended biofiltration swales performance across seasonal and storm specific parameters, with an efficiency of up to 80+% removal of 6PPD-q.

### Identifying Vulnerable Aquatic Areas.

Washington State is developing strategies to focus sampling and stormwater mitigation efforts in locations where 6PPD-q is having a critical environmental impact. The development of these strategies is based on collaboration with tribal governments, community engagement, and available GIS mapping tools containing parameters that are assumed to influence concentrations of 6PPD-q in surface waters.

## What We Know: 6PPD and 6PPD-quinone



Figure 4. Example of a compost-amended biofiltration swale. Stormwater is filtered as it flows along the grass in the swale and infiltrates into the topsoil and compost. Photo: Washington State Department of Transportation.

Factors used to identify these areas include, but are not limited to, level of traffic, impervious surfaces, precipitation, media composition that stormwater travels through to get to receiving water, size of the receiving water, and the presence of sensitive species.

Other ongoing research includes the effectiveness of permeable pavement to capture tire particles, analysis of different compost medias and biochar to determine the effectiveness of organic matter in bioretention systems, and evaluation of existing street-sweeping technologies and practices (timing and frequency) on removal of 6PPD and 6PPD-q from roadways.

### Alternatives to the Use of 6PPD in Tires

Identifying and deploying alternatives to 6PPD in tires can ultimately reduce or eliminate 6PPD-q in the environment. Currently, 6PPD is necessary for tire safety and to extend the life of tires by preventing cracking and degradation caused by ozone. Discussions with tire manufacturers have revealed that an anti-degradant is not currently available to replace 6PPD. Tires are complex products with tire safety as a principal design priority, and a fully functional anti-degradant is a necessity. Research is ongoing to identify safer alternative chemicals that provide the functionality of 6PPD in tires.<sup>62</sup> Due to the complexity of identifying, testing, and implementing a suitable alternative to 6PPD, the U.S. Tire Manufacturers Association cannot estimate the time frame for the replacement at this stage of the process. The states of California and Washington are pursuing policies to promote the advancement of alternatives to 6PPD in tires.

### State Policies and Regulations

Washington State is developing a statewide action plan, funding research to fill in data gaps, assessing other potential tire anti-degradants, and developing specific data requirements and standards to assess the hazards of the alternatives. [Technical Memo: Assessment of Potential Hazards of 6PPD and Alternatives](#)<sup>63</sup> provides an overview of known toxicological hazards of chemicals that are or have been used as anti-degradants in tires. Washington State is currently developing hazard criteria to define “safer” when looking at alternatives to 6PPD. There is currently no estimated timeline for completion of the action plan or alternatives assessment. The [Safer Products for Washington](#) program, which aims to reduce toxic chemicals in consumer products, identified 6PPD as a priority chemical. Washington and California supported

## What We Know: 6PPD and 6PPD-quinone


---

the [Collaborative Innovation Forum: Functional substitutes to 6PPD in tires to develop a road map for identifying safer alternatives to 6PPD](#).<sup>62</sup>

California's Department of Toxic Substances Control will begin regulating [6PPD in motor vehicles](#) through the [Safer Consumer Products Program](#) on October 1, 2023. These regulations require tire manufacturers to analyze the hazards and adverse environmental impacts of potential alternatives to 6PPD, as well as evaluate the benefits and tradeoffs of replacing 6PPD. This process leverages the technical expertise of the tire manufacturers and enables them to meet

vehicle safety and consumer product safety requirements, while providing a rigorous, transparent, and scientific framework to evaluate and compare potential alternatives to 6PPD. The tire manufacturers' initial screening of potential alternatives is due on March 29, 2024.

6PPD is on Minnesota's Toxic Free Kids Act Chemicals of High Concern List,<sup>64</sup> and the state's legislature appropriated nearly half a million dollars for research on 6PPD-q and its effect on state fishes.<sup>65</sup> Maine also includes 6PPD on its Chemicals of Concern list.<sup>66</sup>



### 6PPD Team Contacts

**Tanya Williams** • Washington State Department of Ecology  
[tanya.williams@ecy.wa.gov](mailto:tanya.williams@ecy.wa.gov)

**Kelly Grant** • Department of Toxic Substances Control  
California Environmental Protection Agency  
[kelly.grant@dtsc.ca.gov](mailto:kelly.grant@dtsc.ca.gov)

**Evan Madden** • Interstate Technology and Regulatory Council  
[emadden@ecos.org](mailto:emadden@ecos.org)

September 2023

---

This fact sheet incorporates data through July 2023.

The **Interstate Technology and Regulatory Council (ITRC)** is a state-led environmental coalition working to create innovative solutions and best management practices. ITRC produces documents and training that broaden and deepen technical knowledge and expedite quality regulatory decision-making while protecting human health and the environment. In January 2023, ITRC started the Tire Anti-degradants (6PPD) Team to provide guidance documents and tools on 6PPD and 6PPD-quinone for environmental officials.



ITRC  
1250 H St. NW, Suite 850  
Washington, DC 20005  
[itrcweb.org](http://itrcweb.org)





# What We Know: 6PPD and 6PPD-quinone

---

## References

1. Tian Z, Zhao H, Peter KT, Gonzalez M, Wetzel J, Wu C, Hu X, Prat J, Mudrock E, Hettinger R, et al. A ubiquitous tire rubber-derived chemical induces acute mortality in coho salmon. *Science*. 2021;371(6525):185–189. doi:10.1126/science.abd6951
2. Tian Z, Gonzalez M, Rideout CA, Zhao HN, Hu X, Wetzel J, Mudrock E, James CA, McIntyre JK, Kolodziej EP. 6PPD-quinone: Revised toxicity assessment and quantification with a commercial standard. *Environmental Science & Technology Letters*. 2022 Jan 11:acs.estlett.1c00910. doi:10.1021/acs.estlett.1c00910
3. Brinkmann M, Montgomery D, Selinger S, Miller JGP, Stock E, Alcaraz AJ, Challis JK, Weber L, Janz D, Hecker M, et al. Acute toxicity of the tire rubber-derived chemical 6PPD-quinone to four fishes of commercial, cultural, and ecological importance. *Environmental Science & Technology Letters*. 2022 Mar 2:acs.estlett.2c00050. doi:10.1021/acs.estlett.2c00050
4. Greer JB, Dalsky EM, Lane RF, Hansen JD. Establishing an In Vitro Model to Assess the Toxicity of 6PPD-Quinone and Other Tire Wear Transformation Products. *Environmental Science & Technology Letters*. 2023 May 2 [accessed 2023 May 8]. <https://doi.org/10.1021/acs.estlett.3c00196>. doi:10.1021/acs.estlett.3c00196
5. Di S, Liu Z, Zhao H, Li Y, Qi P, Wang Z, Xu H, Jin Y, Wang X. Chiral perspective evaluations: Enantioselective hydrolysis of 6PPD and 6PPD-quinone in water and enantioselective toxicity to *Gobio cypris rarus* and *Oncorhynchus mykiss*. *Environment International*. 2022;166:107374. doi:10.1016/j.envint.2022.107374
6. Challis JK, Popick H, Prajapati S, Harder P, Giesy JP, McPhedran K, Brinkmann M. Occurrences of Tire Rubber-Derived Contaminants in Cold-Climate Urban Runoff. *Environmental Science & Technology Letters*. 2021 Sep 22:acs.estlett.1c00682. doi:10.1021/acs.estlett.1c00682
7. Rauert C, Charlton N, Okoffo ED, Stanton RS, Agua AR, Pirrung MC, Thomas KV. Concentrations of tire additive chemicals and tire road wear particles in an Australian urban tributary. *Environmental Science & Technology*. 2022 Jan 31:acs.est.1c07451. doi:10.1021/acs.est.1c07451
8. Johannessen C, Helm P, Lashuk B, Yargeau V, Metcalfe CD. The tire wear compounds 6PPD-quinone and 1,3-diphenylguanidine in an urban watershed. *Archives of Environmental Contamination and Toxicology*. 2021 Aug 4 [accessed 2021 Aug 5]. <https://doi.org/10.1007/s00244-021-00878-4>. doi:10.1007/s00244-021-00878-4
9. Cao G, Wang W, Zhang J, Wu P, Zhao X, Yang Z, Hu D, Cai Z. New Evidence of Rubber-Derived Quinones in Water, Air, and Soil. *Environmental Science & Technology*. 2022;56(7):4142–4150. doi:10.1021/acs.est.1c07376
10. Maurer L, Carmona E, Machate O, Schulze T, Krauss M, Brack W. Contamination Pattern and Risk Assessment of Polar Compounds in Snow Melt: An Integrative Proxy of Road Runoffs. *Environmental Science & Technology*. 2023;57(10):4143–4152. doi:10.1021/acs.est.2c05784
11. Cao G, Wang W, Zhang J, Wu P, Zhao X, Yang Z, Hu D, Cai Z. New evidence of rubber-derived quinones in water, air, and soil. *Environmental Science & Technology*. 2022;56(7):4142–4150. doi:10.1021/acs.est.1c07376
12. Huang W, Shi Y, Huang J, Deng C, Tang S, Liu X, Chen D. Occurrence of substituted *p*-phenylenediamine antioxidants in dusts. *Environmental Science & Technology Letters*. 2021;8(5):381–385. doi:10.1021/acs.estlett.1c00148
13. Zhang Y-J, Xu T-T, Ye D-M, Lin Z-Z, Wang F, Guo Y. Widespread *N*-(1,3-Dimethylbutyl)-*N'*-phenyl-*p*-phenylenediamine quinone in size-fractionated atmospheric particles and dust of different indoor environments. *Environmental Science & Technology Letters*. 2022 Apr 25 [accessed 2022 May 2]. <https://doi.org/10.1021/acs.estlett.2c00193>. doi:10.1021/acs.estlett.2c00193
14. Johannessen C, Saini A, Zhang X, Harner T. Air monitoring of tire-derived chemicals in global megacities using passive samplers. *Environmental Pollution*. 2022;314:120206. doi:10.1016/j.envpol.2022.120206

## What We Know: 6PPD and 6PPD-quinone

---

15. Zeng L, Li Y, Sun Y, Liu L-Y, Shen M, Du B. Widespread occurrence and transport of *p*-Phenylenediamines and their quinones in sediments across urban rivers, estuaries, coasts, and deep-sea regions. *Environmental Science & Technology*. 2023 Jan 31;acs.est.2c07652. doi:10.1021/acs.est.2c07652
16. Zhao HN, Hu X, Gonzalez M, Rideout CA, Hobby GC, Fisher MF, McCormick CJ, Dodd MC, Kim KE, Tian Z, et al. Screening *p*-Phenylenediamine antioxidants, their transformation products, and industrial chemical additives in crumb rubber and elastomeric consumer products. *Environmental Science & Technology*. 2023 Feb 9;acs.est.2c07014. doi:10.1021/acs.est.2c07014
17. Du B, Liang B, Li Y, Shen M, Liu L-Y, Zeng L. First report on the occurrence of *N*-(1,3-Dimethylbutyl)-*N*'-phenyl-*p*-phenylenediamine (6PPD) and 6PPD-quinone as pervasive pollutants in human urine from South China. *Environmental Science & Technology Letters*. 2022;9(12):1056–1062. doi:10.1021/acs.estlett.2c00821
18. DTSC. Product-Chemical Profile for Motor Vehicle Tires Containing 6PPD - Department of Toxic Substances Control (DTSC). 2022. [https://dtsc.ca.gov/wp-content/uploads/sites/31/2022/05/6PPD-in-Tires-Priority-Product-Profile\\_FINAL-VERSION\\_accessible.pdf](https://dtsc.ca.gov/wp-content/uploads/sites/31/2022/05/6PPD-in-Tires-Priority-Product-Profile_FINAL-VERSION_accessible.pdf)
19. Maji UJ, Kim K, Yeo I-C, Shim K-Y, Jeong C-B. Toxicological effects of tire rubber-derived 6PPD-quinone, a species-specific toxicant, and dithiobisbenzanilide (DTBBA) in the marine rotifer *Brachionus koreanus*. *Marine Pollution Bulletin*. 2023;192:115002. doi:10.1016/j.marpolbul.2023.115002
20. Botelho MT, Militão GG, Brinkmann M, Umbuzeiro G de A. Toxicity and mutagenicity studies of 6PPD-quinone in a marine invertebrate species and bacteria. *Environmental and Molecular Mutagenesis*. [accessed 2023 Jul 7];n/a(n/a). <https://onlinelibrary.wiley.com/doi/abs/10.1002/em.22560>. doi:10.1002/em.22560
21. Hua X, Feng X, Liang G, Chao J, Wang D. Long-term exposure to tire-derived 6-PPD quinone causes intestinal toxicity by affecting functional state of intestinal barrier in *Caenorhabditis elegans*. *Science of The Total Environment*. 2022 Dec:160591. doi:10.1016/j.scitotenv.2022.160591
22. Hua X, Feng X, Liang G, Chao J, Wang D. Exposure to 6-PPD quinone at environmentally relevant concentrations causes abnormal locomotion behaviors and neurodegeneration in *Caenorhabditis elegans*. *Environmental Science & Technology*. 2023 Mar 13;acs.est.2c08644. doi:10.1021/acs.est.2c08644
23. Hua X, Feng X, Liang G, Chao J, Wang D. Long-term exposure to 6-PPD quinone reduces reproductive capacity by enhancing germline apoptosis associated with activation of both DNA damage and cell corpse engulfment in *Caenorhabditis elegans*. *Journal of Hazardous Materials*. 2023;454:131495. doi:10.1016/j.jhazmat.2023.131495
24. Lo B, Marlatt V, Liao X, Reger S, Gallilee C, Brown T. Acute toxicity of 6PPD-quinone to early life stage juvenile Chinook (*Oncorhynchus tshawytscha*) and coho (*Oncorhynchus kisutch*) salmon. *Environmental Toxicology and Chemistry*. 2023;42:815–822. doi:doi: 10.1002/etc.5568
25. Greer JB, Dalsky EM, Lane RF, Hansen JD. Establishing an in vitro model to assess the toxicity of 6ppd-quinone and other tire wear transformation products. *Environmental Science & Technology Letters*. 2023 May 2 [accessed 2023 May 8]. <https://doi.org/10.1021/acs.estlett.3c00196>. doi:10.1021/acs.estlett.3c00196
26. Hiki K, Yamamoto H. The tire-derived chemical 6PPD-quinone is lethally toxic to the white-spotted char *Salvelinus leucomaenis pluvius* but not to two other salmonid species. *Environmental Science & Technology Letters*. 2022 Nov 7;acs.estlett.2c00683. doi:10.1021/acs.estlett.2c00683
27. French BF, Baldwin DH, Cameron J, Prat J, King K, Davis JW, McIntyre JK, Scholz NL. Urban roadway runoff is lethal to juvenile coho, steelhead, and Chinook salmonids, but not congeneric sockeye. *Environmental Science & Technology Letters*. 2022 Aug 24 [accessed 2022 Aug 29]. <https://doi.org/10.1021/acs.estlett.2c00467>. doi:10.1021/acs.estlett.2c00467

## What We Know: 6PPD and 6PPD-quinone

---

28. Foldvik A, Kryuchkov F, Sandodden R, Uhlig S. Acute Toxicity Testing of the Tire Rubber-Derived Chemical 6PPD-quinone on Atlantic Salmon (*Salmo salar*) and Brown Trout (*Salmo trutta*). *Environmental Toxicology and Chemistry*. 2022;41(12):3041–3045. doi:10.1002/etc.5487
29. Nair P, Sun J, Xie L, Kennedy L, Kozakiewicz D, Kleywegt S, Hao C, Byun H, Barrett H, Baker J, et al. Synthesis and Toxicity Evaluation of Tire Rubber-Derived Quinones (working paper). 2023 [accessed 2023 Jun 26]. <https://chemrxiv.org/engage/chemrxiv/article-details/648ccfec4f8b1884b7669239>. doi:10.26434/chemrxiv-2023-pmxcv
30. McIntyre JK, Prat J, Cameron J, Wetzel J, Mudrock E, Peter KT, Tian Z, Mackenzie C, Lundin J, Stark JD, et al. Treading water: Tire wear particle leachate recreates an urban runoff mortality syndrome in coho but not chum salmon. *Environmental Science*. 2021;8.
31. Hiki K, Asahina K, Kato K, Yamagishi T, Omagari R, Iwasaki Y, Watanabe H, Yamamoto H. Acute toxicity of a tire rubber-derived chemical, 6PPD quinone, to freshwater fish and crustacean species. *Environmental Science & Technology Letters*. 2021;8(9):779–784. doi:10.1021/acs.estlett.1c00453
32. Scholz NL, Myers MS, McCarthy SG, Labenia JS, McIntyre JK, Ylitalo GM, Rhodes LD, Laetz CA, Stehr CM, French BL, et al. Recurrent die-offs of adult coho salmon returning to spawn in Puget Sound lowland urban streams. *PLOS ONE*. 2011;6(12):e28013. doi:10.1371/journal.pone.0028013
33. Blair SI, Barlow CH, McIntyre JK. Acute cerebrovascular effects in juvenile coho salmon exposed to roadway runoff. *Canadian Journal of Fisheries and Aquatic Sciences*. 2021 Feb [accessed 2021 Jan 28]. <https://cdnscepub.com/doi/abs/10.1139/cjfas-2020-0240>. doi:10.1139/cjfas-2020-0240
34. Mahoney H, da Silva Junior FC, Roberts C, Schultz M, Ji X, Alcaraz AJ, Montgomery D, Selinger S, Challis JK, Giesy JP, et al. Exposure to the tire rubber-derived contaminant 6PPD-quinone causes mitochondrial dysfunction *in vitro*. *Environmental Science & Technology Letters*. 2022 Aug 4:acs.estlett.2c00431. doi:10.1021/acs.estlett.2c00431
35. Zhang S-Y, Gan X, Shen B, Jiang J, Shen H, Lei Y, Liang Q, Bai C, Huang C, Wu W, et al. 6PPD and its metabolite 6PPD-q induce different developmental toxicities and phenotypes in embryonic zebrafish. *Journal of Hazardous Materials*. 2023;455:131601. doi:10.1016/j.jhazmat.2023.131601
36. Ricarte M, Prats E, Montemurro N, Bedrossiantz J, Bellot M, Gómez-Canela C, Raldúa D. Environmental concentrations of tire rubber-derived 6PPD-quinone alter CNS function in zebrafish larvae. *Science of The Total Environment*. 2023;896:165240. doi:10.1016/j.scitotenv.2023.165240
37. Hu X, Zhao HN, Tian Z, Peter KT, Dodd MC, Kolodziej EP. Transformation product formation upon heterogeneous ozonation of the tire rubber antioxidant 6PPD (n-(1,3-dimethylbutyl)-n'-phenyl-p-phenylenediamine). *Environmental Science & Technology Letters*. 2022 Apr 12 [accessed 2022 Apr 12]. <https://doi.org/10.1021/acs.estlett.2c00187>. doi:10.1021/acs.estlett.2c00187
38. ToxServices, LLC. N-(1,3-Dimethylbutyl)-N'-Phenyl-P-Phenylenediamine (6PPD) (CAS# 793-24-8) GreenScreen® for Safer Chemicals (GreenScreen®) Assessment. 2021. [https://www.ezview.wa.gov/Portals/\\_1962/Documents/6ppd/GreenScreenExecutiveSummaryFor6PPD.pdf](https://www.ezview.wa.gov/Portals/_1962/Documents/6ppd/GreenScreenExecutiveSummaryFor6PPD.pdf)
39. ECHA. 6PPD: 1,4-Benzenediamine, N1-(1,3-dimethylbutyl)-N4-phenyl- Registration Dossier - European Chemicals Agency (ECHA). 2021 Mar 29 [accessed 2021 Mar 29]. <https://echa.europa.eu/registration-dossier/-/registered-dossier/15367/5/1>
40. Fang L, Fang C, Di S, Yu Y, Wang C, Wang X, Jin Y. Oral exposure to tire rubber-derived contaminant 6PPD and 6PPD-quinone induce hepatotoxicity in mice. *Science of The Total Environment*. 2023;869:161836. doi:10.1016/j.scitotenv.2023.161836
41. Wang W, Cao G, Zhang J, Chen Z, Dong C, Chen J, Cai Z. p-Phenylenediamine-Derived Quinones as New Contributors to the Oxidative Potential of Fine Particulate Matter. *Environmental Science & Technology Letters*. 2022 Aug 10:acs.estlett.2c00484. doi:10.1021/acs.estlett.2c00484

## What We Know: 6PPD and 6PPD-quinone

---

42. Castan S, Sherman A, Peng R, Zumstein MT, Wanek W, Hüffer T, Hofmann T. Uptake, Metabolism, and Accumulation of Tire Wear Particle-Derived Compounds in Lettuce. *Environmental Science & Technology*. 2022 Dec 28 [accessed 2023 Jan 3]. <https://doi.org/10.1021/acs.est.2c05660>. doi:10.1021/acs.est.2c05660
43. Armada D, Martinez-Fernandez A, Celeiro M, Dagnac T, Llompart M. Assessment of the bioaccessibility of PAHs and other hazardous compounds present in recycled tire rubber employed in synthetic football fields. *Science of The Total Environment*. 2023;857:159485. doi:10.1016/j.scitotenv.2022.159485
44. Johannessen C, Metcalfe CD. The occurrence of tire wear compounds and their transformation products in municipal wastewater and drinking water treatment plants. *Environmental Monitoring and Assessment*. 2022;194(10):731. doi:10.1007/s10661-022-10450-9
45. Rowangould GM. A census of the US near-roadway population: Public health and environmental justice considerations. *Transportation Research, Part D: Transport and Environment*. 2013;2013(25):59–67. doi:10.1016/j.trd.2013.08.003
46. Tian N, Xue J, Barzyk TM. Evaluating socioeconomic and racial differences in traffic-related metrics in the United States using a GIS approach. *The Journal of Exposure Science and Environmental Epidemiology*. 2013;(23):215–222.
47. Seiwert B, Nihemaiti M, Troussier M, Weyrauch S, Reemtsma T. Abiotic oxidative transformation of 6-PPD and 6-PPD quinone from tires and occurrence of their products in snow from urban roads and in municipal wastewater. *Water Research*. 2022;212:118122. doi:10.1016/j.watres.2022.118122
48. Zhang H-Y, Huang Z, Liu Y-H, Hu L-X, He L-Y, Liu Y-S, Zhao J-L, Ying G-G. Occurrence and risks of 23 tire additives and their transformation products in an urban water system. *Environment International*. 2023;171:107715. doi:10.1016/j.envint.2022.107715
49. Johannessen C, Metcalfe CD. The occurrence of tire wear compounds and their transformation products in municipal wastewater and drinking water treatment plants. *Environmental Monitoring and Assessment*. 2022;194(10):731. doi:10.1007/s10661-022-10450-9
50. Hu X, Zhao H (Nina), Tian Z, Peter KT, Dodd MC, Kolodziej EP. Chemical characteristics, leaching, and stability of the ubiquitous tire rubber-derived toxicant 6PPD-quinone. *Environmental Science: Processes & Impacts*. 2023;25(5):901–911. doi:10.1039/D3EM00047H
51. Wang W, Cao G, Zhang J, Wu P, Chen Y, Chen Z, Qi Z, Li R, Dong C, Cai Z. Beyond substituted *p*-Phenylenediamine antioxidants: prevalence of their quinone derivatives in PM<sub>2.5</sub>. *Environmental Science & Technology*. 2022 Jul 14:acs.est.2c02463. doi:10.1021/acs.est.2c02463
52. Zhang Y, Xu C, Zhang W, Qi Z, Song Y, Zhu L, Dong C, Chen J, Cai Z. *p*-Phenylenediamine antioxidants in PM<sub>2.5</sub>: The underestimated urban air pollutants. *Environmental Science & Technology*. 2021 Sep 22:acs.est.1c04500. doi:10.1021/acs.est.1c04500
53. Deng C, Huang J, Qi Y, Chen D, Huang W. Distribution patterns of rubber tire-related chemicals with particle size in road and indoor parking lot dust. *Science of The Total Environment*. 2022;844:157144. doi:10.1016/j.scitotenv.2022.157144
54. Zhao HN, Hu X, Tian Z, Gonzalez M, Rideout CA, Peter KT, Dodd MC, Kolodziej EP. Transformation Products of Tire Rubber Antioxidant 6PPD in Heterogeneous Gas-Phase Ozonation: Identification and Environmental Occurrence. *Environmental Science & Technology*. 2023;57(14):5621–5632. doi:10.1021/acs.est.2c08690
55. Johannessen C, Helm P, Metcalfe CD. Detection of selected tire wear compounds in urban receiving waters. *Environmental Pollution*. 2021;287:117659. doi:10.1016/j.envpol.2021.117659
56. Nedrich S. Preliminary Investigation of the Occurrence of 6PPD-Quinone in Michigan's Surface Water. 2022. doi:10.13140/RG.2.2.34478.59204

## What We Know: 6PPD and 6PPD-quinone

---

57. Challis JK, Popick H, Prajapati S, Harder P, Giesy JP, McPhedran K, Brinkmann M. Occurrences of Tire Rubber-Derived Contaminants in Cold-Climate Urban Runoff. *Environmental Science & Technology Letters*. 2021;8(11):961–967. doi:10.1021/acs.estlett.1c00682
58. Washington State Department of Ecology. Standard Operating Procedure (SOP): Extraction and Analysis of 6PPD-Quinone (Mel730136, Version 1.2). 2023.
59. Navickis-Brasch A, Maurer M, Hoffman-Ballard T, Bator S, Diamond J. Stormwater Treatment of Tire Contaminants Best Management Practices Effectiveness. 2022. p. 72. [https://fortress.wa.gov/ecy/ezshare/wq/Permits/Flare/2019SWMMWW/Content/Resources/DocsForDownload/2022\\_SWTreatmentOfTireContaminants-BMPEffectiveness.pdf](https://fortress.wa.gov/ecy/ezshare/wq/Permits/Flare/2019SWMMWW/Content/Resources/DocsForDownload/2022_SWTreatmentOfTireContaminants-BMPEffectiveness.pdf)
60. Spromberg JA, Baldwin DH, Damm SE, McIntyre JK, Huff M, Sloan CA, Anulacion BF, Davis JW, Scholz NL. Coho salmon spawner mortality in western US urban watersheds: bioinfiltration prevents lethal storm water impacts. *Journal of Applied Ecology*. 2016;53(2):398–407. doi:<https://doi.org/10.1111/1365-2664.12534>
61. McIntyre JK, Davis JW, Hinman C, Macneale KH, Anulacion BF, Scholz NL, Stark JD. Soil bioretention protects juvenile salmon and their prey from the toxic impacts of urban stormwater runoff. *Chemosphere*. 2015;132:213–219. doi:10.1016/j.chemosphere.2014.12.052
62. Sustainable Chemistry Catalyst. Collaborative Innovation Forum: Functional Substitutes to 6PPD in Tires. Meeting Report. 2023. <https://static1.squarespace.com/static/633b3dd6649ed62926ed7271/t/63ee6cd15eb30a0fd4f0630d/1676569810601/6PPD-in-Tires-Innovation-Forum-Meeting-Report.pdf>
63. Washington State Department of Ecology. Technical Memo: Assessment of Potential Hazards of 6PPD and Alternatives. 2021. [https://www.ezview.wa.gov/Portals/\\_1962/Documents/6ppd/6PPD%20Alternatives%20Technical%20Memo.pdf](https://www.ezview.wa.gov/Portals/_1962/Documents/6ppd/6PPD%20Alternatives%20Technical%20Memo.pdf)
64. Minnesota Department of Health. Minnesota Toxic Free Kids Act 2019 Chemicals of High Concern. Toxic Free Kids Act: Chemicals of High Concern - EH: Minnesota Department of Health. 2019 [accessed 2021 Mar 30]. <https://www.health.state.mn.us/communities/environment/childenvhealth/tfka/highconcern.html>
65. Hall B. House environment panel OKs bill to appropriate \$47 million from state’s Clean Water Fund - Session Daily - Minnesota House of Representatives. Minnesota Legislature. 2022 Mar 24 [accessed 2023 Jun 17]. <https://www.house.mn.gov/sessiondaily/Story/17300>
66. Maine DEP. Chemicals of Concern. Maine Department of Environmental Protection (DEP). Chemicals of Concern, Safer Chemicals, Maine DEP. 2017 Jul [accessed 2021 Apr 5]. <https://www.maine.gov/dep/safechem/childrens-products/concern/index.html>