

Breck Lebegue MD MPH

Thank you for keeping WA air clean! A vast body of research proves that fossil fuel particulates cause human disease, shorten lives and eventually kill vulnerable kids, seniors and poor people who live near freeways, freight yards and high-traffic areas that produce combustion pollution. Hold strong despite Federal rollbacks of clean air standards, to continue to produce and publicize scientific truth. Thanks Breck Lebegue MD MPH, Co-Chair WA Physicians for Social Responsibility, Climate & Health Task Force www.wpsr.org/transportation



Fueling Sickness:

The Hidden Health Costs of Fossil Fuel Pollution



Fueling Sickness: The Hidden Health Costs of Fossil Fuel Pollution

Endorsed By:



Acknowledgements:

Thank you to the following individuals for their help in drafting and reviewing various sections of this document: Scott Barstow, MSc, Laura Kate Bender, Robert Brook, MD, Robert Byron, MD, MPH, Paige Knappenberger, Shyamala Rajan, Linda Rudolph, MD, MPH, Mona Sarfaty, MD, MPH, FAAFP, Joan Schiller, MD, Liz Scott, Caren Solomon, MD, MPH, Emily Thompson, George Thurston, ScD, Ernesto Villasensor, Diana van Vleet, Skye Wheeler





Table of Contents

Endorsed By/Acknowledgements.....	1
Introduction.....	3
Key Health Impacts.....	4
Health Harms of the Fossil Fuel Life Cycle.....	4
How Do Fossil Fuels Impact Our Health?.....	6
Respiratory Health	6
Cardiovascular Health	7
Neurological/Nervous System Health	8
Mental Health	9
Reproductive Health.....	9
Kidney Health	10
Endocrine System and Metabolic Health	11
Cancer	11
Plastics.....	12
Individuals that Face Higher Risk	13
Climate Change.....	15
Conclusion and Recommendations.....	17
References	18





Introduction

Fossil fuels – including coal, oil and gas – have shaped the modern world, powering advances in industry, mobility and daily life, including in medicine and public health. Yet the human health costs of fossil fuel use are enormous. From extraction to end use, pollution from fossil fuels contaminates the air we breathe, the water we drink and the food we eat, driving illness and premature death across every stage of life and every system of the body.

Air pollution from fossil fuels kills hundreds of thousands of people in the U.S. and millions worldwide every year. Fossil fuel combustion also causes climate change, another profound threat to human health. They also pollute our water and food through their use in making plastics and other petrochemicals.

This brief is not a comprehensive technical report. Instead, it offers a summary of how fossil fuels directly damage health: across the human life span and body systems and across the fossil fuel life cycle, from extraction to end use. It also highlights populations at heightened risk and points to key steps for protecting health. The extensive health impacts of climate change, driven by fossil fuel usage, are addressed after the direct health impacts.

Health and medical professionals see these harms in patients and communities every day. Our aim is to elevate the often-overlooked health costs of fossil fuels in conversations about our energy future. Too often, debates about the energy transition focus on economics, security or consumer choice. But we cannot ignore the devastating health burden of fossil fuels, especially when cleaner, more affordable alternatives are available. As health and medical professionals and organizations, we believe a transition to non-polluting energy is not only possible, but essential – for a healthier future.





Key Health Impacts

Fossil fuels are non-renewable resources. They formed millions of years ago when prehistoric plants and animals died and were gradually buried by layers of rock, then transformed by changing temperatures and pressure into coal, oil and gas. These fossil fuels are now extracted, processed, transported, burned and disposed of in ways that pollute air, water and land. At every stage of their life cycle, they harm human health and the environment.

Everyone in the U.S. is exposed to many toxic pollutants from fossil fuels. These exposures accumulate in our bodies over time and cause long-term health harms.¹ They affect every stage of human life, from before birth through old age. There is no safe threshold for fossil fuel pollution exposure – even very low levels of exposure may be harmful.²

While national and state air quality standards have led to significant improvements in air quality over several decades, air pollution remains a leading environmental health risk in the U.S. Globally, studies estimate that fossil fuel pollution is responsible for between 5 and 8 million deaths each year – including an estimated 350,000 premature deaths in the U.S. alone.^{3,4}

Beyond deaths, fossil fuel pollution is making people sick, including through heart and lung disease, cancer, adverse reproductive outcomes, neurological disorders and other chronic conditions. The healthcare, economic and societal costs of illness and deaths attributable to fossil fuel pollution are immense. These illnesses and deaths are preventable, which is why strong, evidence-based policy action is essential. This report includes recommendations that policymakers can take to protect health and accelerate the transition to a cleaner, healthier future.

Health Harms Across the Fossil Fuel Life Cycle

Each stage of the fossil fuel life cycle – extraction, processing, transportation, combustion and waste – creates significant health harms. Emissions from oil and gas use in the U.S. alone are estimated to cause more than 216,000 cases of childhood asthma and 10,000 preterm births each year. Communities of color and low-income populations bear a greater share of the health burden at nearly every stage of the fossil fuel supply chain.⁵ The cumulative health harms of fossil fuels extend far beyond climate change, imposing an enormous and preventable toll on public health.

Extraction:

Coal mining, oil drilling and hydraulic fracturing (fracking) for gas release harmful air pollutants that endanger workers and nearby communities. Coal extraction also scars the land, pollutes waterways and leaves behind toxic waste that threatens both ecosystems and human health. Fracking for methane gas (called “natural gas”) is linked to water contamination and health problems including childhood cancer, birth defects, kidney and respiratory disease. Gas leaks and flaring (the burning of excess gas) release hazardous pollutants such as volatile organic compounds (VOCs) and nitrogen oxides (NOx), which increase cancer risk and trigger asthma, respectively.



Fueling Sickness: The Hidden Health Costs of Fossil Fuel Pollution

Processing, Refining and Petrochemical Manufacturing:

After extraction, fossil fuels are processed and refined in industrial facilities called petrochemical plants, often located near residential neighborhoods. Workers and nearby residents are exposed to a complex mixture of hazardous air pollutants, including benzene, formaldehyde and other carcinogens, which heighten risks of cancer, respiratory illness and adverse birth outcomes. These facilities also emit particulate matter, nitrogen oxides and sulfur dioxide, worsening asthma and other chronic lung diseases in surrounding communities.

Transportation and Distribution:

Pipelines, trains, ships and trucks each pose hazards throughout the transport of fossil fuels. Pipeline leaks and ruptures happen regularly and contaminate soil and water and can cause explosions. Transport by rail and truck increases risks of accidents, spills and community exposure to harmful pollutants. Port communities near shipping hubs also face high concentrations of diesel exhaust, which is classified as a known carcinogen and linked to asthma, heart disease and premature death.

Combustion:

The greatest health damage arises from burning coal, oil and gas for electricity, transportation, heating and industrial use. Alongside greenhouse gases, combustion generates harmful air pollutants, including fine particulate matter (PM2.5), NOx, sulfur dioxide, ozone and toxic metals – pollutants strongly linked to cardiovascular disease, respiratory illness, adverse birth outcomes and premature death. Transportation is a major source of this pollution, as cars, trucks, trains, ships and other equipment powered by fossil fuels emit large volumes of exhaust that directly harm health, particularly in urban areas and those nearest traffic-related pollution sources.^{6,7}

Waste Products:

Even after combustion, fossil fuel waste continues to harm health. Coal-fired power plants generate coal ash, one of the nation's largest industrial waste streams, containing arsenic, cadmium, lead, mercury and radioactive elements. When stored in unlined ponds or landfills, these contaminants can leach into groundwater, pollute surface waters and contaminate drinking water supplies.⁸ Oil and gas refining produces chemicals that are used to manufacture plastics, pesticides and fertilizers, extending fossil fuel harms into consumer products. Many of these byproducts, such as plastics and agricultural chemicals, are associated with additional health harms from endocrine disruption to cancer (see Plastics section).



Exploration and production

On- and off-shore drilling, flaring,
solid and liquid waste



Transport, storage and processing

Storage tanks, transmission compression
stations, transportation



Refining, distribution, transmission

Refineries, oil and gas pipelines



Combustion and end use

Power plants, buildings and appliances,
vehicles



How Do Fossil Fuels Impact Our Health?

Fossil Fuels and Respiratory Health

The extraction, processing and burning of fossil fuels releases air pollutants that can penetrate deep into the lungs to cause inflammation, cell and tissue damage, immune system disorders, changes in the structure of the airways and chronic respiratory damage. Long-term exposure causes asthma and asthma attacks, chronic obstructive pulmonary disease (COPD), reduced lung function and lung development in children, increased respiratory infections, premature deaths from COPD and lung cancer.^{9,10}

Children are particularly vulnerable: they have greater exposure than adults as they spend more time outdoors, are more physically active and have faster respiratory rates. As children's lungs are still developing, they are more susceptible to damage from pollutants.^{11,12}



People living and working near the extraction or processing of fossil fuels also face significant risk. Coal miners can develop black lung disease, COPD and lung cancer from breathing in coal dust.^{13,14} Those living near coal-burning plants or coal ash storage facilities are at increased risk of respiratory symptoms (such as cough) and respiratory infections. Air pollution from oil and gas production (e.g. drilling and fracking) in one year alone resulted in 410,000 asthma flare-ups, 2,200 new cases of childhood asthma and 7,500 excess deaths.¹⁵ Residents living near mountaintop removal mining are exposed to pollutants such as particulate matter, polycyclic aromatic hydrocarbons (PAHs), heavy metals and hydrogen sulfide, linked to higher rates of birth defects, cardiovascular disease and respiratory illness.¹⁶

Reducing air pollution from fossil fuel facilities can deliver immediate and measurable health benefits. For instance, when a coal coking plant in Pittsburgh, Pennsylvania shut down, nearby communities saw a rapid decline in emergency department visits - over 20% fewer for respiratory symptoms and more than 40% fewer for pediatric asthma.¹⁷

Methane ("natural") gas is used to generate electricity and to power home appliances such as dryers, furnaces, ovens, stoves and water heaters.¹⁸ It is linked to an array of health problems including childhood cancer, birth defects and respiratory issues. Methane gas appliances (like gas stoves and furnaces) in our homes can emit carcinogenic benzene, carbon monoxide, nitrogen dioxide and fine particulate matter, increasing the risk of asthma and other respiratory symptoms, particularly in children and vulnerable populations.¹⁹

Burning fossil fuels releases air pollutants that lead to respiratory harm, such as particulate matter (PM2.5 and P10), ozone, nitrogen oxides (NO₂) and sulfur dioxide (SO₂).



Fueling Sickness: The Hidden Health Costs of Fossil Fuel Pollution

PM_{2.5}

Everyone in the U.S. is exposed to PM_{2.5} and at risk of its harmful effects, which accumulate over time. There is no safe threshold of exposure, meaning even low levels of exposure are associated with health risks. People with higher exposures – most often found in urban and low-income communities and communities of color – are most impacted.^{20,21} Short-term PM_{2.5} exposure can cause coughing, wheezing, phlegm production, increased respiratory infections, asthma attacks, increased emergency department visits and hospitalizations for respiratory conditions and deaths from respiratory illness. Long-term exposure is associated with significantly escalated risks of asthma, chronic respiratory diseases like COPD and pneumonia, lung cancer and premature death, even at low exposure levels routinely experienced in the US.²²

Ozone

Short-term ozone exposure (up to 8 hours) causes injury and inflammation of the respiratory tract leading to airway constriction and coughing. This results in decreased lung function, increased respiratory symptoms (such as coughing, throat irritation, chest discomfort, increased sensitivity to asthma triggers), emergency department visits and hospital admissions from asthma exacerbations and respiratory infections. Even healthy individuals have experienced decreased lung function and respiratory symptoms after short-term ozone exposure.²³ Long-term ozone exposure causes new-onset asthma as well as worsening symptoms in individuals with asthma. It is also associated with altered lung development, COPD and respiratory mortality.²⁴

NO₂

Short-term exposure to nitrogen dioxide causes development of asthma in children, increases airway responsiveness (or sensitivity) in adults with asthma, decreases lung function and increases respiratory-related hospital admissions and emergency department visits.²⁵ Long-term exposure can also decrease lung development in children and increase asthma or chronic bronchitis in adults.²⁶

SO₂

Exposure to sulfur dioxide worsens asthma. SO₂ may increase the risk of lung disease and death, especially in the elderly and people with chronic lung disease or cardiovascular disease.²⁷

Fossil Fuels and Cardiovascular Disease

Fossil fuels significantly increase the risk for cardiovascular diseases – heart attacks, strokes and heart failure – which are leading causes of death both in the U.S. and globally. Of the millions of deaths attributable to fossil fuel pollution, the greatest proportion are due to cardiovascular diseases.²⁸

Hundreds of studies have shown that short-term (hours to days) elevations in outdoor PM_{2.5} levels increase the risks for heart attacks, strokes and deaths from cardiovascular disease. Short-term exposure to particulate matter has also been linked to higher risks of atrial fibrillation, other arrhythmias and heart failure.²⁹ While brief exposures can be harmful, living in locations with poor air quality over the long term is related to much larger increases in cardiovascular risk. Fine particulate matter and NOx are consistently associated with hypertension, myocardial infarction, stroke and ischemic heart disease, while long-term exposures to PM_{2.5} increase the risk of atherosclerosis, incident stroke and stroke mortality.³⁰



Fueling Sickness: The Hidden Health Costs of Fossil Fuel Pollution

Fossil Fuel Use and Impacts on Brain Function

Air pollution from burning fossil fuels can harm the brain. Research shows that exposure to air pollutants - especially PM_{2.5}, NO₂ and ozone - can cause inflammation, stress and changes in brain chemistry. These changes can damage brain cells, interfere with how the brain communicates and increase the risk of neurological diseases such as Alzheimer's or Parkinson's.^{31,32}

Both short-term and long-term exposures to PM_{2.5} raise the risk of stroke.³³ Studies have found that people living in areas with higher levels of PM_{2.5}, NO₂ or ozone are more likely to experience strokes, hospitalization or premature death. A nationwide study of millions of Medicare enrollees found that reducing NO₂ levels by just 12.4 parts per billion - the difference between living in a large city like Los Angeles and a smaller one like Portland- was associated with a 6% reduction in stroke rates.³⁴

Long-term exposure to polluted air has also been linked to dementia and cognitive decline.³⁵ Another large study of Medicare beneficiaries found that every small increase in annual PM_{2.5} levels was associated with a higher risk of hospitalization for Alzheimer's, Parkinson's and other dementias, even at pollution levels below current air quality standards.³⁶



Children's developing brains are especially sensitive to fossil fuel pollution. Studies show that when pregnant women are exposed to high levels of air pollution, their children are more likely to have behavioral and learning problems. In one study, children exposed to more NO₂ before birth and PM_{2.5} in early childhood had more behavioral challenges and performed worse on cognitive tests at ages 4 to 6.³⁷ Other research has found that children exposed to polycyclic aromatic hydrocarbons (PAHs), compounds produced during incomplete fossil fuel combustion, have more difficulty with impulse control later in childhood.³⁸

Exposure to higher levels of air pollution has also been linked to poor cognitive function in adults. Although specific findings vary, a recent review of research found general agreement across studies that NO₂ and PM_{2.5} exposure is associated with lower cognitive performance in adults.³⁹

The neurocognitive harms of fossil fuel pollution disproportionately impact children, older adults and communities already facing environmental and health inequities. People living near busy roadways, diesel truck routes or polluting industries are exposed to higher levels of PM_{2.5} and other air pollutants. These communities are more likely to be low-income or communities of color, compounding the risks.

In children, exposure before and soon after birth can alter brain development, leading to smaller head circumference, reduced brain volumes and long-lasting neurologic effects.⁴⁰ Such exposures are associated with increased risks of autism spectrum disorder, attention deficit hyperactivity disorder (ADHD), behavioral problems, lower IQ and reduced academic performance.⁴¹ Alarmingly, studies have even found early signs of brain changes linked to Alzheimer's and Parkinson's disease in children exposed to high levels of air pollution.⁴²



Fueling Sickness: The Hidden Health Costs of Fossil Fuel Pollution

Fossil Fuels and Mental Health

A substantial and growing body of research demonstrates that fossil fuel pollution has adverse impacts on mental health. Exposure to pollutants such as $PM_{2.5}$ and NO_2 , can cause inflammation, stress and neurotoxicity on the brain and nervous system, which are linked to higher risks of depression, anxiety and other mental health challenges.

Both short-term and long-term exposure to air pollution have been connected to depression.⁴⁴ Risks of depression grow even larger with long-term exposure to elevated levels of $PM_{2.5}$ and NO_2 . Research has also found that long-term exposure to higher levels of PM_{10} and NO_2 is associated with higher risk of anxiety.⁴⁵

Air pollution has also been linked to suicide risk. In one study, short-term exposure to significant increases in particulate matter was found to be associated with a 1-2% increased risk of completing suicide⁴⁶, while a second found that increases in NO_2 were associated with a 3% increased risk.⁴⁷ Separate research has also found a connection between PM_{10} exposure and increased suicide risk over the subsequent 24-48 hours.⁴⁸

Children are especially vulnerable to the mental health effects of fossil fuel pollution.⁴⁹ Research has found that children who grow up breathing higher levels of air pollution are more likely to develop depression and behavioral disorders as teenagers. One study found that children experiencing the highest levels of air pollution at age 12 were up to 4 times more likely to be diagnosed with depression and up to 5 times more likely to be diagnosed with conduct disorder by age 18 - effects comparable to those seen from childhood abuse or trauma.⁵⁰

Pollution exposure may also play a role in more severe mental health conditions. While schizophrenia is largely influenced by genetics, environmental factors also contribute. Research found that people exposed to higher levels of NO_2 from birth through age 10 were 62% more likely to develop schizophrenia by age 37 compared to those growing up with cleaner air.⁵¹

Fossil Fuels and Reproductive Health

Exposure to fossil fuel pollution is linked to poor pregnancy health, pregnancy loss, fetal neurodevelopmental disruption and adverse birth outcomes such as preterm birth, low birth weight and birth defects.⁵² Inequitable distribution of fossil fuel pollution in the US is driving and worsening social and racial inequities in reproductive health.^{53,54,55}

Pregnant women and the developing fetus are especially vulnerable to the impacts of pollution because they experience many crucial and precisely timed physical and psychologic changes. When these processes are disrupted by pollutants, the impacts on pregnancy and the fetus can be profound and sometimes lifelong.⁵⁶ Due to pregnancy stress on the body, pregnant women are at additional risk of many serious diseases that can also be worsened by fossil fuel pollution, such as hypertension⁵⁷ and diabetes.⁵⁸ Pregnant women also breathe in more air than non-pregnant people, exposing them to more air pollution.⁵⁹

Numerous studies find convincing evidence that exposure to $PM_{2.5}$ during pregnancy raises the risks of unfavorable birth outcomes and pregnancy complications, including low birth weight, preterm birth, stillbirth, small for gestational age and birth defects.





Fueling Sickness: The Hidden Health Costs of Fossil Fuel Pollution

Pregnant women exposed to PM_{2.5} had a significantly higher risk of developing hypertensive disorder of pregnancy, gestational diabetes, gestational hypertension and preeclampsia.

Exposure to oil and gas well sites during pregnancy is associated with increased risk of spontaneous preterm birth.⁶⁰ Babies born in Pennsylvania within 1 km of a fracking site were more likely to have low birthweight⁶¹, and fracking has been associated with pre-term birth.⁶² Pregnant women living close to an oil or gas extraction site during pregnancy have an increased risk of hypertensive conditions, such as preeclampsia.⁶³

One of the strongest associations between air pollution and negative birth outcomes like low birth weight and preterm birth are with power plants and petrochemical industries.⁶⁴ Oil refineries produce a wide range of toxics, including benzene, which can harm fetal development and the male reproductive system; carbon monoxide, which during pregnancy can affect fetal brain development and cause pregnancy loss; and lead, a heavy metal that can adversely impact fetal brain developments, cause learning and behavioral problems in children and harm the reproductive systems of both men and women.⁶⁵

A study of exposure to traffic-related pollutants near the end of pregnancy showed that greater exposure was associated with an increase in NICU admissions.⁶⁶ Air pollution can disrupt fetal brain development.⁶⁷ People who live closer to fossil fuel power plants are more likely to have preterm delivery, even if they live up to 20 km (12.4 miles) away.⁶⁸



Other Reproductive Health Harms

Many petrochemicals are endocrine-disrupting chemicals (EDCs), which means they interfere with normal hormonal activity and increase risk for adverse health risks, including infertility.⁶⁹ A growing body of evidence also suggests that EDCs, including from fossil fuel-derived products are associated in studies with female reproductive health diseases such as painful and underdiagnosed conditions like fibroids, breast cancer and girls getting their periods increasingly earlier in life.⁷⁰

Exposure to endocrine disrupting chemicals is associated with reproductive harms (including reduced fertility in both women and men and adverse pregnancy outcomes).⁷¹ Consistent with these observations, women living near fracking sites have increased risks of preterm birth.⁷² Another report documented that women living near coal and oil power plants had a lower rate of preterm births after the plants were closed.⁷³

Fossil Fuels and Kidney Health

Fossil fuel pollution increases the risk of chronic kidney disease (CKD).⁷⁴ It has been estimated that the global burden of CKD attributable to PM_{2.5} is 6.95 million new cases of CKD each year.⁷⁵ The burden is greatest in low- and middle-income countries, where air pollution levels are higher and health risk factors like diabetes and high blood pressure are also more common. Both of these conditions, which can be made worse by fossil fuel pollution, increase the risk of developing CKD.^{76,77}

Both short- and longer-term exposure to PM_{2.5} are associated with significantly increased risks for development of acute kidney injury, progression of CKD and increased mortality in those with end stage kidney disease (ESKD).⁷⁸ Several studies



Fueling Sickness: The Hidden Health Costs of Fossil Fuel Pollution

have demonstrated an association between short-term elevations in PM_{2.5} and higher risk of death from kidney disease as well as emergency department visits.^{79,80} Short-term exposure to PM_{2.5} is also associated with an increased risk of hospital admissions and deaths in patients receiving dialysis for ESKD.⁸¹ A meta-analysis showed a higher risk for CKD and lower kidney function among people living near petrochemical plants than those without such exposure.⁸²

Fossil Fuels and Endocrine (Hormonal) Disease

Fossil-fuel driven air pollution poses substantial risks to the endocrine system, which controls growth and development, metabolism and reproduction. Studies have consistently demonstrated significant associations between levels of PM_{2.5} in the air and risks of obesity⁸³, Type 2 diabetes (and diabetes-associated mortality)⁸⁴ and high blood pressure.⁸⁵ Higher exposure to PM_{2.5} is also associated with higher risk for preterm birth, low birthweight, stillbirth⁸⁶ and preeclampsia.⁸⁷ While most of the evidence linking air pollution to endocrine system dysfunction is observational, a randomized trial in healthy adults showed significantly lower blood pressure, and lower levels of markers of stress and inflammation in healthy adults given air purifiers to reduce PM_{2.5} exposure.⁸⁸

Fossil fuel extraction releases many chemicals that interfere with normal function of the endocrine system. For example, hormone-disrupting chemicals are used in fracking for gas and often contaminate surface water, groundwater and even drinking water. A study of water samples from a region in Colorado with a high density of gas drilling and reports of spills and discharges found significantly elevated endocrine-disrupting activity compared with water from areas without gas drilling.⁸⁹ In addition, coal mining releases toxic metals (including lead, arsenic and mercury) that also act as endocrine disrupters.

Endocrine disrupting chemicals are also produced from fossil fuel products, such as microplastics/nanoplastics and PFAS, which have been detected in a variety of human tissues, including placenta.⁹⁰ Additionally, other fossil-fuel derived chemicals including pesticides act as endocrine disrupters.

Fossil Fuels and Cancer

Fossil fuel pollution is a contributor to the large burden of cancer in the U.S., affecting both the occurrence of cancer and its outcomes. About 2 million Americans will be diagnosed with cancer and more than 600,000 will die from cancer in 2025.⁹¹

Lung cancer is one of the most common cancers worldwide. It is estimated that approximately 15% of lung cancer worldwide is due to air pollution.⁹² The International Agency on Research on Cancer (IARC) has found PM_{2.5} and diesel exhaust are proven to cause cancer, damaging and mutating lung cells.

There is extensive evidence that people exposed to outdoor air pollution from car, truck and diesel exhaust have an increased risk for lung cancer and death.⁹³ Studies show that for each 10 mg/m³ increase in PM_{2.5} in the air, there is about a 13% increase in the risk for death from lung cancer.⁹⁴ Individuals who have never smoked can also face as much as a 27% increase in lung cancer deaths due to PM_{2.5}.⁹⁵ People of color who are diagnosed with lung cancer face worse outcomes compared to white Americans, including: less likely to be diagnosed early, less likely to receive surgical treatment, more likely to receive no treatment and less likely to survive five years.⁹⁶



Fueling Sickness: The Hidden Health Costs of Fossil Fuel Pollution

A growing body of evidence also shows an association between air pollution and increased risk for breast, colorectal, brain, bladder and liver cancers, including many studies linking traffic pollution with increased adult cancer risk.⁹⁷ Similarly, many studies have found an increased risk of brain, kidney, bone cancers and leukemia in children living near heavy-traffic roads or near industrial sites.^{98,99}

Workers and communities can be exposed to cancer-causing substances across the life cycle of fossil fuels. Fracking for oil and gas uses many chemicals - including benzene, 1,3-butadiene and formaldehyde - that are known human carcinogens. Children living near unconventional oil and gas development have shown an increased risk of leukemias and lymphomas and some studies suggest that it may be related to fracking.¹⁰⁰

Refinery and other petroleum workers may have repeated exposures to cancer-causing substances - including asbestos - and have been found to have increased risks of mesothelioma, lung cancer, malignant skin melanoma, acute lymphoid leukemia, multiple myeloma and bladder cancer.¹⁰¹ Residents in fence-line communities near petroleum industry sites and facilities also have elevated risks for leukemia.

Plastics and Health

Plastics are manufactured chemical products, more than 98% of which are made from coal, oil and gas. They persist in the environment for years to decades, contributing to a rapidly growing global pollution burden that now totals an estimated 8 billion tons. Despite efforts to promote plastics as recyclable, fewer than 10% of plastic (and only 1-2% of single use plastics) is recycled globally, leaving the vast majority to accumulate in landfills, waterways and ecosystems.¹⁰²

All plastics contain chemical additives, of which more than 10,000 are in use, and the vast majority have not been tested for toxicity. Of the chemicals that have been closely studied for health impacts, four major categories of chemicals are of particular concern: phthalates (used to soften plastic), bisphenols (BPA - used to produce linings), per- and polyfluoroalkyl substances (PFAS); and brominated flame-retardants.¹⁰³ These additives include known carcinogens (vinyl chloride, 1-3 butadiene and PFAS), neurotoxicants (lead and brominated and organophosphate flame retardants) and endocrine disruptors (phthalates and BPA). As these substances leach from plastics into the environment, they drive many of the health and ecological harms now associated with plastic pollution.

Plastics threaten human health across their entire life cycle – from fossil fuel extraction to production, use and disposal. Workers in fossil fuel extraction, plastic production, plastic textile workers and plastic recycling workers are all exposed to air pollutants and to multiple toxic chemicals. They suffer increased rates of cardiovascular, pulmonary, metabolic and neurologic diseases and cancer.¹⁰⁴



Microplastics and nanoplastics, very tiny plastic particles, have been detected in human hearts, brains and blood vessels, where they are associated with cardiovascular disease. Everyday exposures from food packaging, bottled water, indoor air and dust contribute to infertility, obesity, kidney disease, cancers and other chronic conditions. These chemicals disrupt endocrine function and increase risk for premature births, neurodevelopmental disorders, male reproductive birth defects, obesity, cardiovascular disease, renal disease and cancer. These particles are pervasive in indoor air, household dust, bottled



Fueling Sickness: The Hidden Health Costs of Fossil Fuel Pollution

water, tap water, meat, salt, fruits, vegetables, seafood, baby formula and breast milk.¹⁰⁵ When plastic is heated or damaged, bisphenol A (BPA) and phthalates can leach into food and beverages.¹⁰⁶

Infants and young children are especially vulnerable to plastics. Fetuses in the womb are exposed to plastics chemicals absorbed by their mothers. Prenatal exposures are linked to miscarriages, premature births, stillbirth, low birth weight, birth defects of the reproductive organs, neurodevelopmental impairment, impaired lung growth and childhood cancer. Early-life exposures to plastic chemicals increases risk of heart disease, Type-2 diabetes and obesity in childhood and across the life span.

The harms caused by plastics are not fairly distributed. People of color, Indigenous populations, low-income communities and workers in production and waste facilities face disproportionate exposures and health burdens including have increased risks of premature birth, low birth weight, asthma, leukemia, cardiovascular disease, chronic obstructive pulmonary disease and lung cancer.

People Disproportionately Impacted by the Health Harms of Fossil Fuels

While fossil fuel pollution harms everyone, its health impacts are not evenly distributed. Certain populations, including low-income communities, communities of color, children, older adults and people with preexisting health conditions, face disproportionate exposure and heightened vulnerability. These inequities reflect a combination of environmental, social and physiological factors that place greater health burdens on already at-risk groups.

Vulnerability due to exposure:

Low-income and Communities of Color:

A complex array of interconnected factors (including but not limited to historical redlining, lending discrimination, exclusionary land use policies, disinvestment and urban renewal projects) have resulted in low income and communities of color disproportionately living in areas with a greater concentration of polluting sources, like highways and factories, and thus have greater exposure to multiple pollutants from multiple sources.^{107,108} As a result, a higher percentage of racial minorities are exposed to particulate matter and ozone, contributing to a greater incidence of childhood asthma and other respiratory conditions. More than 1 in 5 African Americans live within a half-mile of an oil or gas production, processing or storage facility.¹⁰⁹ Low-income neighborhoods also tend to have older and less maintained housing that allow for greater penetration of outdoor air pollution into homes, making it difficult to escape the poor air quality.

Individuals Who are Unhoused

People experiencing homelessness are significantly more vulnerable to the adverse health impacts of fossil fuels due to increased exposure to air pollution. As many unhoused individuals seek shelter near highways¹¹⁰, they face long-term exposure to harmful pollutants such as particle pollution coming from car tailpipes.¹¹¹ Additionally, individuals who are unhoused are disproportionately impacted by other chronic and pre-existing conditions and may be less able to access care, which further increases their vulnerability to adverse health risks.¹¹²



Individuals Who Work Outdoors

People who work outdoors can face higher health risks due to their increased amount of time spent outside breathing in air pollution. Additionally, outdoor workers are often engaged in strenuous activity that increases their breathing rate and the amount of polluted air they inhale. Outdoor workers may also have limited options for reducing their exposure without jeopardizing their employment.

Workers in the Fossil Fuel Industry

Individuals who work in the fossil fuel industry face severe occupational health risks and disparities in health outcomes. Miners other than coal workers face disparities in other health outcomes, such as increased mortality from cardiovascular disease and several types of cancer, including mesothelioma.¹¹³ Workers in these industries also face substantial safety hazards, including but not limited to explosions and fires due to the ignition of flammable vapors or gases, falls from platforms or equipment and transportation issues. Highway vehicle crashes are the leading cause of death among oil and gas extraction worker fatalities.¹¹⁴

Vulnerability due to physiology:

Children

Children are uniquely vulnerable to the health harms of fossil fuels because their bodies, organs and immune systems are still developing. The impacts of pollution can exert multiple and cumulative adverse effects starting in utero and through childhood. Compared with adults, they spend more time outdoors, are more physically active and breathe more air in relation to their body size - factors that increase their uptake of pollutants that can penetrate deep into the lungs and bloodstream. Research has linked these exposures to a wide range of health effects in children, including low birth weight, asthma, reduced lung function, respiratory infections and allergies, as well as elevated risks for chronic disease later in life. Children living near fracking sites in Pennsylvania around their birth were two to three times more likely to be diagnosed with leukemia between the ages of 2-7 than children not living near these sites.¹¹⁵ Studies show that children's disproportionate burden stems from both biological and social vulnerabilities. Their developing lungs, brains and immune systems, along with immature detoxification pathways, make them less able to process and recover from toxic exposures.¹¹⁶ Disadvantaged and minority children are more likely to live in areas with higher concentrations of pollution, leaving them with elevated asthma rates and greater lifelong health challenges.^{117,118}

Older Adults

Older adults are more likely to have other pre-existing conditions, such as high blood pressure, that increase their risk of adverse impacts from air pollution. Air pollution can exacerbate COPD and increase the risk of heart attacks and strokes in older adults, especially those who are also obese and/or have diabetes. In terms of fossil fuel extraction, elderly people living near or downwind from gas extraction sites have a higher risk of premature death than seniors who do not.¹¹⁹

Individuals with Chronic Illness

People with asthma, COPD, cardiovascular disease, lung cancer, diabetes or obesity are at particularly high risk from particle pollution from fossil fuel combustion. Fossil fuel pollution causes cellular injury and inflammation that puts more stress on the lungs and heart and other systems already harmed by disease. People with chronic illness who are exposed to pollution may experience worse symptoms than those without, and have more emergency department visits, more hospitalizations and higher risk of premature death.¹²⁰



Fueling Sickness: The Hidden Health Costs of Fossil Fuel Pollution

Pregnant Women and Fetuses

Pregnant women are especially vulnerable to air pollution because they breathe more often. Pregnant women who live in counties with higher levels of fracking in Texas have been found to be at much higher risk of giving birth to children with specific birth defects.¹²¹ Additionally, another study found people living close to fracking sites in Pennsylvania had an increased risk of giving birth prematurely and of having high-risk pregnancies.¹²² Fetal development and health after birth may also be harmed by environmental contaminants that have been shown to cross the placenta.

Fossil Fuels, Climate Change and Health

Human activities are the cause of increased greenhouse gas emissions causing climate change, and burning fossil fuels is the single largest driver of climate change. Health professionals and scientists around the world agree that climate change is the biggest health threat of this century. Reducing fossil fuel pollution is the most effective way to protect current and future generations from the health harms of climate change.¹²³

Climate change causes warmer temperatures, more frequent extreme weather events, changes in drought and rainfall patterns and sea level rise. These changes affect the air, food, water and land on which human life depends, with health impacts already evident in communities across the U.S. Every small rise in global temperatures increases the likelihood of catastrophic health consequences, raising concern that our capacity to adapt or build resilience will be overwhelmed.

Climate Health Impacts

Heat-related deaths are increasing rapidly in the U.S. as extreme heat events become more frequent and severe. Extreme heat increases the risk of heart attack, worsens asthma and COPD, stresses the kidneys, increases the risk of complications from diabetes and worsens mental health. Scientists predict as many as tens of thousands of additional heat deaths in coming decades.¹²⁴

Hotter temperatures also worsen air quality due to the formation of ground-level ozone (smog). Wildfires, driven by hotter, drier conditions, expose millions to PM_{2.5} and toxic pollution that damages lungs, trigger cardiovascular disease, harm brain development and increase cancer risk. Warmer temperatures are even making allergy seasons longer.



Extreme weather events such as hurricanes, floods and severe storms cause injury and death, contaminate drinking water and overwhelm health systems. Power outages and facility closures disrupt medical care, including dialysis, insulin refrigeration and cancer treatment. Many patients - particularly older adults, people with disabilities and those without financial resources or transportation - face added risks because they may be unable to evacuate safely during floods, fires or smoke events.



Fueling Sickness: The Hidden Health Costs of Fossil Fuel Pollution

Climate change also impacts food. Rising temperatures cause increased risk of food poisoning, while droughts and floods can destroy crops, increase prices and worsen food insecurity. Climate change alters the distribution of mosquitoes and ticks, expanding risks for diseases like West Nile virus, dengue and Lyme disease.

Mental health is also deeply impacted. Nearly a third of people who experience a natural disaster may subsequently experience stress, PTSD and depression related to property damage, home loss or loss of loved ones.¹²⁵

Climate Change and Chronic Disease

Climate change compounds existing chronic health burdens, with extreme heat, wildfires, storms and other disasters worsening health outcomes. Heat stress, for example, raises the likelihood of heart attack, stroke, heart failure and dangerous arrhythmias. Heat also can lead to hospitalization for endocrine and metabolic conditions as well as adverse pregnancy outcomes including preterm birth, low birthweight and stillbirth.¹²⁶ Wildfire smoke, extreme cold and hurricanes have also been linked to higher rates of heart disease, stroke and cardiovascular deaths. Cancer risks also rise with increased wildfire smoke exposure.¹²⁷ The increase in vector-borne diseases aggravates kidney disease and may cause endocrine disturbances such as hypoglycemia and hyponatremia. Kidney disease is also worsened by extreme heat and dehydration.

Extreme weather and disasters also disrupt access to critical healthcare, particularly for those who rely on steady, reliable treatments for their chronic condition. For example, after Hurricane Katrina, women with breast cancer and patients with lung cancer experienced worse long-term survival due to disruptions in care.¹²⁸ Similarly, both Hurricane Katrina and Superstorm Sandy were followed by increases in heart attack incidence and mortality.¹²⁹ Heat waves have also been shown to disrupt cancer care delivery, further highlighting how climate-related events can endanger patients who rely on timely medical treatment.¹³⁰

Climate Vulnerability

While climate change threatens everyone's health, some populations bear greater burdens, which exacerbates existing health inequities. Children, pregnant women, the elderly and people with chronic illness and disabilities are more vulnerable to climate health threats. Low-income communities and communities of color face higher exposures and fewer resources to adapt, compounded by historic disinvestment, less green space and higher baseline pollution. Outdoor workers, residents of hotter regions and fence line communities near industrial or transportation facilities are also disproportionately affected.

Reducing Climate Health Impacts

The most powerful health protection strategy is to reduce climate pollution by transitioning away from fossil fuels. The recommendations listed below to reduce pollution from fossil fuels will also deliver long-term benefits of mitigating climate change. At the same time, communities can build climate resilience through cooling centers, greening neighborhoods and weatherizing homes as well as implementing early warning systems and stronger healthcare infrastructure. These measures reduce risk from current climate impacts while we work to prevent even greater harm.



Fueling Sickness: The Hidden Health Costs of Fossil Fuel Pollution

Conclusions and Recommendations

Fossil fuels harm nearly every system of the body across the human life cycle both in the U.S. and worldwide. Fossil fuels are also the leading driver of climate change, which the healthcare community recognizes as the greatest health challenge of this century.

Extensive medical and scientific evidence makes clear: to safeguard health today and for future generations, we must transition away from fossil fuels toward clean, non-polluting energy sources as quickly as possible. This transition is not only possible but increasingly practical thanks to affordable clean energy technologies. The health benefits of rapid action are immediate and profound. Each step away from fossil fuels reduces air pollution, prevents disease and saves lives, while also delivering cost savings through reduced healthcare burdens.

As health and medical professionals and organizations, it is our duty not only to treat those already harmed by fossil fuel pollution, but also to speak out about its dangers. We must inform the public and policymakers about the health risks, and advocate for policies that reduce exposure to fossil fuel pollution for our patients and communities.

Individuals can also play a vital role by engaging with local organizations that are advocating for clean air and a clean energy transition, talking to community members and policymakers about the health harms of fossil fuel pollution and reducing personal reliance on fossil fuels where possible.



But individuals and health professionals alone are not enough. Protecting communities from the health harms of fossil fuel pollution requires bold policy and systemic change.

Recommendations for Policymakers

- Transition rapidly from fossil fuels to clean, non-polluting energy and transportation.
- Protect communities from exposure to fossil fuel pollution by protecting, enforcing and strengthening clean air and water standards.
- Implement a public health campaign to educate the public about the health harms of fossil fuel pollution, even at low, everyday levels of exposure.
- Hold the fossil fuel industry accountable for its health harms.
- Invest in public health systems, healthcare infrastructure, research and innovation to protect and promote health and accelerate clean technology progress.

The costs of inaction are measured in illness, lives lost and mounting healthcare burdens. The benefits of a clean energy transition are measured in healthier communities and a safer future for our children and grandchildren. Protecting health demands nothing less than urgent action.



Fueling Sickness: The Hidden Health Costs of Fossil Fuel Pollution

References:

Environmental Protection Agency. Integrated Science Assessment (ISA) for Particulate Matter. January 2025. [Integrated Science Assessment \(ISA\) for Particulate Matter | US EPA](#)

² Harvard Health Publishing. Harvard Medical School. Even low-level air pollution may harm health. February 2022. [Even low-level air pollution may harm health - Harvard Health](#)

³ Harvard T.H. Chan School of Public Health. Fossil fuel air pollution responsible for 1 in 5 deaths worldwide. February 2021. [Fossil fuel air pollution responsible for 1 in 5 deaths worldwide | Harvard T.H. Chan School of Public Health](#)

⁴ Lelieveld, J., Evans, J. S., Fnais, M., Giannadaki, D., & Pozzer, A. (2019). Effects of fossil fuel and total anthropogenic emission removal on public health and climate. *Proceedings of the National Academy of Sciences*, 116(15), 7192–7197. [Effects of fossil fuel and total anthropogenic emission removal on public health and climate | PNAS](#)

⁵ Karn Vohra et al., The health burden and racial-ethnic disparities of air pollution from the major oil and gas lifecycle stages in the United States. *Sci. Adv.* 11, eadu2241 (2025). DOI: [10.1126/sciadv.adu2241](#)

⁶ Vimercati L. Traffic related air pollution and respiratory morbidity. *Lung India*. 2011 Oct; 28(4):238. doi: [10.4103/0970-2113.85682](#). PMID: 22084534; PMCID: PMC3213707.

⁷ Health Effects Institute. Special Report 23. Systematic Review and Meta-analysis of Selected Health Effects of Long-Term Exposure to Traffic-Related Air Pollution. April 2023. [https://www.healtheffects.org/publication/systematic-review-and-meta-analysis-selected-health-effects-long-term-exposure-traffic](#)

⁸ Sears CG, Zierold KM. Health of Children Living Near Coal Ash. *Glob Pediatr Health*. 2017 Jul 25;4:2333794X17720330. doi: [10.1177/2333794X17720330](#). PMID: 28804748; PMCID: PMC5533260. [Health of Children Living Near Coal Ash - PMC](#)

⁹ Jiang XQ, Mei XD, Feng D. Air pollution and chronic airway diseases: what should people know and do? *J Thorac Dis*. 2016 Jan;8(1):E31-40. doi: [10.3978/j.issn.2072-1439.2015.11.50](#). PMID: 26904251; PMCID: PMC4740163. [Air pollution and chronic airway diseases: what should people know and do? - PMC](#)

¹⁰ American Lung Association. State of the Air 2025. Health Impact of Air Pollution. [Health Impact of Pollution | State of the Air | American Lung Association](#)

¹¹ Su JG, Aslebagh S, Shahriary E, Barrett M, Balmes JR. Impacts from air pollution on respiratory disease outcomes: a meta-analysis. *Front Public Health*. 2024 Oct 9;12:1417450. doi: [10.3389/fpubh.2024.1417450](#). PMID: 39444957; PMCID: PMC11497638. [Impacts from air pollution on respiratory disease outcomes: a meta-analysis - PMC](#)

¹² Garcia E, Rice MB, Gold DR. Air pollution and lung function in children. *J Allergy Clin Immunol*. 2021 Jul;148(1):1-14. doi: [10.1016/j.jaci.2021.05.006](#). PMID: 34238501; PMCID: PMC8274324. [Air pollution and lung function in children - PMC](#)

¹³ Laney AS, Weissman DN. Respiratory diseases caused by coal mine dust. *J Occup Environ Med*. 2014 Oct;56 Suppl 10(0 10):S18-22. doi: [10.1097/JOM.0000000000000260](#). PMID: 25285970; PMCID: PMC4556416. [Respiratory Diseases Caused by Coal Mine Dust - PMC](#)

¹⁴ Centers for Disease Control and Prevention. Modern Coal Miners Have Higher Death Rates From Lung Diseases Than Their Predecessors. February 2023. [Modern Coal Miners Have Higher Death Rates From Lung Diseases Than Their Predecessors | Blogs | CDC](#)

¹⁵ Jonathan J Buonocore et al 2023 *Environ. Res.: Health* 1 021006 [Air pollution and health impacts of oil & gas production in the United States - IOPscience](#)

¹⁶ Boyles AL, Blain RB, Rochester JR, Avanasi R, Goldhaber SB, McComb S, Holmgren SD, Masten SA, Thayer KA. Systematic review of community health impacts of mountaintop removal mining. *Environ Int*. 2017 Oct;107:163-172. doi: [10.1016/j.envint.2017.07.002](#). Epub 2017 Jul 21. PMID: 28738262; PMCID: PMC5562233. [Systematic Review of Community Health Impacts of Mountaintop Removal Mining - PMC](#)

¹⁷ American Journal of Respiratory and Critical Care Medicine. Reductions in Respiratory Hospital Visits after a Coal Coking Plant Closure: A Natural Experiment. October 2024. [Reductions in Respiratory Hospital Visits after a Coal Coking Plant Closure: A Natural Experiment | American Journal of Respiratory and Critical Care Medicine | Articles in Press](#)



Fueling Sickness: The Hidden Health Costs of Fossil Fuel Pollution

¹⁸ US Energy Information Administration. Use of natural gas-fired generation differs in the United States by technology and region. February 2024. [Use of natural gas-fired generation differs in the United States by technology and region - U.S. Energy Information Administration \(EIA\)](#)

¹⁹ Kashtan YS, Nicholson M, Finnegan C, Ouyang Z, Lebel ED, Michanowicz DR, Shonkoff SBC, Jackson RB. Gas and Propane Combustion from Stoves Emits Benzene and Increases Indoor Air Pollution. *Environ Sci Technol.* 2023 Jul 4;57(26):9653-9663. doi: [10.1021/acs.est.2c09289](https://doi.org/10.1021/acs.est.2c09289). Epub 2023 Jun 15. PMID: 37319002; PMCID: PMC10324305.

²⁰ Liu J, Clark LP, Bechle MJ, Hajat A, Kim S-Y, Robinson AL, Sheppard L, Szpiro AA, Marshall JD. Disparities in air pollution exposure in the United States by race/ethnicity and income, 1990–2010. *Environ Health Perspect.* 2021; 129(12). [Disparities in Air Pollution Exposure in the United States by Race/Ethnicity and Income, 1990–2010 | Environmental Health Perspectives | Vol. 129, No. 12](#)

²¹ Martenies SE, Milando CW, Williams GO, Batterman SA. Disease and Health Inequalities Attributable to Air Pollutant Exposure in Detroit, Michigan. *Int J Environ Res Public Health.* 2017 Oct 19;14(10):1243. doi: 10.3390/ijerph14101243. PMID: 29048385; PMCID: PMC5664744. [Disease and Health Inequalities Attributable to Air Pollutant Exposure in Detroit, Michigan - PMC](#)

²² US Environmental Protection Agency. Supplement to the 2019 Integrated Science Assessment for Particulate Matter (Final Report, 2022). May 2024. [Supplement to the 2019 Integrated Science Assessment for Particulate Matter \(Final Report, 2022\) | Integrated Science Assessments | Environmental Assessment | US EPA](#)

²³ US Environmental Protection Agency. Health Effects of Ozone in the General Population. March 2025. [Health Effects of Ozone in the General Population | US EPA](#)

²⁴ US Environmental Protection Agency. Integrated Science Assessment for Ozone and Related Photochemical Oxidants. April 2020. https://ordspub.epa.gov/ords/eims/eimscomm.getfile?p_download_id=540022;

²⁵ US Environmental Protection Agency. Integrated Science Assessment for Oxides of Nitrogen - Health Criteria. January 2016. https://ordspub.epa.gov/ords/eims/eimscomm.getfile?p_download_id=526855

²⁶ Anenberg, Susan C.; Mohegh, Arash; Goldberg, Daniel L.; Kerr, Gaige H.; Brauer, Michael; Burkart, Katrin; Hystad, Perry; Larkin, Andrew; Wozniak, Sarah; and Lamsal, Lok, “Long-term trends in urban NO₂ concentrations and associated paediatric asthma incidence: estimates from global datasets” (2022). GW Authored Works. Paper 65. [“Long-term trends in urban NO₂ concentrations and associated paediatric” by Susan C. Anenberg, Arash Mohegh et al.](#)

²⁷ California Air Resources Board. [Sulfur Dioxide & Health. Sulfur Dioxide & Health | California Air Resources Board](#)

²⁸ Murray, Christopher J L et al. The Lancet, Volume 396, Issue 10258, 1223 - 1249. Global burden of 87 risk factors in 204 countries and territories, 1990–2019: a systematic analysis for the Global Burden of Disease Study 2019. [Global burden of 87 risk factors in 204 countries and territories, 1990–2019: a systematic analysis for the Global Burden of Disease Study 2019 - The Lancet](#)

²⁹ Orellano P, Reynoso J, Quaranta N, Bardach A, Ciapponi A. Short-term exposure to particulate matter (PM₁₀ and PM_{2.5}), nitrogen dioxide (NO₂), and ozone (O₃) and all-cause and cause-specific mortality: Systematic review and meta-analysis. *Environ Int.* 2020 Sep;142:105876. doi: [10.1016/j.envint.2020.105876](https://doi.org/10.1016/j.envint.2020.105876). Epub 2020 Jun 23. PMID: 32590284.

³⁰ Alexeeff, Stacey E., Noelle S. Liao, Xi Liu, Stephen K. Van Den Eeden, and Stephen Sidney. 2021. “Long-Term PM_{2.5} Exposure and Risks of Ischemic Heart Disease and Stroke Events: Review and Meta-Analysis.” *Journal of the American Heart Association* 10 (1): e016890. <https://doi.org/10.1161/JAHA.120.016890>

³¹ Block ML, Calderón-Garcidueñas L. Air pollution: mechanisms of neuroinflammation and CNS disease. *Trends Neurosci.* 2009 Sep;32(9):506-16. doi: 10.1016/j.tins.2009.05.009. Epub 2009 Aug 26. PMID: 19716187; PMCID: PMC2743793. [Air pollution: mechanisms of neuroinflammation and CNS disease - PubMed](#)

³² Costa LG, Cole TB, Dao K, Chang YC, Coburn J, Garrick JM. Effects of air pollution on the nervous system and its possible role in neurodevelopmental and neurodegenerative disorders. *Pharmacol Ther.* 2020 Jun;210:107523. doi: 10.1016/j.pharmthera.2020.107523. Epub 2020 Mar 9. PMID: 32165138; PMCID: PMC7245732. [Effects of air pollution on the nervous system and its possible role in neurodevelopmental and neurodegenerative disorders - PMC](#)



Fueling Sickness: The Hidden Health Costs of Fossil Fuel Pollution

³³ Kulick ER, Kaufman JD, Sack C. Ambient Air Pollution and Stroke: An Updated Review. *Stroke*. 2023 Mar;54(3):882-893. doi: 10.1161/STROKEAHA.122.035498. Epub 2022 Dec 29. PMID: 36579640; PMCID: PMC10421613. [Ambient Air Pollution and Stroke – An Updated Review - PMC](#)

³⁴ Ma T, Yazdi MD, Schwartz J, Réquia WJ, Di Q, Wei Y, Chang HH, Vaccarino V, Liu P, Shi L. Long-term air pollution exposure and incident stroke in American older adults: A national cohort study. *Glob Epidemiol*. 2022 Dec;4:100073. doi: 10.1016/j.gloepi.2022.100073. Epub 2022 Apr 16. PMID: 36644436; PMCID: PMC9838077. [Long-term air pollution exposure and incident stroke in American older adults: A national cohort study - PMC](#)

³⁵ Delgado-Saborit JM, Guercio V, Gowers AM, Shaddick G, Fox NC, Love S. A critical review of the epidemiological evidence of effects of air pollution on dementia, cognitive function and cognitive decline in adult population. *Sci Total Environ*. 2021 Feb 25;757:143734. doi: 10.1016/j.scitotenv.2020.143734. Epub 2020 Nov 25. PMID: 33340865. [A critical review of the epidemiological evidence of effects of air pollution on dementia, cognitive function and cognitive decline in adult population - PubMed](#)

³⁶ Shi L, Wu X, Danesh Yazdi M, Braun D, Abu Awad Y, Wei Y, Liu P, Di Q, Wang Y, Schwartz J, Dominici F, Kioumourtzoglou MA, Zanobetti A. Long-term effects of PM_{2.5} on neurological disorders in the American Medicare population: a longitudinal cohort study. *Lancet Planet Health*. 2020 Dec;4(12):e557-e565. doi: 10.1016/S2542-5196(20)30227-8. Epub 2020 Oct 19. PMID: 33091388; PMCID: PMC7720425. [Long-term effects of PM2.5 on neurological disorders in the American Medicare population: a longitudinal cohort study - PMC](#)

³⁷ Ni Y, Loftus CT, Szpiro AA, Young MT, Hazlehurst MF, Murphy LE, Tylavsky FA, Mason WA, LeWinn KZ, Sathyaranayana S, Barrett ES, Bush NR, Karr CJ. Associations of Pre- and Postnatal Air Pollution Exposures with Child Behavioral Problems and Cognitive Performance: A U.S. Multi-Cohort Study. *Environ Health Perspect*. 2022 Jun;130(6):67008. doi: 10.1289/EHP10248. Epub 2022 Jun 23. PMID: 35737514; PMCID: PMC9222764. [Associations of Pre- and Postnatal Air Pollution Exposures with Child Behavioral Problems and Cognitive Performance: A U.S. Multi-Cohort Study - PMC](#)

³⁸ Amy E. Margolis, Bruce Ramphal, David Pagliaccio, Sarah Bunker, Ena Selmanovic, Lauren V. Thomas, Pam Factor-Litvak, Frederica Perera, Bradley S. Peterson, Andrew Rundle, Julie B. Herbstman, Jeff Goldsmith, Virginia Rauh, Prenatal exposure to air pollution is associated with childhood inhibitory control and adolescent academic achievement, *Environmental Research*, Volume 202, 2021, 111570, ISSN 0013-9351, <https://doi.org/10.1016/j.envres.2021.111570>

³⁹ Thompson R, Smith RB, Karim YB, Shen C, Drummond K, Teng C, Toledo MB. Air pollution and human cognition: A systematic review and meta-analysis. *Sci Total Environ*. 2023 Feb 10;859(Pt 2):160234. doi: 10.1016/j.scitotenv.2022.160234. Epub 2022 Nov 23. PMID: 36427724. [Air pollution and human cognition: A systematic review and meta-analysis - PubMed](#)

⁴⁰ Peterson BS, Bansal R, Sawardekar S, Nati C, Elgabalawy ER, Hoepner LA, Garcia W, Hao X, Margolis A, Perera F, Rauh V. Prenatal exposure to air pollution is associated with altered brain structure, function, and metabolism in childhood. *J Child Psychol Psychiatry*. 2022 Nov;63(11):1316-1331. doi: 10.1111/jcpp.13578. Epub 2022 Feb 14. PMID: 35165899. [Prenatal exposure to air pollution is associated with altered brain structure, function, and metabolism in childhood - PubMed](#)

⁴¹ Kaur S, Morales-Hidalgo P, Arija V, Canals J. Prenatal Exposure to Air Pollutants and Attentional Deficit Hyperactivity Disorder Development in Children: A Systematic Review. *Int J Environ Res Public Health*. 2023 Apr 7;20(8):5443. doi: 10.3390/ijerph20085443. PMID: 37107725; PMCID: PMC10138804. [Prenatal Exposure to Air Pollutants and Attentional Deficit Hyperactivity Disorder Development in Children: A Systematic Review - PMC](#)

⁴² Lilian Calderón-Garcidueñas, Angélica González-Maciel, Rafael Reynoso-Robles, Jessica Hammond, Randy Kulesza, Ingolf Lachmann, Ricardo Torres-Jardón, Partha S. Mukherjee, Barbara A. Maher, Quadruple abnormal protein aggregates in brain-stem pathology and exogenous metal-rich magnetic nanoparticles (and engineered Ti-rich nanorods). The substantia nigrae is a very early target in young urbanites and the gastrointestinal tract a key brainstem portal, *Environmental Research*, Volume 191, 2020, 110139, ISSN 0013-9351, <https://doi.org/10.1016/j.envres.2020.110139>.

⁴³ Fu, et al. The association between PM 2.5 exposure and neurological disorders: A systematic review and meta-analysis. *Sci Total Environ* 2019 Mar 10;655:1240-1248 [The association between PM2.5 exposure and neurological disorders: A systematic review and meta-analysis - PubMed](#)

⁴⁴ Borroni E, Pesatori AC, Bollati V, Buoli M, Carugno M. Air pollution exposure and depression: A comprehensive updated systematic review and meta-analysis. *Environ Pollut*. 2022 Jan 1;292(Pt A):118245. doi: 10.1016/j.envpol.2021.118245. Epub 2021 Sep 29. PMID: 34600062. [Air pollution exposure and depression: A comprehensive updated systematic review and meta-analysis - PubMed](#)



Fueling Sickness: The Hidden Health Costs of Fossil Fuel Pollution

⁴⁵ Federica Nobile, Anna Forastiere, Paola Michelozzi, Francesco Forastiere, Massimo Stafoggia, Long-term exposure to air pollution and incidence of mental disorders. A large longitudinal cohort study of adults within an urban area, *Environment International*, Volume 181, 2023, 108302, ISSN 0160-4120, <https://doi.org/10.1016/j.envint.2023.108302>.

⁴⁶ Liu Q, Wang W, Gu X, Deng F, Wang X, Lin H, Guo X, Wu S. Association between particulate matter air pollution and risk of depression and suicide: a systematic review and meta-analysis. *Environ Sci Pollut Res Int*. 2021 Feb;28(8):9029-9049. doi: 10.1007/s11356-021-12357-3. Epub 2021 Jan 22. PMID: 33481201. [Association between particulate matter air pollution and risk of depression and suicide: a systematic review and meta-analysis - PubMed](#)

⁴⁷ Heo S, Lee W, Bell ML. Suicide and Associations with Air Pollution and Ambient Temperature: A Systematic Review and Meta-Analysis. *Int J Environ Res Public Health*. 2021 Jul 20;18(14):7699. doi: 10.3390/ijerph18147699. PMID: 34300149; PMCID: PMC8303705. [Suicide and Associations with Air Pollution and Ambient Temperature: A Systematic Review and Meta-Analysis - PubMed](#)

⁴⁸ Braithwaite I, Zhang S, Kirkbride JB, Osborn DPJ, Hayes JF. Air Pollution (Particulate Matter) Exposure and Associations with Depression, Anxiety, Bipolar, Psychosis and Suicide Risk: A Systematic Review and Meta-Analysis. *Environ Health Perspect*. 2019 Dec;127(12):126002. doi: 10.1289/EHP4595. Epub 2019 Dec 18. PMID: 31850801; PMCID: PMC6957283. [Air Pollution \(Particulate Matter\) Exposure and Associations with Depression, Anxiety, Bipolar, Psychosis and Suicide Risk: A Systematic Review and Meta-Analysis - PubMed](#)

⁴⁹ Zundel CG, Ryan P, Brokamp C, Heeter A, Huang Y, Strawn JR, Marusak HA. Air pollution, depressive and anxiety disorders, and brain effects: A systematic review. *Neurotoxicology*. 2022 Dec;93:272-300. doi: 10.1016/j.neuro.2022.10.011. Epub 2022 Oct 21. PMID: 36280190; PMCID: PMC10015654. [Air Pollution, Depressive and Anxiety Disorders, and Brain Effects: A Systematic Review - PMC](#)

⁵⁰ Roberts S, Arseneault L, Barratt B, Beevers S, Danese A, Odgers CL, Moffitt TE, Reuben A, Kelly FJ, Fisher HL. Exploration of NO₂ and PM_{2.5} air pollution and mental health problems using high-resolution data in London-based children from a UK longitudinal cohort study. *Psychiatry Res*. 2019 Feb;272:8-17. doi: 10.1016/j.psychres.2018.12.050. Epub 2018 Dec 10. PMID: 30576995; PMCID: PMC6401205. [Exploration of NO₂ and PM_{2.5} air pollution and mental health problems using high-resolution data in London-based children from a UK longitudinal cohort study - PMC](#)

⁵¹ Antonsen S, Mok PLH, Webb RT, Mortensen PB, McGrath JJ, Agerbo E, Brandt J, Geels C, Christensen JH, Pedersen CB. Exposure to air pollution during childhood and risk of developing schizophrenia: a national cohort study. *Lancet Planet Health*. 2020 Feb;4(2):e64-e73. doi: 10.1016/S2542-5196(20)30004-8. Epub 2020 Feb 26. PMID: 32112749. [Exposure to air pollution during childhood and risk of developing schizophrenia: a national cohort study - PubMed](#)

⁵² March of Dimes, “Healthy Environment for a Healthy Start: Promoting Environmental Justice for Equitable Birth Outcomes”, especially, “Brief 1: Air pollution worsens inequities in maternal and newborn health” and “Brief 5: The fossil fuel cycle worsens maternal and infant health inequities” 2024. <https://ignitingimpacttogether.marchofdimes.org/environmentaljustice> (accessed July 30, 2025).

⁵³ Trowbridge, Jessica, Goin, Dana E, Sklar, Rachel et al. Fossil fuel is the common denominator between climate change and petrochemical exposures, and effects on women and children’s health, <https://doi.org/10.1002/ijgo.14408>, September 7, 2022. *International Journal of Gynecology and Obstetrics*.

⁵⁴ Reducing Prenatal Exposure to Toxic Environmental Agents, ACOG Committee Opinion, Number 832, *Obstetrics & Gynecology* [138\(1\):p e40-e54, July 2021](https://doi.org/10.1002/jgo.14408). DOI: 10.1097/AOG.0000000000004449

⁵⁵ Women’s Earth and Climate Action Network (WECAN), “Gendered and Racial Impacts of the Fossil Fuel Industry in North America and Complicit Financial Institutions”, 4th Edition, September 2024, <https://www.wecaninternational.org/divestment-report>, (accessed July 30, 2025).

⁵⁶ Animal studies suggest that even very low levels of toxics if administered at crucial moments in fetal development can change how an organism is constructed and functions throughout its life. See TedX, The Endocrine Disruption Exchange, “The Fossil Fuel Connection”, fact sheet [The fossil fuel connection 6-3-19.pdf](https://tedxendocrinodisruption.org/wp-content/uploads/2019/04/The-fossil-fuel-connection-6-3-19.pdf). See also Guidance from the Western States Pediatric Environmental Health Specialty Unit, Climate Change and Pregnancy, undated, https://wspehsu.ucsf.edu/wp-content/uploads/2025/01/Climate-Change-and-Pregnancy-factsheet_04_08_25_II.pdf



Fueling Sickness: The Hidden Health Costs of Fossil Fuel Pollution

⁵⁷ Hoimonty Mazumder, Fariha Hoque Rimu, Monir Hossain Shimul, Jyoti Das, Easter Protiva Gain, Winston Liaw, M. Mahbub Hossain, Maternal health outcomes associated with ambient air pollution: An umbrella review of systematic reviews and meta-analyses, *Science of The Total Environment*, Volume 914, 2024, 169792, ISSN 0048-9697, <https://doi.org/10.1016/j.scitotenv.2023.169792>.

⁵⁸ Nazarpour S, Ramezani Tehrani F, Valizadeh R, Amiri M. The relationship between air pollutants and gestational diabetes: an updated systematic review and meta-analysis. *J Endocrinol Invest*. 2023 Jul;46(7):1317-1332. doi: [10.1007/s40618-023-02037-z](https://doi.org/10.1007/s40618-023-02037-z). Epub 2023 Feb 18. PMID: 36807891

⁵⁹ Guidance from the Western States Pediatric Environmental Health Specialty Unit, Climate Change and Pregnancy, undated, https://wspehsu.ucsf.edu/wp-content/uploads/2025/01/Climate-Change-and-Pregnancy-factsheet_04_08_25_ll.pdf

⁶⁰ Gonzalez, David J. X.a; Sherris, Allison R.a; Yang, Weib; Stevenson, David K.b; Padula, Amy M.c; Baiocchi, Michaeld; Burke, Marshalle; Cullen, Mark R.f; Shaw, Gary M.b. Oil and gas production and spontaneous preterm birth in the San Joaquin Valley, CA: A case-control study. *Environmental Epidemiology* 4(4): p e099, August 2020. DOI: [10.1097/EE.0000000000000099](https://doi.org/10.1097/EE.0000000000000099)

⁶¹ Liu J, Clark LP, Bechle MJ, Hajat A, Kim S-Y, Robinson AL, Sheppard L, Szpiro AA, Marshall JD. Disparities in Air Pollution Exposure in the United States by Race/ Ethnicity and Income, 1990–2010. 2021 *Environmental Health Perspectives* 129:12 CID: 127005 DOI: [10.1289/EHP8584](https://doi.org/10.1289/EHP8584)

⁶² Deziel, N. C., Brokovich, E., Grotto, I., Clark, C. J., Barnett-Itzhaki, Z., Broday, D., & Agay-Shay, K. (2020). Unconventional oil and gas development and health outcomes: A scoping review of the epidemiological research. *Environmental research*, 182, 109124. <https://doi.org/10.1016/j.envres.2020.109124>

⁶³ Mary D Willis, Elaine L Hill, Molly L Kile, Susan Carozza, Perry Hystad, Associations between residential proximity to oil and gas extraction and hypertensive conditions during pregnancy: a difference-in-differences analysis in Texas, 1996–2009, *International Journal of Epidemiology*, Volume 51, Issue 2, April 2022, Pages 525–536, <https://doi.org/10.1093/ije/dyab246>

⁶⁴ Veber T, Dahal U, Lang K, Orru K, Orru H. Industrial Air Pollution Leads to Adverse Birth Outcomes: A Systematized Review of Different Exposure Metrics and Health Effects in Newborns. *Public Health Rev*. 2022 Aug 10;43:1604775. doi: [10.3389/phrs.2022.1604775](https://doi.org/10.3389/phrs.2022.1604775). PMID: 36035982; PMCID: PMC9400407.

⁶⁵ Proposition 65 Warnings Office of Environmental Health Hazard Assessment, California, “Petroleum Products Environmental Exposure from Refineries” factsheet, https://www.p65warnings.ca.gov/sites/default/files/downloads/factsheets/petroleum_products_fs_-_refinery.pdf

⁶⁶ Phiri, Y.V.A., Canty, T., Nobles, C. et al. Neonatal intensive care admissions and exposure to satellite-derived air pollutants in the United States, 2018. *Sci Rep* 15, 420 (2025). <https://doi.org/10.1038/s41598-024-84755-9> SEE ALSO Indulaxmi Seen, Andrew Williams, Carrie Nobles, Zhen Chen, Seth Sherman, Pauline Mendola, Acute air pollution exposure and NICU admission: a case-crossover analysis, *Annals of Epidemiology*, Volume 37, 2019, Pages 64-70.e2, ISSN 1047-2797, <https://doi.org/10.1016/j.anepidem.2019.07.008>.

⁶⁷ Air pollution and fetal brain morphological development: a prospective cohort study, Gómez-Herrera, Laura et al. *The Lancet Planetary Health*, Volume 9, Issue 6, e480 - e490. [Air pollution and fetal brain morphological development: a prospective cohort study - The Lancet Planetary Health](https://doi.org/10.1016/j.lanepe.2025.e480)

⁶⁸ Sandie Ha and others, Associations Between Residential Proximity to Power Plants and Adverse Birth Outcomes, *American Journal of Epidemiology*, Volume 182, Issue 3, 1 August 2015, Pages 215–224, <https://doi.org/10.1093/aje/kwv042>

⁶⁹ Woodruff, Tracey J. Health Effects of Fossil Fuel-Derived Endocrine Disruptors, March 6, 2024, *The New England Journal of Medicine*, 2024;390:922-933 DOI: [10.1056/NEJMra2300476](https://doi.org/10.1056/NEJMra2300476)

⁷⁰ Gore, A.C., La Merrill, M.A., Patisaul, H.B., and Sargis, R. Endocrine Disrupting Chemicals: Threats to Human Health. The Endocrine Society and IPEN. February 2024. [edc_report-2024-final-compressed.pdf](https://doi.org/10.1016/j.jen.2024.01.001)

⁷¹ Woodruff T. Health Effects of Fossil Fuel-Derived Endocrine Disruptors. *N Engl J Med* 2024;390:922-933. doi: [10.1056/NEJMra2300476](https://doi.org/10.1056/NEJMra2300476). PMID: 38446677.



Fueling Sickness: The Hidden Health Costs of Fossil Fuel Pollution

⁷² Casey JA, Savitz DA, Rasmussen SG et al. Unconventional Natural Gas Development and Birth Outcomes in Pennsylvania, USA. *Epidemiology* 2016; 27: 163-72. [Unconventional natural gas development and birth outcomes in Pennsylvania, USA - PMC](#)

⁷³ Casey JA, Karasek D, Ogburn EL et al. Retirements of Coal and Oil Power Plants in California: Association With Reduced Preterm Birth Among Populations Nearby. *Am J Epidemiol* 2018; 187: 1586-1594. [Retirements of Coal and Oil Power Plants in California: Association With Reduced Preterm Birth Among Populations Nearby - PMC](#)

⁷⁴ Lao XQ, Bo Y, Chen D, Zhang K and Szeto CC. Environmental pollution to kidney disease: an updated review of current knowledge and future directions. *Kidney Int*. 2024;106:214-225. doi: [10.1016/j.kint.2024.04.021](#). Epub 2024 May 24. PMID: 38797324.

⁷⁵ Bowe B, Xie Y, Li T, Yan Y, Xian H and Al-Aly Z. Estimates of the 2016 global burden of kidney disease attributable to ambient fine particulate matter air pollution. *BMJ Open*. 2019;9:e022450. doi: [10.1136/bmjopen-2018-022450](#). PMID: 31072847; PMCID: PMC6528010.

⁷⁶ Rajagopalan S, Brook RD, Salerno P, Bourges-Sevenier B, Landrigan P, Nieuwenhuijsen MJ, Munzel T, Deo SV and Al-Kindi S. Air pollution exposure and cardiometabolic risk. *Lancet Diabetes Endocrinol*. 2024;12:196-208. doi: [10.1016/S2213-8587\(23\)00361-3](#). Epub 2024 Feb 1. PMID: 38310921; PMCID: PMC11264310.

⁷⁷ GBD 2019 Diabetes and Air Pollution Collaborators. Estimates, trends, and drivers of the global burden of type 2 diabetes attributable to PM(2.5) air pollution, 1990-2019: an analysis of data from the Global Burden of Disease Study 2019. *Lancet Planet Health*. 2022;6:e586-e600. doi: [10.1016/S2542-5196\(22\)00122-X](#). PMID: 35809588; PMCID: PMC9278144.

⁷⁸ Lao XQ, Bo Y, Chen D, Zhang K and Szeto CC. Environmental pollution to kidney disease: an updated review of current knowledge and future directions. *Kidney Int*. 2024;106:214-225. doi: [10.1016/j.kint.2024.04.021](#). Epub 2024 May 24. PMID: 38797324.

⁷⁹ Bowe B, Xie Y, Li T, Yan Y, Xian H and Al-Aly Z. Estimates of the 2016 global burden of kidney disease attributable to ambient fine particulate matter air pollution. *BMJ Open*. 2019;9:e022450. doi: [10.1136/bmjopen-2018-022450](#). PMID: 31072847; PMCID: PMC6528010.

⁸⁰ Rajagopalan S, Brook RD and Munzel T. Environmental Hypertensionology and the Mosaic Theory of Hypertension. *Hypertension*. 2025;82:561-572. doi: [10.1161/HYPERTENSIONAHA.124.18733](#). Epub 2025 Feb 19. PMID: 39968647; PMCID: PMC11975430.

⁸¹ Bi J, Barry V, Weil EJ, Chang HH and Ebelt S. Short-term exposure to fine particulate air pollution and emergency department visits for kidney diseases in the Atlanta metropolitan area. *Environ Epidemiol*. 2021 Aug 6;5(4):e164. doi: [10.1097/EE.000000000000164](#). PMID: 34414347; PMCID: PMC8367053.

⁸² Okoye OC, Carnegie E and Mora L. Air Pollution and Chronic Kidney Disease Risk in Oil and Gas- Situated Communities: A Systematic Review and Meta-Analysis. *Int J Public Health*. 2022 Apr 11;67:1604522. doi: [10.3389/ijph.2022.1604522](#). PMID: 35479765; PMCID: PMC9035494.

⁸³ Luo C, Wei T, Joang W et al. The association between air pollution and obesity: an umbrella review of meta-analyses and systematic reviews. *BMC Public Health* 24, 1856 (2024). <https://doi.org/10.1186/s12889-024-19370-4>

⁸⁴ Yang B-Y , Fan S, Thiering E et al. Ambient air pollution and diabetes: A systematic review and meta-analysis. *Environ Res* 2020 Jan;180:108817. doi: [10.1016/j.envres.2019.108817](#). Epub 2019 Oct 12. PMID: 31627156.

⁸⁵ Chen H, Burnett RT, Kwong JC, Villeneuve PJ, Goldberg MS, Brook RD, van Donkelaar A, Jerrett M, Martin RV, Kopp A, Brook JR, Copes R. [Spatial association between ambient fine particulate matter and incident hypertension](#). *Circulation*. 2014;129:562-9.

⁸⁶ Bekkar B, Pacheco S, Basu R et al. Association of Air Pollution and Heat Exposure With Preterm Birth, Low Birth Weight, and Stillbirth in the USA Systematic Review. *JAMA Network Open* 2020; 3: e208243. doi: [10.1001/jamanetworkopen.2020.8243](#)

⁸⁷ Yu H, Yin Y, Zhang J, Zhou R. The impact of particulate matter 2.5 on the risk of preeclampsia: an updated systematic review and meta-analysis. *Environ Sci Pollut Res Int* 2020; 27: 37527-37539. doi: [10.1007/s11356-020-10112-8](#). Epub 2020 Aug 1. PMID: 32740838; PMCID: PMC7496023.



Fueling Sickness: The Hidden Health Costs of Fossil Fuel Pollution

⁸⁸ Li H, Cai J, Chen R et al. Particulate Matter Exposure and Stress Hormone Levels: A Randomized, Double-Blind, Crossover Trial of Air Purification. *Circ* 2017; 136: <https://doi.org/10.1161/CIRCULATIONAHA.116.026796>

⁸⁹ Kassotis CD, Tillitt DE, Davis JW, Hormann AM, Nagel SC. 2014. Estrogen and androgen receptor activities of hydraulic fracturing chemicals and surface and ground water in a drilling-dense region. *Endocrinology* 155:897-907 <https://pubmed.ncbi.nlm.nih.gov/24424034/>

⁹⁰ Ragusa A, Svelato A, Santacroce C et al. Plasticenta: First evidence of microplastics in human placenta. *Environ Int* 2021; 146: 10624. doi: [10.1016/j.envint.2020.106274](https://doi.org/10.1016/j.envint.2020.106274).

⁹¹ National Cancer Institute. Cancer Stat Facts: Common Cancer Sites. [Common Cancer Sites — Cancer Stat Facts](#)

⁹² Hiatt, R.A. and Beyeler, N. (2023), Women's cancers and climate change. *Int J Gynecol Obstet*, 160: 374-377. <https://doi.org/10.1002/ijgo.14407>

⁹³ American Cancer Society. Diesel Exhaust and Cancer Risk. February 2024. [Diesel Exhaust and Cancer Risk | American Cancer Society](#)

⁹⁴ Hamra GB, Guha N, Cohen A, et al. Outdoor particulate matter exposure and lung cancer: a systematic review and meta-analysis. *Environ Health Perspect*. 2014;122:906-11. doi: [10.1289/ehp.1408092](https://doi.org/10.1289/ehp.1408092)

⁹⁵ Turner MC, Andersen ZJ, Baccarelli A, et al: Outdoor air pollution and cancer: An overview of the current evidence and public health recommendations. *CA Cancer J Clin*, 2020. Aug 25:10.3322/caac.21632. doi: [10.3322/caac.21632](https://doi.org/10.3322/caac.21632). Epub ahead of print. PMID: 32964460; PMCID: PMC7904962.

⁹⁶ American Lung Association. State of Lung Cancer. Racial and Ethnic Disparities. 2024. [State of Lung Cancer | Racial and Ethnic Disparities | American Lung Association](#)

⁹⁷ White AJ, Fisher JA, Sweeney MR, et al: Ambient fine particulate matter and breast cancer incidence in a large prospective US cohort. *J Natl Cancer Inst* 116:53-60, 2024; Praud D, Deygas F, Amadou A, et al: Traffic-Related Air Pollution and Breast Cancer Risk: A Systematic Review and Meta-Analysis of Observational Studies. *Cancers (Basel)* 15, 2023; Ramamoorthy T, Nath A, Singh S, et al: Assessing the Global Impact of Ambient Air Pollution on Cancer Incidence and Mortality: A Comprehensive Meta-Analysis. *JCO Glob Oncol* 10:e2300427, 2024 [Air pollution and its impact on cancer incidence, cancer care and cancer outcomes - PMC](#)

⁹⁹ Boothe VL, Boehmer TK, Wendel AM, Yip FY. Residential traffic exposure and childhood leukemia: a systematic review and meta-analysis. *Am J Prev Med*. 2014 Apr;46(4):413-22. doi: 10.1016/j.amepre.2013.11.004. PMID: 24650845; PMCID: PMC5779082. [Residential Traffic Exposure and Childhood Leukemia: A Systematic Review and Meta-analysis - PMC](#)

¹⁰⁰ Clark CJ, Johnson NP, Soriano M, Jr., et al: Unconventional Oil and Gas Development Exposure and Risk of Childhood Acute Lymphoblastic Leukemia: A Case-Control Study in Pennsylvania, 2009-2017. *Environ Health Perspect* 130:87001, 2022. <https://doi.org/10.1289/EHP11092>

¹⁰¹ De Guzman R, Schiller J. Air pollution and its impact on cancer incidence, cancer care and cancer outcomes. *BMJ Oncol*. 2025 Mar 25;4(1):e000535. doi: 10.1136/bmjonc-2024-000535. PMID: 40165831; PMCID: PMC11956401. [Air pollution and its impact on cancer incidence, cancer care and cancer outcomes - PMC](#)

¹⁰² NPR. How Big Oil Misled the Public Into Believing Plastic Would be Recycled. September 2020. [Is Plastic Recycling A Lie? Oil Companies Touted Recycling To Sell More Plastic : NPR](#)

¹⁰³ Trasande L. The role of plastics in allergy, immunology, and human health: What the clinician needs to know and can do about it. *Ann Allergy Asthma Immunol*. 2025 Jan;134(1):46-52. doi: 10.1016/j.anai.2024.06.029. Epub 2024 Jun 28. PMID: 38945394. [The role of plastics in allergy, immunology, and human health: What the clinician needs to know and can do about it - PubMed](#)



Fueling Sickness: The Hidden Health Costs of Fossil Fuel Pollution

¹⁰⁴ Landrigan PJ, Raps H, Cropper M, Bald C, Brunner M, Canonizado EM, Charles D, Chiles TC, Donohue MJ, Enck J, Fenichel P, Fleming LE, Ferrier-Pages C, Fordham R, Gozt A, Griffin C, Hahn ME, Haryanto B, Hixson R, Ianelli H, James BD, Kumar P, Laborde A, Law KL, Martin K, Mu J, Mulders Y, Mustapha A, Niu J, Pahl S, Park Y, Pedrotti ML, Pitt JA, Ruchirawat M, Seewoo BJ, Spring M, Stegeman JJ, Suk W, Symeonides C, Takada H, Thompson RC, Vicini A, Wang Z, Whitman E, Wirth D, Wolff M, You-suf AK, Dunlop S. The Minderoo-Monaco Commission on Plastics and Human Health. *Ann Glob Health*. 2023 Mar 21;89(1):23. doi: 10.5334/aogh.4056. Erratum in: *Ann Glob Health*. 2023 Oct 11;89(1):71. doi: 10.5334/aogh.4331. PMID: 36969097; PMCID: PMC10038118. [The Minderoo-Monaco Commission on Plastics and Human Health - PubMed](#)

¹⁰⁵ Yale Climate Connections. What are the health impacts of plastic? May 2025. [What are the health impacts of plastic? » Yale Climate Connections](#)

¹⁰⁶ Baj J, Dring JC, Czeczelewski M, et al: Derivatives of Plastics as Potential Carcinogenic Factors: The Current State of Knowledge. *Cancers (Basel)* 14, 2022. doi: [10.3390/cancers14194637](https://doi.org/10.3390/cancers14194637). PMID: 36230560; PMCID: PMC9562888.

¹⁰⁷ MorelloFrosch, R., & Obasogie, O. K. (2023). The climate gap and the color line — Racial health inequities and climate change. *New England Journal of Medicine*, 388(10), 943949. <https://doi.org/10.1056/NEJMsb2213250>

¹⁰⁸ Timothy Q. Donaghy, Noel Healy, Charles Y. Jiang, Colette Pichon Battle, Fossil fuel racism in the United States: How phasing out coal, oil, and gas can protect communities, *Energy Research & Social Science*, Volume 100, 2023, 103104, ISSN 2214-6296, <https://doi.org/10.1016/j.erss.2023.103104>

¹⁰⁹ NAACP and Clean Air Task Force. Fumes Across the Fence-Line. The Health Impacts of Air Pollution from Oil & Gas Facilities on African American Communities. [Fumes Across the Fence-Line: The Health Impacts of Air Pollution from Oil & Gas Facilities on African American Communities | NAACP](#)

¹¹⁰ Jacob L. Wasserman, Anastasia Loukaitou-Sideris, Hao Ding, Claire Nelischer, The Road, Home: Challenges of and Responses to Homelessness in State Transportation Environments, *Transportation Research Interdisciplinary Perspectives*, Volume 21, 2023, 100890, ISSN 2590-1982, <https://doi.org/10.1016/j.trip.2023.100890>.

¹¹¹ American Lung Association. Who is Most Affected by Outdoor Air Pollution? September 2024. [Who is Most Affected by Outdoor Air Pollution? | American Lung Association](#)

¹¹² CDC. About Homelessness and Health. October 2024. [About Homelessness and Health | Homelessness and Health | CDC](#)

¹¹³ Yeoman K, Sussell A, Retzer K, Poplin G. Health Risk Factors Among Miners, Oil and Gas Extraction Workers, Other Manual Labor Workers, and Nonmanual Labor Workers, BRFSS 2013-2017, 32 States. *Workplace Health Saf*. 2020 Aug;68(8):391-401. doi: 10.1177/2165079920909136. Epub 2020 Jun 2. PMID: 32484732; PMCID: PMC8340646. [Health risk factors among miners, oil and gas extraction workers, other manual labor workers, and non-manual labor workers, BRFSS 2013–2017, 32 States – PMC](#)

¹¹⁴ Occupational Safety and Health Administration. Safety Hazards Associated with Oil and Gas Extraction Activities. [Oil and Gas Extraction - Hazards | Occupational Safety and Health Administration](#)

¹¹⁵ Yale School of Medicine. Yale environmental health expert discusses concerns about fracking. November 2024. [Yale environmental health expert discusses concerns about fracking < Yale School of Medicine](#)

¹¹⁶ Perera, Frederica & Nadeau, Kari. (2022). Climate Change, Fossil-Fuel Pollution, and Children's Health. *New England Journal of Medicine*. 386. 2303-2314. 10.1056/NEJMra2117706. [Climate Change, Fossil-Fuel Pollution, and Children's Health](#)

¹¹⁷ Perera, Frederica, and Kari Nadeau. "Climate Change, Fossil-Fuel Pollution, and Children's Health." *New England Journal of Medicine*, vol. 386, no. 24, 16 June 2022, pp. 2303–2314. *New England Journal of Medicine*, doi:10.1056/NEJMra2117706. [Climate Change, Fossil-Fuel Pollution, and Children's Health | New England Journal of Medicine](#)

¹¹⁸ Yale School of Medicine. Yale environmental health expert discusses concerns about fracking. November 2024. [Yale environmental health expert discusses concerns about fracking < Yale School of Medicine](#)

¹¹⁹ Yale School of the Environment. YaleEnvironment360. As Evidence Mounts, New Concerns About Fracking and Health. November 2022. [As Evidence Mounts, New Concerns About Fracking and Health - Yale E360](#)

¹²⁰ American Lung Association. State of the Air. Health Impact of Air Pollution. April 2025. [Health Impact of Pollution | State of the Air | American Lung Association](#)



Fueling Sickness: The Hidden Health Costs of Fossil Fuel Pollution

¹²¹ JeongWon Han, Bangning Zhang, Xiao Zhang, Ke Huang, Vixey Fang, Xiaohui Xu, Associations between occurrence of birth defects and hydraulic fracturing activities in Barnett shale region, Texas, *Helijon*, Volume 9, Issue 4, 2023, e15213, ISSN 2405-8440, <https://doi.org/10.1016/j.heliyon.2023.e15213>.

¹²² Johns Hopkins Bloomberg School of Public Health. Study: Fracking Industry Wells Associated With Premature Birth. October 2015. [Study: Fracking Industry Wells Associated With Premature Birth | Johns Hopkins | Bloomberg School of Public Health](#)

¹²³ Lancet Countdown on Health and Climate Change. Policy Brief for the United States of America. 2024. [2024 Lancet Countdown U.S. Brief - LANCET COUNTDOWN](#):

¹²⁴ Khatana SAM, Szeto JJ, Eberly LA, Nathan AS, Puvvula J, Chen A. Projections of Extreme Temperature–Related Deaths in the US. *JAMA Netw Open*. 2024;7(9):e2434942. doi: [10.1001/jamanetworkopen.2024.34942](https://doi.org/10.1001/jamanetworkopen.2024.34942)

¹²⁵ Heanoy EZ, Brown NR. Impact of Natural Disasters on Mental Health: Evidence and Implications. *Healthcare (Basel)*. 2024 Sep 10;12(18):1812. doi: [10.3390/healthcare12181812](https://doi.org/10.3390/healthcare12181812). PMID: 39337153; PMCID: PMC11430943.

¹²⁶ White NJ, Warrell DA, Chanthavanich P et al. Severe Hypoglycemia and Hyperinsulinemia in Falciparum Malaria. *N Engl J Med*. 1983 Jul 14;309(2):61-6. doi: [10.1056/NEJM198307143090201](https://doi.org/10.1056/NEJM198307143090201). PMID: 6343877.

¹²⁷ Long-term exposure to wildfires and cancer incidence in Canada: a population-based observational cohort study. Korsiak, Jill et al. *The Lancet Planetary Health*, Volume 6, Issue 5, e400 – e409 [Long-term exposure to wildfires and cancer incidence in Canada: a population-based observational cohort study - The Lancet Planetary Health](#)

¹²⁸ Bell, S.A., Banerjee, M., Griggs, J.J. et al. The Effect of Exposure to Disaster on Cancer Survival. *J GEN INTERN MED* **35**, 380–382 (2020). <https://doi.org/10.1007/s11606-019-05465-x>

¹²⁹ Thomas Münzel, Haitham Khraishah, Alexandra Schneider, Jos Lelieveld, Andreas Daiber, Sanjay Rajagopalan, Challenges posed by climate hazards to cardiovascular health and cardiac intensive care: implications for mitigation and adaptation, *European Heart Journal. Acute Cardiovascular Care*, Volume 13, Issue 10, October 2024, Pages 731–744, <https://doi.org/10.1093/ehjacc/zuae113>

¹³⁰ Hassan AM, Nogueira L, Lin YL, Rogers JE, Nori-Sarma A, Offodile AC 2nd. Impact of Heatwaves on Cancer Care Delivery: Potential Mechanisms, Health Equity Concerns, and Adaptation Strategies. *J Clin Oncol*. 2023 Jun 10;41(17):3104-3109. doi: 10.1200/JCO.22.01951. Epub 2023 Apr 25. PMID: 37098249. [Impact of Heatwaves on Cancer Care Delivery: Potential Mechanisms, Health Equity Concerns, and Adaptation Strategies - PubMed](#)



1930 18th St NW Ste B2 PMB 2338, Washington, DC 20009 | psr.org

Published November 2025

CLIMATE AND HEALTH TASK FORCE




WASHINGTON PHYSICIANS
FOR SOCIAL RESPONSIBILITY

WHAT IS THE TASK FORCE?

WPSR's climate and health task force - comprised primarily of healthcare professionals - seeks to bring attention to climate change as a serious individual and public health issue. We enlist healthcare leaders to speak out on the health impacts of the climate crisis using their trusted medical voices to promote policies that protect our climate and our health.

TASK FORCE MEMBERS IN ACTION



OUR CLIMATE & HEALTH PRIORITIES



ELECTRIFICATION



FOSSIL FUELS



TRANSPORTATION

WHAT CAN I WORK ON?

WPSR Task Force members meet monthly to discuss challenges and opportunities, to learn, and to plan advocacy efforts on the policies that support our mission to foster a healthy climate.

Task Force members attend coalition meetings, write letters to the editor and op-eds, meet with elected officials, deliver trainings, and speak publicly in our effort to inform and influence decision-makers of the health impacts of climate policies.

Task force members engage as their skills, expertise, and time allow. There is a place for all regardless of prior knowledge or expertise.

LEARN MORE & JOIN

WPSR membership is open to all, and you can sign up as a member and to receive action alert emails on our website at www.wpsr.org. To engage with WPSR's Climate & Health program, please email wpsr@wpsr.org!



Use this QR Code
to sign up for
WPSR Climate &
Health Action
Alerts.

Contact: James Moschella, Climate and Health Program Manager, james@wpsr.org