

VACCINE	Birth	1 Month	2 months	4 months	6 months	8 months	9 months	12 months	15 months	18 months	19-23 months	2-3 years	4-6 years	7-10 years	11-12 years	13-15 years	16 years	17-18 years	
Respiratory syncytial virus (RSV-mAb [nirsevimab, clesrovimab])	1 dose depending on maternal RSV vaccination status (<8 months)					1 dose (8–19 months)													
Hepatitis B (HepB)	1 <sup>st</sup> dose	2 <sup>nd</sup> dose		3 <sup>rd</sup> dose															
Rotavirus (RV): RV1 (2-dose series), RV5 (3-dose series)			1 <sup>st</sup> dose	2 <sup>nd</sup> dose	3 <sup>rd</sup> dose*														
Diphtheria, tetanus, and acellular pertussis (DTaP <7 yrs)			1 <sup>st</sup> dose	2 <sup>nd</sup> dose	3 <sup>rd</sup> dose	4 <sup>th</sup> dose			5 <sup>th</sup> dose										
Haemophilus influenzae type b (Hib)			1 <sup>st</sup> dose	2 <sup>nd</sup> dose	3 <sup>rd</sup> dose	4 <sup>th</sup> dose													
Pneumococcal conjugate (PCV15, PCV20)			1 <sup>st</sup> dose	2 <sup>nd</sup> dose	3 <sup>rd</sup> dose	4 <sup>th</sup> dose													
Inactivated poliovirus (IPV)			1 <sup>st</sup> dose	2 <sup>nd</sup> dose	3 <sup>rd</sup> dose					4 <sup>th</sup> dose									
COVID-19 (1vCOV-mRNA, 1vCOV-aPS)			1 or 2 doses**					Yearly based on medical necessity											
Influenza			1 or 2 doses annually***											1 dose annually (9 years and above)					
Measles, mumps, and rubella (MMR)							1 <sup>st</sup> dose						2 <sup>nd</sup> dose						

VACCINE	Birth	1 Month	2 months	4 months	6 months	8 months	9 months	12 months	15 months	18 months	19-23 months	2-3 years	4-6 years	7-10 years	11-12 years	13-15 years	16 years	17-18 years
Varicella (VAR)								1 <sup>st</sup> dose					2 <sup>nd</sup> dose					
Hepatitis A (HepA)								2 dose series										
Tetanus, diphtheria, and acellular pertussis (Tdap ≥7 yrs)															1 dose			
Human papillomavirus (HPV)															2 dose series			
Meningococcal ACWY															1 <sup>st</sup> dose		2 <sup>nd</sup> dose	
Meningococcal B																	1 dose	

This routine pediatric vaccine schedule is reflective of evidence-based guidelines around the recommended standard timing of child and adolescent vaccines. For further information on vaccine interval recommendations, as well as additional tables listing timings for catch-up vaccines and guidelines around specific medical indications [and contraindications](#)<sup>1</sup>, please refer to the [most current](#) immunization tables which can be found at: <https://downloads.aap.org/AAP/PDF/AAP-Immunization-Schedule.pdf>

[For adult vaccine coverage, refer to AMPM Policy 310-M.](#)<sup>2</sup>

~~For pediatric vaccine recommendations: Refer to Table 1 – Recommended Child and Adolescent Immunization Schedule for Ages 18 Years or Younger, United States, 2025.~~

<sup>1</sup> Unduplicated language referring to the immunization schedule tables.

<sup>2</sup> Added reference to AMPM Policy 310-M for adult vaccine coverage.

~~For catch-up vaccines: Refer to Table 2 – Recommended Catch-up Immunization Schedule for Children and Adolescents Who Start Late or Who Are More than 1 Month Behind, United States, 2025~~

~~For specific medical indications and contraindications: Refer to Table 3: Recommended Child and Adolescent Immunization Schedule by Medical Indication, United States, 2025~~

\* Rotavirus vaccination is a 2 or 3 dose series depending on the vaccine manufacturer. If any dose in the series is of unknown type, default to 3-dose series.

\*\* 1-2 doses depending on COVID-19 vaccine manufacturer guidelines. See [American Academy of Pediatrics \(AAP\)](#) vaccine schedule for further guidance.

\*\*\* For ages 6 months to 8 years 2 doses of influenza are needed for the first year of vaccination only.